arXiv:cond-mat/9701012v1 3 Jan 1997

Fermi Systems with Strong Forward Scattering

Walter Metzner
Sektion Physik, Universität München
Theresienstraße 37, D-80333 München, Germany

Claudio Castellani and Carlo Di Castro
Dipartimento di Fisica, Università "La Sapienza",
P.le A. Moro 2, 00185 Roma, Italy

February 1, 2008

Abstract

We review the theory of interacting Fermi systems whose low-energy physics is dominated by forward scattering, i.e. scattering processes generated by effective interactions with small momentum transfers. These systems include Fermi liquids as well as several important non-Fermi liquid phases: one-dimensional Luttinger liquids, systems with long-range interactions, and fermions coupled to a gauge field. We report results for the critical dimensions separating different "universality classes", and discuss the behavior of physical quantities such as the momentum distribution function, the single-particle propagator and low-energy response functions in each class.

The renormalization group for Fermi systems will be reviewed and applied as a link between microscopic models and effective low-energy theories. Particular attention is payed to conservation laws, which constrain any effective low-energy theory of interacting Fermi systems. In scattering processes with small momentum transfers the velocity of each scattering particle is (almost) conserved. This asymptotic conservation law leads to non-trivial cancellations of Feynman diagrams and other simplifications, making thus possible a non-perturbative treatment of forward scattering via Ward identities or bosonization techniques.

PACS: 71.10.-w, 71.10.Ay, 71.10.Hf, 71.10.Pm

CONTENTS

1. Introduction **1**

1.1. Normal phases of interacting fermions 1

1.2. Theoretical routes to non-Fermi liquid behavior 3

1.3. Dominant forward scattering 7

1.4. Contents of this article 10

2. Renormalization group **12**

2.1. Functional integral representation 13

2.2. Mode elimination and effective actions 15

2.3. Low-energy coupling-space 21

3. Exact conservation laws **26**

3.1. Charge conservation 26

3.2. Spin conservation 29

3.3. Renormalization of response functions 31

4. Fermi liquid **34**

4.1. Microscopic Fermi liquid theory 34

4.2. Effective action and fixed point 36

4.3. Response functions 39

4.4. Subleading corrections 40

4.5. Instabilities 41

5. Forward scattering **43**

5.1. Global charge and spin conservation 45

5.2. Velocity conservation 47
5.2.1. Loop cancellation and response functions 48
5.2.2. Density and current vertex 51
5.2.3. Single-particle propagator 52
5.3. Bosonization 56

6. One-dimensional Luttinger liquid **63**

6.1. The g-ology model 64
6.2. Renormalization group and cutoffs 66
6.3. Charge/spin conservation and Ward identities 68
6.3.1. Total charge and spin conservation 68
6.3.2. Separate left/right conservation laws 70
6.4. Luttinger liquid fixed point 71
6.5. Instabilities 77

7. Short-range interactions in d dimensions **79**

7.1. Continuation to non-integer dimensions 80
7.2. Perturbative results 81
7.2.1. Particle-hole bubble 81
7.2.2. Particle-particle bubble 83
7.2.3. Second order self-energy 84
7.3. Random phase approximation 87
7.3.1. Effective interaction 87
7.3.2. Density-density response 88
7.3.3. RPA self-energy 88
7.4. Resummation of forward scattering 90
7.5. Critical dimensions and crossover 94

7.5.1. Leading low-energy behavior 94
7.5.2. Subleading corrections 96

8. Long-range density-density interactions **98**

9. Fermions coupled to a gauge-field **103**

9.1. Action 103
9.2. Gauge-field propagator 105
9.3. Fermion propagator 108
9.4. Response functions 114

10. Conclusions **116**

Acknowledgements **122**

Appendices **123**

A: Loop cancellation for $N = 3$ 123
B: Spectral representation of $\bar{D}$ 123
C: Bubbles for a quadratic dispersion relation 125
D: Perturbative self-energy in one dimension 127
E: Limit $d \to 1$ for the propagator 127

References **129**

Figure captions **137**

1. INTRODUCTION

1.1. NORMAL PHASES OF INTERACTING FERMIONS

Observed phases of interacting Fermi systems such as liquid ^{3}He and electrons in metals fall into two broad categories: symmetry-broken and "*normal*". The former contains superconductors or superfluids, magnetically ordered phases, charge density waves, and several others. While the variety of ordered phases has long been known to be very rich, the low-energy behavior of normal Fermi systems seemed to be governed by a remarkably small number of "universality classes".

Actually most normal metals, as well as normal ^{3}He, follow the same type of low-energy behavior, the one described by *Fermi liquid theory* [1, 2]. This originally purely phenomenological theory makes very specific predictions on the low-energy scaling (i.e. power-law) behavior of thermodynamic and transport properties as a function of temperature, frequency, and other small energy or momentum scales. These predictions have been experimentally verified to an extremely high precision in liquid ^{3}He, a rather exceptional substance in its high purity and its perfect isotropy [3]. In metals, the anisotropy imposed by the crystal structure and the unavoidable impurities blur some of the more sensitive properties of pure and isotropic Fermi liquids, but many others are robust with respect to such complications. Among these robust properties are the existence of a Fermi surface, a specific heat proportional to temperature, a constant spin-susceptibility, a finite density of low-energy single-particle excitations and, of course, a metallic DC-conductivity which increases for decreasing temperature, implying that charged excitations are gapless. These properties behave qualitatively as in a non-interacting Fermi gas. In Fermi liquid theory they are phenomenologically, but quantitatively, described in terms of Landau's "*quasi-particles*", i.e. fermionic elementary excitations which have momenta near the Fermi surface and obey a free-fermion-like energy-momentum relation. To leading order (in energy scales), the quasi-particles are mutually independent, except for a Hartree-type interaction. The *parameters* characterizing these almost-free excitations vary drastically from one substance to another. In particular, the effective mass of quasi-particles in metals may be as much as a factor thousand bigger than the bare electron mass, as has been observed in the so-called heavy fermion systems [4]. In addition, the energy or temperature *scale* controlled by Fermi liquid theory is strongly material dependent, too. Usually at least three distinct scales must be distinguished: the Fermi temperature T_F given

roughly by the average kinetic energy, a coherence temperature T^* below which Fermi liquid behavior sets in, and a transition temperature T_c associated with an instability of the normal Fermi liquid towards some sort of symmetry breaking. For electrons in metals, T_F is usually much higher than room temperature. Phase transition temperatures may be anything from zero to $1388\,K$, the Curie temperature of cobalt. Both T_c and T^* are always much smaller than T_F. Usually T_c is also much smaller than T^*, but sometimes they are comparable, such that only very gross aspects of Fermi liquid behavior can be really observed.

Going to small but finite energy scales, there are subleading corrections due to quasi-particle scattering. Among these are a quasi-particle decay rate proportional to the square of its energy, a contribution proportional to $T^3 \log T$ to the specific heat (in three dimensions) and a contribution of order T^2 to the electrical resistivity due to electron-electron interactions in metals. In metals, subleading terms due to electron-electron scattering cannot always be observed experimentally, because they are often superposed by other, much larger, contributions due to phonons or disorder.

Up to recently, Fermi liquid theory seemed universally applicable at least to all sufficiently pure interacting Fermi systems, and its more gross features even to quite dirty ones, provided the normal phase is not destroyed by symmetry breaking before the coherence temperature T^* is reached. This situation has changed during the last years with the discovery of new materials, where unusual scaling behavior is observed above T_c, which differs in many respects from the predictions of Fermi liquid theory.

High temperature superconductors are certainly the most famous among these new materials. Their unusual and unexpected properties have stimulated a profound reinvestigation of basic concepts in the theory of metals. The high superconducting transition temperature is hard to obtain from the conventional phonon-induced pairing, and the coherence length is too short for applying simple BCS mean field theory, whatever the pairing mechanism may be. Even more striking is the behavior of the normal metallic phase in these materials, which does not fit in a simple Fermi liquid type description. Although photoemission experiments yield convincing evidence for the existence of a large Fermi surface, many transport properties are incompatible with a Fermi liquid picture of low-lying excitations [5, 6]. The most prominent among many other quantitities showing non-Fermi-liquid behavior is the electrical resistivity, whose temperature dependence deviates significantly from the T^2 behavior at low T predicted by Fermi liquid theory. The layer structure and the giant anisotropy observed in various properties like the resistivity indicate that the normal phase is governed by electrons confined essentially to two dimensions. The vicinity to an antiferromagnetic Mott insulating phase reveals the

importance of electron-electron interactions. These observations have stimulated speculations on the existence of non-Fermi-liquid metallic phases in two-dimensional interacting electron systems [7, 8].

Asymptotic low-energy behavior different from Fermi liquid predictions in a normal metallic phase has also been found recently in various heavy-fermion alloys [9, 10, 11, 12, 13]. A specific heat proportional to $T \log T$ and a T-linear contribution to the electrical resistivity was observed. The magnetic susceptibility as a function of temperature has a cusp in one of these materials, and an algebraic divergence for $T \to 0$ in another.

1.2. THEORETICAL ROUTES TO NON-FERMI LIQUID BEHAVIOR

Theoretically, the almost universal validity of Fermi liquid theory in normal Fermi systems has been made plausible in various ways. A justification within perturbation theory has already been given by Landau and his coworkers [14]. To push the validity of the theory beyond the perturbatively accessible regime in coupling space, only few basic assumptions on the asymptotic low-energy behavior of propagators and vertex functions need to be made [2, 14].

A new point of view has recently been developed with the adaption of Wilson's *renormalization group* to interacting Fermi systems [15, 16, 17, 18, 19]. The idea is to integrate out high energy degrees of freedom in a path-integral representation of the full interacting theory. At least in perturbation theory this corresponds to integrating out fermions with momenta far from the Fermi surface. Successive reduction of the momentum cutoff generates a sequence of effective actions with renormalized kinetic terms and interactions. One may then classify different types of low-energy behavior in terms of different asymptotic effective actions, defined on a thin shell around the Fermi surface of the interacting system.[1] Consistency of a certain hypothetical low-energy action can be systematically checked by calculating the renormalization group flow of its couplings. The renormalization group thus provides effective actions representative of the low-energy behavior of the system. These asymptotic low-energy theories are prototype models of possible (metallic) phases, which can often be solved exactly by various techniques.

An effective low-energy action with finite (not singular) renormalized interactions will usually lead to Fermi liquid behavior, with the possibility of a Cooper instability towards superconductivity. Different low-energy behavior may be found if the effective theory has

[1] It is important to give a sufficiently general definition of a *Fermi surface*. In a Fermi liquid, the momentum distribution function $n_{\mathbf{k}}$ is discontinuous across the Fermi surface, and this property is often used to define the surface itself. There are other systems where $n_{\mathbf{k}}$ is continuous everywhere, but where one can still define a Fermi surface as a surface in momentum space where the excitations with lowest energy are located.

one of the following peculiar features:

a) dimensionality $d = 1$,
b) special Fermi surfaces,
c) vanishing Fermi velocities,
d) singular interactions,
e) coupling to gauge fields,
f) coupling to other soft modes,
g) local degrees of freedom.

A *normal* metallic phase of interacting fermions with a low-energy behavior that is not described by Fermi liquid theory is generally called a "*non-Fermi liquid*". "Metallic" behavior implies that there are gapless charge fluctuations. More generally, including neutral fermions such as ^{3}He, one would require the existence of gapless density fluctuations. Let us now address the various routes to non-Fermi liquid behavior one by one.

a) The breakdown of Fermi liquid theory in *one-dimensional* interacting Fermi systems shows up already in second order perturbation theory: the perturbative contributions to the quasi-particle weight diverge logarithmically at the Fermi surface of the non-interacting system. The problem of treating these divergencies has been solved by the renormalization group, using an effective low-energy theory known as "g-ology" model [20]. Assuming a scaling ansatz for the vertex functions, one approaches the Fermi surface by rescaling the fields and the coupling constants (a small number of "g's"). Depending on the values of the bare couplings the renormalized couplings flow either to strong coupling, and hence out of the perturbatively controlled regime, or to the Luttinger model [21], which is exactly solvable [22] . In the latter case the system is a "*Luttinger liquid*", i.e. a normal (not symmetry-broken) metallic phase characterized by i) a continuous momentum distribution with a power-law singularity at the Fermi surface, the exponent η being non-universal; ii) a single-particle density of states which vanishes as ω^η near the Fermi energy, implying the absence of fermionic quasi-particles; iii) finite charge and spin density response for small wave vectors and the existence of collective bosonic charge and spin density modes; iv) power-law singularities in Cooper pair correlation functions and density correlation functions with large wave vectors; v) separation of spin and charge degrees of freedom [20, 23, 24]. As introduced by Haldane [25], the term "Luttinger liquid" denotes the universality class of systems whose fixed point Hamiltonian is the Luttinger model.

b) It is easy to see, in weak coupling analysis, that certain special shapes of Fermi surfaces make a Fermi liquid unstable. In particular "*nesting*", i.e. the property that a finite fraction of the Fermi surface can be shifted to other points of the surface by adding a certain

fixed momentum transfer, usually leads to formation of density waves and thus breaking of translational invariance. If a Fermi surface contains extended almost flat pieces, a whole host of competing infrared singularities appears in perturbation theory. The perturbative quasi-particle decay rate is then linear in temperature (or frequency), signalling a breakdown of Fermi liquid theory. To obtain the real low-energy behavior in such a case is a formidable task even at weak coupling, because the necessary renormalization group calculation is complicated by the huge coupling space one has to consider, and one must go at least to two-loop order to include singularities involving single-particle excitations. It has been suggested that the shape of Fermi surfaces in high-T_c superconductors might be responsible at least for some of their anomalous properties [26], but to our knowledge a controlled calculation that takes into account all the singular channels has not yet been performed. Hence it is not clear whether the Fermi surface shapes can quantitatively explain the strange low-energy behavior in these materials over a rather wide range of different compositions.

c) The Fermi velocity may vanish on or near parts of the Fermi surface. Saddle points of the dispersion relation in momentum space lead to *van Hove singularities* in the single-particle density of states. In three dimensions these singularities are only cusps, while in two dimensions logarithmic divergencies are obtained. If such a divergence comes close to the Fermi energy, it will affect the low-energy behavior: second order perturbation theory yields a quasi-particle decay rate that depends linearly on energy [27]. It has been proposed that this linear behavior can explain the anomalous transport in high-T_c superconductors [27, 28]. On the other hand, it has been pointed out that transport properties are related to decay rates for currents rather than single-particle decay, and, within perturbation theory, these decay rates relevant for transport properties are not significantly affected by the van Hove singularities [29]. In any case, the presence of singularities makes the application of standard perturbation theory problematic, and one has to wait for a more sophisticated analysis.

d) Anderson [7] has suggested that in a two-dimensional Fermi system perturbation theory and Fermi liquid theory should break down even at *weak* coupling, having in mind two intriguing signals: i) a finite phase shift for two interacting particles on the same point of the Fermi surface, and ii) an "antibound state" in the two-particle spectrum, separated by an energy gap from the continuum of all other states. He argued that conventional many body theory would miss these effects, while they might be included by introducing a *singular "pseudo-potential"* acting between electrons. The resulting state should not be a Fermi liquid, but rather a two-dimensional "Luttinger liquid"[2] with properties similar

[2] According to Anderson's terminology, the term "Luttinger liquid" is no more restricted to one-

to those known for one-dimensional systems. So far, nobody has succeeded to construct a sound theory based on these intuitive ideas. Several authors have analyzed the phase shift in considerable detail, but they all concluded that one cannot infer a breakdown of Fermi liquid theory from that effect [30, 31, 32, 33]. Recent rigorous results confirm the general expectation that the Cooper instability is the only weak coupling instability of generic two-dimensional Fermi liquids with short-range interactions [34]. It has also been shown that the mechanism leading to Luttinger liquid behavior in one dimension does not destroy the Landau quasi-particle in any (possibly non-integer) dimension $d > 1$ [35]. However, certain ad hoc *long-range* interactions (with singular Fourier transforms in momentum space) can indeed destabilize the Fermi liquid in dimensions $d > 1$ [36, 37, 38]. To overwhelm screening, which usually makes long-range interactions in an electron system effectively short-ranged, these interactions must be even more singular than the Coulomb interaction.

e) For lattice models of interacting fermions with a local constraint (arising from very strong repulsion) an effective low-energy theory involving fermions and bosons coupled to a ficticious *gauge field* has been derived [39, 40, 41]. The coupling to the gauge field leads to infrared divergences in perturbation theory. In an approximate evaluation of the theory non-Fermi liquid behavior in several transport properties has been obtained [40, 41]. A controlled derivation of the low-energy behavior is however very difficult, and has not yet been fully achieved. The real electromagnetic gauge field also leads to divergences when coupled to a many fermion system [42, 43]. The ensuing physical effects are however only of order v_F/c (where v_F is the Fermi velocity and c the velocity of light) and are thus practically unobservable. Fermions coupled to a gauge field may also describe interacting fermions in a strong magnetic field tuned such that the highest occupied Landau level is precisely half-filled. Halperin, Lee and Read [44] and Kalmeyer and Zhang [45] have indeed proposed that this system can be mapped on a system of spinless fermions in zero (average) magnetic field, but coupled to a fluctuating gauge field.

f) Anomalous scattering mediated by soft modes appears to be a generic consequence of proximity to a critical point. This mechanism has been invoked to explain various non-Fermi liquid features in heavy fermion systems close to a second-order phase transition [46, 47]. Proximity to a zero temperature antiferromagnetic and/or charge instability has been also suggested to be responsible for the anomalous metallic behavior in high-T_c cuprates.

Close to an antiferromagnetic instability, *antiferromagnetic spin fluctuations* lead to a

dimensional systems, but may also denote higher-dimensional non-Fermi liquids, provided they have a Fermi surface in the generalized sense described above.

strongly enhanced quasi-particle decay rate, pointing towards non-Fermi liquid behavior. It has been argued that the anomalous quasi-particle decay carries over to anomalous transport properties, such as those observed in high temperature superconductors [48, 49]. On the other hand, analyzing non-trivial solutions of the Boltzmann equation which take into account that spin fluctuations enhance scattering strongly only on special points of the Fermi surface ("hot spots"), Hlubina and Rice [29] have found that standard Fermi liquid behavior (for transport) holds up to a rather high crossover scale, which is too high to explain the observed behavior in the high-T_c materials.

Charge fluctuations also lead to strongly singular scattering and non-Fermi liquid behavior [50, 51, 52]. In particular, in the proximity of phase separation the effective interactions are singular at small $\mathbf{q}$ and lead to infrared singularities which share many similarities with those generated by coupling to a gauge field [51]. When the long-range Coulomb force ensures macroscopic charge neutrality an incommensurate charge density wave may instead occur at a finite wave vector $\mathbf{q}_{\rm CDW}$. Close to this instability the effective interaction becomes singular at $\mathbf{q}_{\rm CDW}$ giving rise to non-Fermi liquid contributions in the quasi-particle decay rate and in transport properties [51].

All the above results are mainly based on lowest order perturbation theory whose validity is questionable close to an instability. A consistent treatment of non-Fermi liquid properties produced by critical fluctuations in the proximity of an instability has not yet been achieved. It is however a formidable project which is worth carrying out.

g) Non-Fermi liquid behavior has been well-established in special single-impurity models, where a sea of conduction electrons is coupled to a *local degree of freedom*. The first instance was given by a two-channel Kondo model, where a localized spin is coupled to two degenerate channels of conduction electrons [53, 54, 55, 56, 57]. Since then many other single impurity models with non-Fermi liquid behavior have been discovered. A non-Fermi liquid phase has also been found in the infinite-dimensional Falikov-Kimball model, a (translation invariant) lattice model with two interacting species of electrons, one itinerant, the other localized [58]. In the Falikov-Kimball model localized and itinerant electrons do not hybridize, but non-Fermi liquid behavior has been reported also for a more general class of infinite-dimensional two-band models, where localized and itinerant degrees of freedom are connected by a hybridization term [59, 60].

1.3. DOMINANT FORWARD SCATTERING

Forward scattering, i.e. scattering with a *small momentum transfer* $\mathbf{q}$, plays a prominent role in Fermi liquids, as well as in several of the above-mentioned non-Fermi liquid systems: one-dimensional Luttinger liquids, systems with singular interactions, and fermions coupled to a gauge field. In all these systems forward scattering governs the

leading low-energy long-wavelength behavior of response functions, and in the non-Fermi liquid phases forward scattering drives the instability of Landau's quasi-particles. The breakdown of Fermi liquid theory in these systems is associated with divergences in perturbation theory, which require a suitable resummation of all orders in the coupling constant. Sometimes the divergencies can be treated by perturbative renormalization group methods, but in general a non-perturbative construction is necessary.

Fortunately, in forward-scattering-dominated systems there is a small parameter other than the coupling strength: the dimensionless ratio $|\mathbf{q}|/k_F$, where $\mathbf{q}$ is a typical momentum transfer in a scattering process, and k_F is the Fermi momentum (or, in anisotropic systems, an effective radius in momentum space measuring the local curvature of the Fermi surface). The smallness of this parameter can be exploited in several ways. To make things precise, it is useful to introduce a cutoff $q_c \ll k_F$, and to allow only for scattering processes with a momentum transfer $|\mathbf{q}| < q_c$. This allows one to consider a formal expansion with respect to the small parameter q_c/k_F. Apart from certain finite renormalizations of parameters, the asymptotic low-energy behavior of forward-scattering-dominated systems turns out to be actually independent of q_c, showing that scattering processes with $|\mathbf{q}| > q_c$ are ultimately irrelevant in the low-energy limit, however small q_c may be. In some cases even subleading corrections to the leading low-energy behavior are completely determined by forward scattering, but in general contributions with any generic $\mathbf{q}$ contribute to these corrections. For example, the decay rate of quasi-particles in systems with short-range interactions is dominated by forward scattering in dimensions $d < 2$ [35], while processes with any $\mathbf{q}$ contribute to the asymptotic decay rate in $d \geq 2$.

To leading order in q_c/k_F, the *velocity* of the particles is *conserved* in each scattering process. This leads to dramatic simplifications in diagrammatic perturbation theory, especially:
i) *"Loop cancellation"*: Fermionic loops containing more than two insertions (density, current or interaction type) cancel each other.
ii) *"Density-current relation"*: The irreducible current vertex $\mathbf{\Lambda}(p;q)$ is related to the irreducible density vertex $\Lambda^0(p;q)$ via

$$\mathbf{\Lambda}(p;q) \;\sim\; \mathbf{v_p}\,\Lambda^0(p;q)$$

for small momenta $\mathbf{q}$. Here $\mathbf{v_p}$ is the velocity of a fermion with momentum $\mathbf{p}$, and p, q are energy-momentum variables, e.g. $p = (p_0, \mathbf{p})$.

The loop-cancellation has been noticed long ago in the analysis of the one-dimensional Luttinger model, where it is exact for arbitrary q_c [61]. In higher dimensions this cancellation has been more or less implicit in many works on forward-scattering-dominated problems. An explicit proof has been presented recently by Kopietz et al. [62]; an alterna-

tive derivation obtained independently by us will be given in Sec. 5. The loop-cancellation directly implies that self-energy and vertex corrections cancel each other in polarization bubbles with small momenta, i.e. the random phase approximation (summing bare bubble-chains) yields the exact low-energy long-wavelength response functions to leading order in q_c/k_F, even in cases where the quasi-particle pole in the single-particle propagator is destroyed by small-$\mathbf{q}$ scattering. Similarly, bare bubbles is all one needs to construct effective interactions and dressed gauge-field propagators.

The density-current relation has first appeared (implicitly) in the work by Dzyaloshinski and Larkin [61] on one-dimensional systems. Its validity as an asymptotic relation in higher dimensions has been first pointed out by the present authors [35], and also, in the context of gauge theories, by Ioffe et al. [63]. Combining the density-current relation with a Ward identity reflecting global charge and spin conservation, one obtains a simple relation for the density and current vertex, expressed in terms of the single particle propagator G. This latter relation can be viewed as an *asymptotic Ward identity* associated with conservation of charge and spin separately on each point of the Fermi surface. In the Luttinger model, where the Fermi "surface" is a discrete set consisting of two Fermi points, this identity is exact, while in higher dimensions it holds only asymptotically for $q_c \ll k_F$. Inserting the asymptotic Ward identity into a Dyson equation for the self-energy, one obtains a complete system of equations for the single particle propagator G, valid to all orders in the coupling constant, to leading order in q_c/k_F [35].

Asymptotic Ward identities have been applied to several problems where forward scattering is dominant: low-dimensional systems with short-range interactions [35] or long-range interactions [38], and fermions coupled to gauge fields [63].

An alternative way of treating forward-scattering dominated problems is *bosonization*, where fermionic creation and annihilation operators are represented in terms of (bosonic) density fluctuation operators. This technique has been introduced by Mattis and Lieb [22] and by Luther and Peschel [65] to the analysis of one-dimensional systems, and has subsequently proved to be very useful [20, 25, 23, 24]. An early generalization of bosonization techniques to higher (than one) dimensions by Luther [67] has met only with limited success, and remained essentially unnoticed. Recently, however, Haldane [68] pioneered a different extension to $d > 1$, which turned out to be more successful. Haldane's scheme of d-dimensional bosonization has been elaborated in detail by Houghton and Marston [69], and others [70, 71, 72]. An alternative formulation of Haldane's bosonization, using functional integrals instead of operators and Hamiltionians, has been proposed by Kopietz and Schönhammer [73], and further elaborated by Kopietz et al. [62]. This latter version allows one to derive at least formal expressions for the corrections to the "non-interacting boson approximation", which yields only the leading terms in the small-$\mathbf{q}$ limit correctly.

The explicit evaluation of corrections turned out to be quite difficult, however, and has so far been achieved only to finite order in the coupling constant for a relatively simple quantity, i.e. the long-wavelength density-density correlation function [62].

To leading order in q_c/k_F, the results obtained from the bosonization technique are identical to those from the asymptotic Ward identity approach. Hence, both techniques can be regarded as different versions of an expansion in q_c/k_F. In both techniques the calculation of corrections to the leading order in q_c/k_F is possible in principle but very difficult in practice.

A third approach that has been used in the analysis of singular interactions [36] and the gauge problem [64] is the so-called *eikonal approximation*. This approximation can also be motivated via an expansion in q_c/k_F, and yields, for small q_c, essentially the same results as the other two methods [72].

1.4. CONTENTS OF THIS ARTICLE

The main purpose of this review is to describe the structure of conventional Fermi liquids and forward-scattering-dominated non-Fermi liquids in one framework, making thus common aspects of at first sight quite different systems obvious. Such a framework is provided by the combined power of the renormalization group, conservation laws and asymptotic properties of forward scattering processes. The latter become crucial in non-Fermi liquids and also in Fermi liquids with large subleading corrections due to residual forward scattering.

The renormalization group, which provides a link between microscopic systems and effective low-energy theories, will be introduced in Sec. 2. The role of exact conservation laws, such as global charge and spin conservation is discussed in Sec. 3. Special properties of and techniques for forward scattering are the subject of Sec. 5. We derive the important loop-cancellation and present the asymptotic Ward identity approach in considerable detail, because no such presentation has so far been published. Bosonization techniques will also be reviewed, and the reader will be referred to the vast literature on the subject.

The remaining sections are devoted to a discussion of several distinct systems and universality classes. In Sec. 4 the structure of Fermi liquids is reviewed from a renormalization group point of view. This section contains merely a description of the most common universality class, not a detailed investigation of its stability. In Sec. 6 we apply the general concepts from Secs. 2,3 and 5 to one-dimensional systems, obtaining thus in particular the well-known low-energy structure of Luttinger liquids. Sec. 7 contains a quantitative analysis of residual forward scattering in d-dimensional systems with short-range interactions, where d is continued to non-integer values. Such a continuation is very instructive, since it connects otherwise isolated results. The critical dimension for the sta-

bility of quasi-particles with respect to residual forward scattering and the crossover from Luttinger liquid behavior in one dimension to Fermi liquid behavior in higher-dimensional systems are determined. Sec. 8 addresses similar issues as Sec. 7, but now for long-range density-density interactions. In Sec. 9 we discuss the low-energy physics of Fermi systems coupled to abelian gauge-fields.

As a byproduct of our analysis of systems with arbitrary dimensionality we have obtained several new explicit analytic results for various quantities such as particle-hole and particle-particle bubbles in d dimensions, which, except for results for the particle-hole bubble in systems with a quadratic dispersion relation [74, 37], can so far be found in the literature only separately in $d = 1, 2, 3$. For a linearized dispersion relation this material, which may be useful also in other contexts (e.g. ϵ-expansions around a critical dimension), is contained in Sec. 7, while results for a quadratic dispersion are listed in the Appendix.

We emphasize that this is *not* a comprehensive review of *all* known non-Fermi liquids, but only of those where the break-down of Fermi liquid theory is driven by *forward scattering*. In particular, models where non-Fermi liquid behavior is due to local degrees of freedom, special shapes of the Fermi surface, special band-structure, or strong scattering with large momentum transfers have been addressed only briefly in Sec. 1.2, but will not be discussed in the bulk of the paper. Note also that we deal only with *pure* systems, i.e. disorder effects are not treated.

2. RENORMALIZATION GROUP

To one-dimensional Fermi systems, renormalization group (RG) methods have already been applied twenty years ago, driven by the necessity to deal with a variety of competing infrared divergences obtained in weak coupling perturbation theory [20]. Problems with singularities in many different channels cannot be solved by common resummations of subsets of diagrams; rather a systematic loop expansion in a renormalization group framework must be done. For today's standards this is quite simple in one-dimensional systems, where the Fermi surface consists only of two discrete points, leading to a very limited finite number of non-irrelevant coupling constants.

In higher dimensions, a general renormalization group for Fermi systems has been formulated only much later. In $d > 1$ there is an infinite number of non-irrelevant couplings and the scale k_F does not simply drop out of the low-energy theory as in one dimension. Nevertheless the reason for the delayed development of a systematic RG approach for interacting Fermi systems in higher dimensions was certainly not that it was too difficult to do; actually it might have been tackled immediately after Wilson's [75] works. The reason was rather that nobody seemed to find it useful, since no generic failure of perturbation theory was encountered. The few infrared singularities appearing in two- and three-dimensional Fermi systems, especially in the Cooper channel, could usually be handled by simpler means. A notable exception: disordered systems, non-interacting or interacting, required special RG treatments which, driven by necessity, have indeed been successfully developed [76]. In addition, *qualitative* renormalization group ideas have been underlying many phenomenological approaches, especially Fermi liquid theory. Much of RG language has tacitly crept into the theory of interacting Fermi systems [77]. Nevertheless, a quantitative and general formulation has been lacking up to recently. Motivated by the issue of non-Fermi liquid behavior, and the related discussion on the validity of perturbation theory, such a formulation has now been worked out almost simultaneously by several groups, all of them following Wilson's idea of integrating out fast modes.

Feldman and Trubowitz [15], and independently Benfatto and Gallavotti [16], have provided a rigorous version of Wilson's RG, aiming at a non-perturbative control of interacting Fermi systems at least within a certain finite radius of convergence in coupling space. Rigorous results have indeed been obtained in one-dimensional [16, 78, 79] and two-dimensional [17, 80] systems. According to these studies, there is a finite radius of

convergence in coupling space (at least for short-range interactions) inside which no hitherto unknown instabilities and/or non-perturbative effects can occur. Higher dimensions than two, where surprises are even less expected, seem to be technically more difficult, such that no significant rigorous results have so far been obtained. An instructive review of the above developments has been given by Chen, Fröhlich and Seifert [81].

A more intuitive formulation of Wilson's RG for interacting Fermi systems has been presented by Shankar [18, 19]. Basic renormalization group ideas, many technical details, and relations between RG and Fermi liquid theory are explained at length in Shankar's recent review article [19].

A brief heuristic description of the renormalization group for Fermi systems from an elementary particle physicists viewpoint has been provided by Polchinski [82]. Fermi liquid theory is viewed as a natural effective (low-energy) field theory of a normal interacting Fermi system.

In this section we will present a concise introduction to the renormalization group for Fermi systems, focussing mainly on those aspects that are important for the rest of the paper. We shall use the RG as a link between microscopic systems and effective low-energy theories, which are specified by renormalized effective actions with running (i.e. cutoff-dependent) coupling functions. The effective action is obtained (at least in principle) from the microscopic theory by integrating out high-energy states outside a thin shell of width Λ around the Fermi surface. In many cases of interest one reaches a *fixed point* in the low-energy limit, i.e. an effective action $\bar{S}^*$ which is asymptotically invariant under further reduction of the cutoff Λ. This fixed point action can often be solved exactly, i.e. the leading low-energy behavior can be expressed exactly in terms of fixed point couplings. Fermi liquids and one-dimensional Luttinger liquids provide two examples for this favorable situation. In the former case the fixed point action yields the quasi-particle velocity and the quasi-particle interactions.

2.1. FUNCTIONAL INTEGRAL REPRESENTATION

We consider a d-dimensional interacting Fermi system, which may be either a one-band lattice system or a continuum system. The dynamics is specified by a Hamiltonian

$$H = \sum_{\mathbf{k},\sigma} \epsilon_{\mathbf{k}}\, a^\dagger_{\mathbf{k}\sigma} a_{\mathbf{k}\sigma} + \tfrac{1}{2V} \sum_{\mathbf{k},\mathbf{k}',\mathbf{q}} \sum_{\sigma\sigma'} g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}';\mathbf{q}}\, a^\dagger_{\mathbf{k}-\mathbf{q}/2,\sigma} a_{\mathbf{k}+\mathbf{q}/2,\sigma} a^\dagger_{\mathbf{k}'+\mathbf{q}/2,\sigma'} a_{\mathbf{k}'-\mathbf{q}/2,\sigma'} \tag{2.1}$$

where $a^\dagger_{\mathbf{k}\sigma}$ and $a_{\mathbf{k}\sigma}$ are the usual creation and annihilation operators for spin-$\frac{1}{2}$ fermions with momentum $\mathbf{k}$ and spin projection σ. Here $\epsilon_{\mathbf{k}}$ is a general dispersion relation, while $g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}';\mathbf{q}}$ parametrizes a two-particle interaction, and V is the volume of the system. We are interested in the *low-energy* thermodynamics, response and correlation functions.

The thermodynamics of the system can be derived from the partition function $\mathcal{Z}$, which can be written as [83, 84]

$$\mathcal{Z} = \int \prod_k d\psi_k d\psi_k^* \, e^{S_0[\psi,\psi^*]+S_I[\psi,\psi^*]} \tag{2.2}$$

Here ψ_k, ψ_k^* are Grassmann variables, where the index $k = (k_0, \mathbf{k})$ includes fermionic Matsubara energies k_0 and momenta $\mathbf{k}$. Spin variables have been suppressed here to avoid proliferation of indices; they are very easy to include whenever necessary. The action $S = S_0 + S_I$ contains a quadratic term

$$S_0 = \int_k \psi_k^*(ik_0 - \epsilon_{\mathbf{k}} + \mu)\psi_k \tag{2.3}$$

which includes the kinetic energy $\epsilon_{\mathbf{k}}$ and the chemical potential μ, and a quartic term

$$S_I = -\tfrac{1}{2} \int_{k,k',q} g_{\mathbf{k}\mathbf{k}';\mathbf{q}} \, \psi^*_{k-q/2}\psi^*_{k'+q/2}\psi_{k'-q/2}\psi_{k+q/2} \tag{2.4}$$

describing two-body interactions. At zero temperature, $\int_k$ is a shorthand notation for $(2\pi)^{-(d+1)} \int dk_0 \int d^d\mathbf{k}$. For finite temperatures T, the energy integrals have to be replaced by Matsubara sums, i.e. $(2\pi)^{-1} \int dk_0 \mapsto T \sum_{k_0}$, where the summation runs over discrete energies $(2n-1)\pi T$ with integer n. The interaction term can also be written as

$$S_I = -\tfrac{1}{4} \int_{k,k',q} \Gamma_0(k, k'; q) \, \psi^*_{k-q/2}\psi^*_{k'+q/2}\psi_{k'-q/2}\psi_{k+q/2} \tag{2.5}$$

where Γ_0 is the bare vertex or antisymmetrized coupling, which is related to g by

$$\Gamma_0(k, k'; q) = g_{\mathbf{k}\mathbf{k}';\mathbf{q}} - g_{(\mathbf{k}+\mathbf{k}'-\mathbf{q})/2,(\mathbf{k}+\mathbf{k}'+\mathbf{q})/2;\mathbf{k}-\mathbf{k}'} \tag{2.6}$$

A diagrammatic representation of the bare interaction is shown in Fig. 2.1.

Correlation functions can be obtained from a generating functional [83, 84]

$$\mathcal{G}[\chi, \chi^*] = \log \Big\{ \mathcal{Z}^{-1} \int \prod_k d\psi_k d\psi_k^* \, e^{S_0[\psi,\psi^*]+S_I[\psi,\psi^*]+\int_k(\psi_k^*\chi_k+\chi_k^*\psi_k)} \Big\} \tag{2.7}$$

where a source term with Grassmann variables χ_k, χ_k^* as source fields has been added to the action. Functional derivatives with respect to the source fields generate connected Euclidean n-particle Green functions:

$$\tilde{G}_n(k_1, .., k_n; k_1', .., k_n') = (-)^n \langle \psi_{k_1} \dots \psi_{k_n} \psi^*_{k_n'} \dots \psi^*_{k_1'} \rangle_c =$$

$$\frac{\delta^n}{\delta\chi^*_{k_1} \dots \delta\chi^*_{k_n}} \frac{\delta^n}{\delta\chi_{k_n'} \dots \delta\chi_{k_1'}} \mathcal{G}[\chi, \chi^*] \Big|_{(\chi,\chi^*)=0} \tag{2.8}$$

where $\langle \ldots \rangle_c$ is the connected part of the average

$$\langle \ldots \rangle = \mathcal{Z}^{-1} \int \prod_k d\psi_k d\psi_k^* \, e^{S_0[\psi,\psi^*]+S_I[\psi,\psi^*]} \ldots \tag{2.9}$$

Energy and momentum conservation implies that $\tilde{G}_n$ can be written as

$$\tilde{G}_n(k_1,..,k_n;k_1',..,k_n') = (2\pi)^{d+1}\delta(k_1'+...+k_n'-k_1-...-k_n)\, G_n(k_1,..,k_n;k_1',..,k_n') \tag{2.10}$$

and in particular

$$\tilde{G}_1(k;k') = (2\pi)^{d+1}\delta(k'-k)\, G(k) \tag{2.11}$$

Useful shorthand notations are $\mathcal{D}[\psi,\psi^*] := \prod_k d\psi_k d\psi_k^*$ for the integration measure and $[\psi^*\chi + \chi^*\psi] := \int_k (\psi_k^*\chi_k + \chi_k^*\psi_k)$ for the source term. Thus, in compact notation, the partition function reads

$$\mathcal{Z} = \int \mathcal{D}[\psi,\psi^*] \, e^{S_0[\psi,\psi^*]+S_I[\psi,\psi^*]} \tag{2.12}$$

and the generating functional becomes

$$\mathcal{G}[\chi,\chi^*] = \log\Big\{ \mathcal{Z}^{-1} \int \mathcal{D}[\psi,\psi^*] \, e^{S_0[\psi,\psi^*]+S_I[\psi,\psi^*]+[\psi^*\chi+\chi^*\psi]} \Big\} \tag{2.13}$$

To deal with correlation functions for *composite* operators it is often convenient to introduce suitable additional source terms. For example, correlation functions involving density fluctuation operators can be generated by adding a term

$$[\phi\rho] := \int_q \phi_q \rho_q \quad \text{where} \quad \rho_q = \int_k \psi_{k-q/2}^* \psi_{k+q/2} \tag{2.14}$$

to the action, and taking functional derivatives with respect to the source field ϕ_q.

Concerning the momentum integrals we note that in condensed matter physics there is always a natural cutoff for large momenta: For lattice systems, such as electrons in a metal, momentum space is compact anyway (Brillouin zone). Continuum systems such as ^{3}He have a physical cutoff given roughly by the inverse atomic length scale.

2.2. MODE ELIMINATION AND EFFECTIVE ACTIONS

The non-interacting single-particle propagator

$$G_0(k) = \frac{1}{ik_0 - \epsilon_{\mathbf{k}} + \mu} \tag{2.15}$$

is singular for $k_0 \to 0$ and $\epsilon_{\mathbf{k}} \to \mu$. The *Fermi surface* of the non-interacting system

$$\partial\mathcal{F} = \{\mathbf{k} : \epsilon_{\mathbf{k}} = \mu\} \tag{2.16}$$

separates the Fermi sea $\mathcal{F} = \{\mathbf{k} : \epsilon_{\mathbf{k}} < \mu\}$ from its complement $\bar{\mathcal{F}}$ in momentum space.

We now integrate out "*fast modes*" ψ_k, ψ_k^* with momenta $\mathbf{k}$ far from the Fermi surface $\partial\mathcal{F}$, such that only "*slow modes*", with momenta whose distance $d(\mathbf{k}, \partial\mathcal{F})$ from the Fermi surface is smaller than a certain cutoff Λ, remain to be integrated (see Fig. 2.2). This yields

$$\mathcal{Z} = \int \mathcal{D}^{<\Lambda}[\psi, \psi^*]\, e^{S^\Lambda[\psi,\psi^*]} \tag{2.17}$$

where the *effective action* S^Λ depends on slow modes only, and is given by

$$e^{S^\Lambda[\psi,\psi^*]} := \int \mathcal{D}^{>\Lambda}[\psi, \psi^*]\, e^{S[\psi,\psi^*]} \tag{2.18}$$

Here $\mathcal{D}^{>\Lambda}[\psi, \psi^*]$ and $\mathcal{D}^{<\Lambda}[\psi, \psi^*]$ denote integration of fast and slow modes, respectively. Restricting the source variables χ_k, χ_k^* to momenta $\mathbf{k}$ with $d(\mathbf{k}, \partial\mathcal{F}) < \Lambda$, the generating functional $\mathcal{G}$ can be expressed in terms of the effective action as

$$\mathcal{G}[\chi, \chi^*] = \log\left\{ \mathcal{Z}^{-1} \int \mathcal{D}^{<\Lambda}[\psi, \psi^*]\, e^{S^\Lambda[\psi,\psi^*] + \int_k^{<\Lambda} (\psi_k^* \chi_k + \chi_k^* \psi_k)} \right\} \tag{2.19}$$

Green functions with momenta inside the Λ-shell defined by $d(\mathbf{k}, \partial\mathcal{F}) < \Lambda$ can be generated from this restricted functional by functional derivatives as in (2.8).

We have not yet specified our "distance" $d(\mathbf{k}, \partial\mathcal{F})$. In fact, there are several possibilities. For isotropic systems, one may simply choose the Euclidean distance in momentum space, $d(\mathbf{k}, \partial\mathcal{F}) = ||\mathbf{k}| - k_F|$, where k_F is the Fermi momentum, i.e. the radius of the Fermi sphere. For anisotropic systems, especially in cases where the Fermi surface is close to the Brillouin zone boundary, it is better to define a distance as $d(\mathbf{k}, \partial\mathcal{F}) = |\epsilon_\mathbf{k} - \mu|$; in this case Λ has the dimension of energy instead of momentum. In the mathematical literature distances $d(k, \partial\mathcal{F}) = [k_0^2 + (\epsilon_\mathbf{k} - \mu)^2]^{1/2}$ in $(k_0, \mathbf{k})$-space are used [17]. A choice of a distance that depends only on $\mathbf{k}$ (not on k_0) allows for an interpretation of the effective action as an action for a system with a restricted set of single-particle states.

The effective action contains quadratic, quartic and higher order monomials in the fields which remain to be integrated:

$$\begin{aligned} S^\Lambda[\psi, \psi^*] = &\int_k^{<\Lambda} [G^\Lambda(k)]^{-1} \psi_k^* \psi_k - \tfrac{1}{4} \int_{k,k',q}^{<\Lambda} \Gamma^\Lambda(k, k'; q)\, \psi_{k-q/2}^* \psi_{k'+q/2}^* \psi_{k'-q/2} \psi_{k+q/2} \\ &+ \mathcal{O}[(\psi^* \psi)^3] \end{aligned} \tag{2.20}$$

where all *fermionic* momenta (not the momentum transfer $\mathbf{q}$) are now restricted by the cutoff Λ. All the terms in S^Λ can be expanded in powers of the bare interactions, and may be represented by Feynman diagrams for connected n-particle Green functions with external lines amputated. Internal lines are integrated only over momenta outside the Λ-shell. In particular, the quadratic term in S^Λ is given by $[G^\Lambda(k)]^{-1} = G_0^{-1}(k) - \Sigma^\Lambda(k)$, where $G_0^{-1}(k) = ik_0 - \epsilon_\mathbf{k} + \mu$ and $\Sigma^\Lambda(k)$ is the self-energy with internal lines restricted

to momenta far from the Fermi surface. Note that in the quadratic part no one-particle reducible diagrams contribute, because $\mathbf{k}$ in (2.20) must be inside the Λ-shell, while momenta on internal lines must be outside. The quartic term is given by the two-particle vertex function $\Gamma^{\Lambda}(k, k'; q)$, i.e. the two-particle Green function with external propagators amputated (no one-particle reducible terms exist in this case). Higher order terms are given by amputated n-particle Green functions. Note that for odd $n > 1$ one-particle reducible diagrams contribute. For more details on the perturbation expansion and its diagrammatic representation, see Refs. [15, 16, 81, 19].

In general the Fermi surface is shifted by interactions (with a change of shape, if the system is anisotropic). To obtain a well behaved perturbation series, one must not expand around S_0, but rather around another suitably chosen quadratic action with a Fermi surface already in its interacting position. This is achieved by adding a counterterm of the form $\int_k \delta\mu_{\mathbf{k}} \psi^*_k \psi_k$ to S_0, and subtracting it from S_I, i.e. one defines

$$S'_0 = S_0 + \int_k \delta\mu_{\mathbf{k}}\, \psi^*_k \psi_k \,, \quad S'_I = S_I - \int_k \delta\mu_{\mathbf{k}}\, \psi^*_k \psi_k \tag{2.21}$$

The perturbation expansion is then carried out around S'_0. The actual value of $\delta\mu_{\mathbf{k}}$ is of course not known a priori, but can be determined (order by order) by a self-consistency condition.[3] A $\mathbf{k}$-dependent quadratic counterterm can also be used to adjust the Fermi velocity (s. below).

In this work we focus on normal state properties, and therefore consider energy scales well above scales dominated by possible symmetry breaking. We note, however, that symmetry breaking can be treated by adding suitable quadratic counterterms. To deal with superconductivity, for example, anomalous quadratic terms $\psi\psi$ and $\psi^*\psi^*$ must be included in S'_0 (and accordingly subtracted in S'_I) [17].

The mode elimination can be iterated, by calculating $S^{\Lambda'}$ with $\Lambda' < \Lambda$ in terms of S^{Λ}. At a generic scale Λ, the effective action S^{Λ} can be decomposed in a quadratic part S_0^{Λ} and a rest S_I^{Λ} (containing interactions and quadratic counterterms) such that S_0^{Λ} has the Fermi surface of the interacting system.

In the following we will frequently consider *isotropic* systems for pedagogical reasons. In this case the Fermi surface is spherically symmetric irrespective of interactions and the counterterm in (2.21) reduces to a constant shift $\delta\mu$ of the chemical potential. It is not hard to extend the formalism to anisotropic situations, while concrete calculations become of course more tedious.

Expanding the kernel of the (purely) quadratic part S_0^{Λ} of the action around the Fermi

[3] To show rigorously that this procedure really works, to any order in perturbation theory, is a subtle problem that has been recently solved by Feldman et al. [85].

surface, one gets

$$S_0^\Lambda = \int_k^{<\Lambda} (Z_{\mathbf{k}_F}^\Lambda)^{-1} \psi_k^* [ik_0 - \bar{v}_{\mathbf{k}_F}^\Lambda k_r - \bar{\Sigma}^\Lambda(k)] \psi_k \tag{2.22}$$

Here momenta $\mathbf{k}$ are represented by the pair $(k_r, \mathbf{k}_F)$ where $\mathbf{k}_F$ is the projection of $\mathbf{k}$ on the Fermi surface, while k_r is its oriented distance (defined positive for $\mathbf{k}$ outside the Fermi surface, and negative inside), i.e.

$$k_r := |\mathbf{k}| - k_F \tag{2.23}$$

for isotropic systems. The *field renormalization factor*

$$Z_{\mathbf{k}_F}^\Lambda = \left[1 - \partial\Sigma^\Lambda(k_0, \mathbf{k})/\partial(ik_0)\right]^{-1}\Big|_{(0,\mathbf{k}_F)} \tag{2.24}$$

is a positive constant < 1. The effective *Fermi velocity* $\bar{\mathbf{v}}_{\mathbf{k}_F}^\Lambda$ is related to the self-energy by

$$\bar{\mathbf{v}}_{\mathbf{k}_F}^\Lambda = \bar{\mathbf{v}}_{\mathbf{k}_F}^{c\Lambda} + \delta\bar{\mathbf{v}}_{\mathbf{k}_F}^\Lambda \tag{2.25}$$

where

$$\bar{\mathbf{v}}_{\mathbf{k}_F}^{c\Lambda} = Z_{\mathbf{k}_F}^\Lambda \left[\mathbf{v}_{\mathbf{k}} + \partial\Sigma^\Lambda(k_0, \mathbf{k})/\partial\mathbf{k}\right]\Big|_{(0,\mathbf{k}_F)} \tag{2.26}$$

and $\delta\bar{\mathbf{v}}_{\mathbf{k}_F}^\Lambda = Z_{\mathbf{k}_F}^\Lambda \, \delta\mathbf{v}_{\mathbf{k}_F}^\Lambda$ is a (renormalized) counterterm. Here $\mathbf{v}_{\mathbf{k}} = \nabla\epsilon_{\mathbf{k}}$ is the bare velocity of particles with momentum $\mathbf{k}$. Note that the limit $\Lambda \to 0$ does not commute (in general) with the $\mathbf{k}$-derivative of Σ^Λ on the Fermi surface, because particle-hole excitations with infinitesimal excitation energy can yield a finite contribution to $\partial\Sigma/\partial\mathbf{k}|_{(0,\mathbf{k}_F)}$. The velocity $\bar{\mathbf{v}}_{\mathbf{k}_F}^{c\Lambda}$ determines the renormalized *current* operator (s. Sec. 3.3). A counterterm $-\int_k \delta v_{\mathbf{k}_F}^\Lambda k_r \, \psi_k^* \psi_k$ has been added to S_0 to make $\bar{\mathbf{v}}_{\mathbf{k}_F}^\Lambda$ converge to the true Fermi velocity in the limit $\Lambda \to 0$. The residual self-energy $\bar{\Sigma}^\Lambda(k)$ vanishes faster than linearly for $k_0 \to 0$, $\mathbf{k} \to \partial\mathcal{F}$. For isotropic systems, $Z_{\mathbf{k}_F}^\Lambda = Z^\Lambda$ and $\bar{v}_{\mathbf{k}_F}^\Lambda = \bar{v}_F^\Lambda$ are constant all over the Fermi surface.

The factor $(Z^\Lambda)^{-1}$ in the quadratic part of the action can be eliminated by introducing *renormalized fields*

$$(\bar{\psi}_k, \bar{\psi}_k^*) = (Z^\Lambda)^{-1/2} (\psi_k, \psi_k^*) \tag{2.27}$$

as the new (functional) integration variables.[4] Of course Z^Λ now shows up in interaction and source terms, but can be absorbed in renormalized source fields

$$(\bar{\chi}_k, \bar{\chi}_k^*) = (Z^\Lambda)^{1/2} (\chi_k, \chi_k^*) \tag{2.28}$$

in the *renormalized vertex functions*

[4]The corresponding Jacobian is a constant and cancels in $\mathcal{G}[\chi, \chi^*]$.

$$\bar{\Gamma}^{\Lambda}(\{k_i\};\{k_i'\}) = (Z^{\Lambda})^n \Gamma^{\Lambda}(\{k_i\};\{k_i'\}) \tag{2.29}$$

for n-particle interactions, and in a renormalized chemical potential shift $\delta\bar{\mu}^{\Lambda} = Z^{\Lambda}\delta\mu^{\Lambda}$. Thus the *renormalized* effective action becomes

$$\begin{aligned}\bar{S}^{\Lambda}[\bar{\psi},\bar{\psi}^*] = &\int_k^{<\Lambda} \bar{\psi}_k^*[ik_0 - \bar{v}_F^{\Lambda}k_r - \bar{\Sigma}^{\Lambda}(k)]\bar{\psi}_k - \int_k^{<\Lambda}(\delta\bar{\mu}^{\Lambda} - \delta\bar{v}_F^{\Lambda}k_r)\,\bar{\psi}_k^*\bar{\psi}_k \\ &-\tfrac{1}{4}\int_{k,k',q}^{<\Lambda} \bar{\Gamma}^{\Lambda}(k,k';q)\bar{\psi}^*_{k-q/2}\bar{\psi}^*_{k'+q/2}\bar{\psi}_{k'-q/2}\bar{\psi}_{k+q/2} + \mathcal{O}((\bar{\psi}^*\bar{\psi})^3)\end{aligned} \tag{2.30}$$

In general the vertex function $\bar{\Gamma}^{\Lambda}(k,k';q)$ depends on three independent $(d+1)$ dimensional energy-momentum variables. However, many details of these dependences are actually irrelevant for the low-energy physics, which depends only on the behavior of $\bar{\Gamma}^{\Lambda}(k,k';q)$ in certain limits. Hence the vertex function in the effective action can be replaced by renormalized *coupling functions* $\bar{g}^{\Lambda}$ with less variables. For example, in most cases the coupling function

$$\bar{g}^{\Lambda}_{\mathbf{k}\mathbf{k}'}(q) = \lim_{k\to(0,\mathbf{k})}\lim_{k'\to(0,\mathbf{k}')} \bar{\Gamma}^{\Lambda}(k,k';q) \tag{2.31}$$

contains enough information. Neglecting all irrelevant terms one thus finds

$$\begin{aligned}\bar{S}^{\Lambda}[\bar{\psi},\bar{\psi}^*] = &\int_k^{<\Lambda} \bar{\psi}_k^*[ik_0 - \bar{v}_F^{\Lambda}k_r]\bar{\psi}_k - \int_k^{<\Lambda}(\delta\bar{\mu}^{\Lambda} - \delta\bar{v}_F^{\Lambda}k_r)\,\bar{\psi}_k^*\bar{\psi}_k \\ &-\tfrac{1}{4}\int_{k,k',q}^{<\Lambda} \bar{g}^{\Lambda}_{\mathbf{k}\mathbf{k}'}(q)\,\bar{\psi}^*_{k-q/2}\bar{\psi}^*_{k'+q/2}\bar{\psi}_{k'-q/2}\bar{\psi}_{k+q/2}\end{aligned} \tag{2.32}$$

In many specific cases (Fermi liquids etc.) the number of relevant variables in $\bar{g}^{\Lambda}$ can be further reduced, especially by replacing (regular) coupling functions by their asymptotic values on the Fermi surface.

The renormalized source term reads $\int_k^{<\Lambda}(\bar{\psi}_k^*\bar{\chi}_k + \bar{\chi}_k^*\bar{\psi}_k)$. The exact (unrenormalized) correlation functions can be obtained as functional derivatives of the renormalized generating functional

$$\bar{\mathcal{G}}[\bar{\chi},\bar{\chi}^*] := \mathcal{G}[\bar{\chi}/(Z^{\Lambda})^{1/2}, \bar{\chi}^*/(Z^{\Lambda})^{1/2}] \tag{2.33}$$

with respect to the renormalized source fields $\bar{\chi}$ and $\bar{\chi}^*$, followed by a multiplication with the respective Z-factors:

$$G_n(k_1,...,k_n;k_1',...,k_n') = (Z^{\Lambda})^n\,\bar{G}_n^{\Lambda}(k_1,...,k_n;k_1',...,k_n') \tag{2.34}$$

where $\bar{G}_n^{\Lambda}$ are the *renormalized* Green functions given by

$$(2\pi)^{d+1}\delta(k_1'+...+k_n'-k_1-...-k_n)\,\bar{G}_n^{\Lambda}(k_1,...,k_n;k_1',...,k_n') =$$

$$\frac{\delta^n}{\delta\bar{\chi}^*_{k_1}\ldots\delta\bar{\chi}^*_{k_n}}\frac{\delta^n}{\delta\bar{\chi}_{k'_n}\ldots\delta\bar{\chi}_{k'_1}}\bar{\mathcal{G}}[\chi,\chi^*]\Big|_{(\bar{\chi},\bar{\chi}^*)=0} \tag{2.35}$$

Note that renormalized correlation functions depend on the (variable) cutoff Λ, while unrenormalized correlation functions depend only on the cutoff determined by the microscopic theory.

What have we gained so far? The renormalized effective action $\bar{S}^\Lambda$ has usually a much more complicated form than the bare action S. So why analyse low energy physics in terms of effective actions instead of the bare one? The point is that for small Λ a new small *expansion parameter* has emerged: Λ/k_F. In many cases this allows one to express the exact leading and subleading low-energy behavior in terms of renormalized couplings even if these are not small. Expressing thermodynamics and correlation functions in terms of the effective parameters defining $\bar{S}^\Lambda$, and expanding in powers of small energy scales (temperature, frequency etc.), one realizes that only few terms in $\bar{S}^\Lambda$ contribute in leading order, a finite number of additional terms to the first subleading order, and so on. The asymptotic infrared behavior of the system can thus be expressed in terms of relatively few parameters.

The effective couplings $\bar{g}^\Lambda$ can be classified as *relevant*, *marginal*, or *irrelevant*, depending on whether their importance relative to the quadratic part of the action grows, remains invariant, or decreases in the limit $\Lambda \to 0$.[5]

More-than-two body interactions with a finite low-energy limit are usually irrelevant. More precisely, the contributions from finite (n+1)-particle interactions are suppressed by a factor Λ with respect to those from n-particle interactions. As an illustration, let us estimate the order of magnitude of the three self-energy diagrams in Fig. 2.3 by naive power-counting. The shaded boxes represent effective two-particle and three-particle interactions. Internal lines correspond to propagators $G_0^\Lambda(k) = [ik_0 - \bar{v}_F^\Lambda k_r - \bar{\Sigma}^\Lambda(k)]^{-1}$. Momenta on internal lines must lie inside the thin Λ-shell around the Fermi surface defined by $d(\mathbf{k}, \partial\mathcal{F}) < \Lambda$, corresponding to small excitation energies $\bar{v}_F^\Lambda k_r$. The propagators become big (of order $(\bar{v}_F^\Lambda\Lambda)^{-1}$), if the energy variables are also small, i.e. $k_0 < \bar{v}_F^\Lambda\Lambda$. Hence, energy-momentum variables leading to a big propagator fill a volume proportional to Λ^2 in $(k_0, \mathbf{k})$-space. Notice that this volume is independent of the space dimension d because the propagator is singular on a $(d-1)$-dimensional surface, which reduces to a point-singularity (as in standard critical phenomena) only for $d = 1$. Thus, according to naive power-counting, the value of a diagram with l internal lines and r integrations (equal to the number of loops) is proportional to Λ^{2r-l}. In particular, the first two

[5]Note that this classification depends on the choice of $\bar{S}_0^\Lambda$! Relevant couplings may sometimes be avoided by a clever choice of the "non-interacting" part $\bar{S}_0^\Lambda$.

diagrams (containing only two-particle interactions) in Fig. 2.3 are of order Λ, while the last one (containing a 3-particle interaction) is of order Λ^2. It is easy to see that the replacement of an n-particle interaction in a diagram by an (n+1)-particle interaction enhances the number of internal lines and integrations by one, leading to an extra factor Λ in the power-counting estimate. Actually it turns out that in dimensions $d > 1$ the naive power-counting often over-estimates the value of a diagram, since geometrical phase-space restrictions reduce the actual integration volume, leading to additional Λ-powers in most cases.

The above power-counting holds only for interactions that remain finite in the low-energy limit. However, it is easy to see that in a Wilson renormalization scheme effective n-particle interactions of order Λ^{2-n} may be generated. An example for a contribution to an effective three-particle interaction with a low-energy limit of order Λ^{-1} is shown in Fig. 2.4; the internal line may have momenta outside but close to the Λ-shell even if all external lines carry momenta inside the Λ-shell. Inserting the effective interaction in Fig. 2.4 into the third self-energy diagram in Fig. 2.3, one obtains contribution of order Λ within naive power-counting, i.e. that three-particle interaction seems to be as important as finite two-particle interactions. Fortunately, in practice (especially for the purposes of the present article) one can usually avoid dealing with these many-particle interactions, for different reasons in different cases. In particular, in one-dimensional systems the Wilson RG can be replaced by a field-theoretic RG in the low-energy regime, with a suitable effective two-body action that incorporates all higher order interactions. The book-keeping of generated terms is different in the field-theoretic RG,[6] making effective many-particle interactions ($n > 2$) generally irrelevant. In higher dimensions additional Λ-powers coming from geometrical phase space contraints suppress the effects of many-particle interactions more strongly than the above simple power-counting would predict. A deeper understanding of the role of effective many-particle interactions in Wilson's RG version for Fermi systems has been reached in the mathematical literature [15, 16]. In the following we will not consider effective n-particle interactions with $n > 2$ any more, assuming that they are either irrelevant or somehow effectively incorporated in one-particle and two-particle terms in the low-energy action.

2.3. LOW-ENERGY COUPLING-SPACE

Let us now try to become acquainted with the huge coupling space for two-particle interactions in Fermi systems, and find a suitable classification of couplings. In general, two-particle interactions scatter energy momenta k_1 and k_2 into k_1' and k_2', where energy-momentum conservation imposes the restriction $k_1 + k_2 = k_1' + k_2'$, which can be manifestly

[6] For a short discussion on this point, see Shankar [19].

built in by parametrizing the vertex function in terms of three variables k, k' and q such that $k_1 = k + q/2$, $k_2 = k' - q/2$, $k_1' = k - q/2$ and $k_2' = k' + q/2$.

For small cutoffs Λ, the restriction of all momenta of fermions to a thin shell around the Fermi surface combined with total momentum conservation in scattering processes leads to drastic geometric constraints on the *angles* between the momenta of the two ingoing and outgoing particles, in addition to the radial constraint directly imposed by the cutoff. To see this, let us consider the limit $\Lambda \to 0$, where all the particles must be situated *on* the Fermi surface, which we assume to be spherical for simplicity.

Let us start with *two-dimensional* systems. Here it is easy to see that there are three distinct types of possible (i.e. momentum conserving) scattering processes, which can be parametrized by a single angle each (see Fig. 2.5):

$$\begin{array}{ll} \text{"}\textit{forward}\text{" (F) scattering:} & \mathbf{k}_1' = \mathbf{k}_1 \text{ and } \mathbf{k}_2' = \mathbf{k}_2 \\ \text{"}\textit{exchange}\text{" (E) scattering:} & \mathbf{k}_1' = \mathbf{k}_2 \text{ and } \mathbf{k}_2' = \mathbf{k}_1 \\ \text{"}\textit{Cooper}\text{" (C) scattering:} & \mathbf{k}_1 + \mathbf{k}_2 = 0. \end{array} \tag{2.36}$$

Actually forward and exchange scattering are equivalent for spinless fermions or fermions with the same spin projection $\sigma_1 = \sigma_2$, since the vertex is antisymmetric with respect to interchange of incoming (or outgoing) particles. Forward and exchange scattering are however distinct for $\sigma_1 \neq \sigma_2$. Forward and exchange scattering can be uniquely parametrized by the angle between the momenta of the incoming particles $\theta = \angle(\mathbf{k}_1, \mathbf{k}_2)$, while Cooper scattering is parametrized by the angle defined by the momentum transfer $\phi = \angle(\mathbf{k}_1, \mathbf{k}_1')$. Whereever the vertex function $\bar{\Gamma}^\Lambda$ is regular near the Fermi surface, it can be replaced by the three coupling functions $\bar{g}_F^\Lambda(\theta)$, $\bar{g}_E^\Lambda(\theta)$ and $\bar{g}_C^\Lambda(\phi)$, which are given by the three Fermi surface limits of $\bar{\Gamma}^\Lambda(k, k'; q)$ according to the above classification.

The above classification of scattering processes represents a canonical generalization of the *g-ology* classification in *one-dimensional* systems [20] to two dimensions. Specializing to one dimension, where only two angles (0 and π) exist, one finds the following correspondence [86]:

$$\begin{array}{ll} g_F(0) = g_4, & g_F(\pi) = g_2 \\ g_E(0) = g_4, & g_E(\pi) = g_1 \\ g_C(0) = g_2, & g_C(\pi) = g_1 \end{array} \tag{2.37}$$

where the numbering $g_1, .., g_4$ follows the usual g-ology convention (here we have suppressed bars and Λ's for easier readability). Note that processes which are generically distinct in $d = 2$ may become equivalent for the special angles 0 and π, which are the only angles in $d = 1$. The g-ology coupling g_3 does not appear here since it describes umklapp processes which have not been considered. Umklapp processes exist only in lattice

systems; at low energy they are important only in special cases (at specific fillings).

In *three-dimensional* systems, two particles whose momenta span an angle θ may scatter into new states with momenta spanning the same angle θ (and their sum pointing in the same direction of course). For $\theta \neq \pi$ this leaves a one-dimensional degree of freedom, which can be parametrized by the angle ϕ spanned by the initial and the final momentum of one of the scattering particles. In two dimensions, ϕ could be either 0 or θ, corresponding to "forward" and "exchange" scattering, respectively. In three dimensions these two extreme cases are continuously connected in one class of scattering processes, which we call "*normal*" scattering. For $\theta = \pi$ one has Cooper scattering, with two degrees of freedom for the tranferred momentum; for the isotropic case considered here the corresponding coupling strength can however be parametrized by a single angle ϕ, e.g. the one spanned by the initial and the final momentum of one of the particles. Thus, in three dimensions we have two distinct classes of two-particle interactions on the Fermi surface:

$$\begin{aligned} &\text{"}\textit{normal}\text{" (N) scattering:} \quad \angle(\mathbf{k}_1, \mathbf{k}_2) = \angle(\mathbf{k}_1', \mathbf{k}_2') \neq \pi \\ &\text{"}\textit{Cooper}\text{" (C) scattering:} \quad \angle(\mathbf{k}_1, \mathbf{k}_2) = \angle(\mathbf{k}_1', \mathbf{k}_2') = \pi. \end{aligned} \tag{2.38}$$

with coupling functions $\bar{g}_N^\Lambda(\theta, \phi)$ and $\bar{g}_C^\Lambda(\phi)$, respectively. The coupling $\bar{g}_F^\Lambda(\theta) = \bar{g}_N^\Lambda(\theta, 0)$, describing *forward* scattering, plays a special role in Fermi liquid theory.

The above analysis has led to a purely *kinematic* classification of all scattering processes that are geometrically possible near the Fermi surface. To see how the corresponding effective interactions behave in the low-energy limit one must estimate the *phase-space* for scattering processes, i.e. estimate Feynman diagrams. The qualitative behavior of effective interactions depends on the properties of the bare interactions, such as signs and regularity properties (bounded or singular).

As a specific example, let us discuss an isotropic Fermi system with short-range interactions in three dimensions.[7] Our aim is to understand the behavior of effective two-particle couplings $\bar{g}^\Lambda$ in the low-energy regime and to reveal possible sources of instabilities. As a first step one would try to calculate $\bar{g}^\Lambda$ in perturbation theory by evaluating Feynman diagrams for the two-particle vertex, with momenta on internal propagators outside a Λ-shell around the Fermi surface. The diagrams contributing to second order in the coupling constant are listed in Fig. 2.6. For almost all k, k', and q a direct expansion of $\bar{\Gamma}^\Lambda(k, k'; q)$ in powers of bare couplings has finite coefficients even in the limit $\Lambda \to 0$. In these cases one can actually integrate all momenta down to the Fermi surface in one shot. However, for $k' = -k$ (the Cooper channel), already the second order diagram (a) in Fig. 2.6 leads

[7] Interactions which are short-ranged in real space are bounded in momentum space.

to a logarithmically divergent contribution to $\bar{\Gamma}^{\Lambda}_{kk';q}$ in the limit $\Lambda \to 0$. This is a case for the renormalization group transformation.

To calculate the flow of $\bar{\Gamma}^{\Lambda}(k, k'; q)$ with $k' = -k$ under infinitesimal reductions of the cutoff, let us parametrize interactions that scatter p and $-p$ into p' and $-p'$ by a coupling function $\bar{g}^{\Lambda}_{pp'}$. To obtain simple (i.e. instructive) analytic results, we assume that $\bar{g}^{\Lambda}_{pp'}$ depends only on the angle ϕ between initial and final momenta, not on frequencies or moduli of momenta, i.e. $\bar{g}^{\Lambda}_{pp'} = \bar{g}^{\Lambda}_{C}(\phi_{\mathbf{pp}'})$. We will see that this assumption is self-consistent in the low-energy limit in the sense that no other dependences will be generated by the flow. Note that we are interested in the flow close to the Fermi surface, i.e. $\Lambda \ll k_F$, where 3-particle interactions are already irrelevant. Hence, to second order in the coupling, we can calculate the flow by computing the Feynman diagrams in Fig. 2.6 for ingoing external energy-momenta p and $-p$, with internal momenta restricted to the infinitesimal region between a Λ-shell and a Λ'-shell with $\Lambda' = \Lambda - d\Lambda$.

We first consider the contribution due to the diagram (a), i.e. the diagram that produces the logarithmic divergence in perturbation theory. The Feynman rules yield a contribution

$$d\bar{g}^{\Lambda}_{pp'} = \int_{\Lambda'}^{\Lambda} \frac{d^{d+1}k}{(2\pi)^{d+1}}\, \bar{g}^{\Lambda}_{pk}\, \bar{g}^{\Lambda}_{kp'}\, \frac{1}{ik_0 - \xi_{\mathbf{k}}}\, \frac{1}{-ik_0 - \xi_{-\mathbf{k}}} \tag{2.39}$$

where $\xi_{\mathbf{k}} = \epsilon_{\mathbf{k}} - \mu$ and the $\mathbf{k}$-integration is restricted to the above-mentioned infinitesimal region. The k-integration can be decomposed in integrals over k_0, k_r and the solid angle $\Omega_{\mathbf{k}}$, where k_r is restricted by the condition $\Lambda' < |k_r| < \Lambda$. Since $\bar{g}^{\Lambda}_{pk} = \bar{g}^{\Lambda}(\phi_{\mathbf{pk}})$ is independent of modula and frequencies, and $\xi_{\mathbf{k}} = \bar{v}^{\Lambda}_F k_r$ in the low-energy limit, the integrations over k_0 and k_r can be carried out explicitly, yielding (at $d = 3$)

$$d\bar{g}^{\Lambda}_{C}(\phi_{\mathbf{pp}'}) = \frac{d\Lambda}{\Lambda} \frac{k_F^2}{(2\pi)^3 \bar{v}^{\Lambda}_F} \int d\Omega_{\mathbf{k}}\, \bar{g}^{\Lambda}_{C}(\phi_{\mathbf{pk}})\, \bar{g}^{\Lambda}_{C}(\phi_{\mathbf{kp}'}) \tag{2.40}$$

Inserting a partial wave decomposition $\bar{g}^{\Lambda}_{C}(\phi) = \sum_{l=0}^{\infty} \bar{g}^{\Lambda}_{l} P_l(\cos\phi)$, and using the addition theorem for the Legendre polynomials P_l, one obtains decoupled flow equations for each angular momentum sector:

$$\frac{d\bar{g}^{\Lambda}_{l}}{d\log\Lambda} \equiv \beta_l(\bar{v}^{\Lambda}_F, \bar{g}^{\Lambda}_{l}) = \frac{1}{2\pi^2(2l+1)} \frac{k_F^2}{\bar{v}^{\Lambda}_F} (\bar{g}^{\Lambda}_{l})^2 \tag{2.41}$$

We note that for spinless fermions the coupling function must be antisymmetric in the initial (and final) momenta, i.e. only odd angular momenta l contribute here. Reinserting spin one obtains separate equations for the singlet channel, with even l, and the triplet channel, with odd l.

It is easy to see that the diagrams (b) and (c) yield a contribution of order $(d\Lambda)^2$ to the flow of the Cooper couplings, which is negligible. The flow equation (2.41), or its two-dimensional analogue, has been derived by Feldman and Trubowitz [15], and by Shankar

[18]. Of course the behavior of the two-particle vertex for small total momenta of the particles has been investigated long before. A scaling equation which is equivalent to the above flow equation can be found already in the book by Abrikosov et al. [14].

The above flow for $\bar{g}_l^\Lambda$ makes negative couplings grow in the low-energy limit, signalling an instability towards a qualitatively different phase. This is the renormalization group version of the *Cooper instability*. Positive couplings scale logarithmically to zero, i.e. they disappear in the low-energy limit, albeit very slowly.

We emphasize that the flow equation (2.41) is valid only to leading order in Λ/k_F (since irrelevant terms have been neglected) and to second order in the interaction. Including subleading orders in Λ/k_F one finds that different angular momenta couple and that non-Cooper couplings influence the flow. It has been shown that these corrections always generate some negative couplings $\bar{g}_l^\Lambda$ in the low-energy limit, even if one has started with a purely repulsive interaction [15, 19]. Ultimately, in the limit $\Lambda/k_F \to 0$, these couplings will be governed by the flow (2.41) and thus drive a Cooper instability. This renormalization group result substantiates an old, more intuitive argument by Kohn and Luttinger [87]. A quantitative numerical analysis of the flow equation shows that in isotropic systems with purely repulsive interaction the instability is strongest in the p-wave ($l = 1$) channel [88].

Perturbation theory for $\bar{\Gamma}^\Lambda(k, k'; q)$ also reveals a singularity in the limit of small momentum transfer, $q \to 0$. In contrast to the Cooper instability, this singularity is not a divergence (for imaginary frequencies), but only a non-uniqueness of limits: the limits $q_0 \to 0$, $\mathbf{q} \to 0$ and $\Lambda \to 0$ do not commute. This singularity does not signal any instability of the system, and a standard perturbation expansion is consistent in the sense that higher orders in the coupling are small if the interaction is weak. Note, however, that this latter statement holds only for the Euclidean theory, i.e. for Green functions with imaginary times or frequencies: perturbative contributions blow up, if one tries to continue to *real* frequencies. These divergencies are associated with collective modes in the interacting system (zero sound) and can be treated by a suitable resummation of Feynman diagrams (introducing particle-hole irreducible vertices). More on this will follow in the section on Fermi liquid theory.

3. EXACT CONSERVATION LAWS

In this section we will discuss important exact conservation laws, namely charge and spin conservation, and derive the associated Ward identities. We emphasize that most of these identities are well-known, at least for the case of continuum systems (see, for example, Nozières [2]). Nevertheless we find it useful to provide a concise collection of those identities which play a role in our article, and present derivations from general principles that do not depend on the assumptions of Fermi liquid theory. An important consequence of Ward identities is the cancellation of the field renormalization Z^Λ in the response of a Fermi system to low-energy long-wavelength fields that couple to the density or current (see 3.3). More stringent conservation laws obeyed by forward scattering processes and the associated Ward identities will be treated later, in Secs. 5 and 6.

3.1. CHARGE CONSERVATION

We start by deriving relations following from charge (or particle number) conservation. The charge-density fluctuation operator is defined as

$$\rho(\mathbf{q}) = \sum_{\mathbf{k}\sigma} a^\dagger_{\mathbf{k}-\mathbf{q}/2,\sigma}\, a_{\mathbf{k}+\mathbf{q}/2,\sigma} \tag{3.1}$$

For $\mathbf{q} = \mathbf{0}$, $\rho(\mathbf{0}) = N$ is the particle number operator. The (imaginary) time evolution of charge-density fluctuations is given by $\rho(\tau, \mathbf{q}) = e^{K\tau}\rho(\mathbf{q})e^{-K\tau}$ where $K = H - \mu N$ is the grand-canonical Hamiltonian. The equation of motion for $\rho(\tau, \mathbf{q})$ can be written as

$$\partial_\tau \rho(\tau, \mathbf{q}) = [H, \rho(\tau, \mathbf{q})] \tag{3.2}$$

since $[N, \rho(\mathbf{q})] = 0$. As a consequence of charge conservation, i.e. $[N, H] = 0$, the right hand side of (3.2) vanishes for $\mathbf{q} \to 0$. In all cases of interest there is a current operator $\mathbf{j}(\mathbf{q})$ such that the equation of motion for $\rho(\mathbf{q})$ assumes the form of a *continuity equation*

$$\partial_\tau \rho(\tau, \mathbf{q}) = -\,\mathbf{q} \cdot \mathbf{j}(\tau, \mathbf{q}) \tag{3.3}$$

at least for small $\mathbf{q}$. In general, the current operator depends explicitly on interactions. Inserting the decomposition $H = H_0 + H_I$ of the Hamiltonian into the commutator in (3.2), one obtains a corresponding decomposition of the current operator, $\mathbf{j} = \mathbf{j}_0 + \mathbf{j}_I$. For continuum systems with a free-particle dispersion relation $\epsilon_\mathbf{k} = \mathbf{k}^2/2m$ and a pure

density-density interaction $H_I = \frac{1}{2V}\sum_{\mathbf{q}}\, g(\mathbf{q})\,\rho(\mathbf{q})\rho(-\mathbf{q})$, one has $\mathbf{j}_I = 0$ and

$$\mathbf{j}(\mathbf{q}) = \mathbf{j}_0(\mathbf{q}) = \sum_{\mathbf{k}\sigma} (\mathbf{k}/m)\; a^\dagger_{\mathbf{k}-\mathbf{q}/2,\sigma}\; a_{\mathbf{k}+\mathbf{q}/2,\sigma} \tag{3.4}$$

For a lattice system a continuity equation of the form (3.3) holds only for small $\mathbf{q}$ (much smaller than the inverse lattice spacing), which is however the most interesting case. In the absence of hopping terms in the interaction H_I, one still has $\mathbf{j}_I = 0$. For small $\mathbf{q}$, the commutator of $\rho(\mathbf{q})$ with H_0 gives rise to a continuity equation with a current operator of the form

$$\mathbf{j}(\mathbf{q}) = \mathbf{j}_0(\mathbf{q}) = \sum_{\mathbf{k}\sigma} \nabla_{\mathbf{k}}\epsilon_{\mathbf{k}}\; a^\dagger_{\mathbf{k}-\mathbf{q}/2,\sigma}\; a_{\mathbf{k}+\mathbf{q}/2,\sigma} \tag{3.5}$$

which generalizes (3.4) to systems with a non-quadratic dispersion relation $\epsilon_{\mathbf{k}}$.

It is often useful to collect density and current operators in a (d+1)-dimensional vector $j^\mu = (\rho, \mathbf{j})$, where $\mu = 0, 1, \ldots, d$, which will also be referred to as a "current" operator.

The continuity equation (3.2) implies *Ward identities* for correlation (or Green) functions involving charge-currents. Two such correlation functions are particularly important: The (charge) current-current correlation function

$$J^{\mu\nu}(q) = -\,\tfrac{1}{V}\,\langle j^\mu(q)\, j^\nu(-q)\rangle \tag{3.6}$$

and the (charge) current vertex part

$$\Gamma^\mu_\sigma(p;q) = \langle j^\mu(q)\; a_{p-q/2,\sigma}\; a^\dagger_{p+q/2,\sigma}\rangle_{tr} \tag{3.7}$$

The above expressions are short-hand notations for thermal expectation values of (imaginary) time ordered operator products, with a subsequent time-to-frequency Fourier-transformation. The index "tr" means truncation of external legs (in a diagrammatic language), i.e. division by $G_\sigma(p+q/2)G_\sigma(p-q/2)$.

The Ward identity for $J^{\mu\nu}$ can be obtained by deriving the ($\mu = 0$)-component of the current-current correlator in time-representation

$$J^{\mu\nu}(\tau-\tau', \mathbf{q}) = -\,\tfrac{1}{V}\,\langle \mathcal{T}\; j^\mu(\tau, \mathbf{q})\; j^\nu(\tau', -\mathbf{q})\rangle \tag{3.8}$$

with respect to τ. One gets

$$\partial_\tau J^{0\nu}(\tau-\tau', \mathbf{q}) = \tfrac{1}{V}\,\langle \mathcal{T}\; \mathbf{q}\cdot\mathbf{j}(\tau, \mathbf{q})\; j^\nu(\tau', -\mathbf{q})\rangle \;-\; \tfrac{1}{V}\,\delta(\tau-\tau')\langle[j^0(\tau, \mathbf{q}), j^\nu(\tau, -\mathbf{q})]\rangle \tag{3.9}$$

where the first term on the right hand side is due to the time-derivative of $j^0(\tau, \mathbf{q})$, as given by the continuity equation, while the second one has been generated by the time-derivative of the time-ordering operator $\mathcal{T}$. Fourier transforming back to frequencies yields the Ward identity

$$(iq_0, \mathbf{q})_\mu J^{\mu\nu}(q) = \tfrac{1}{V}\,\langle[j^0(\mathbf{q}), j^\nu(-\mathbf{q})]\rangle =: c^\nu(\mathbf{q}) \tag{3.10}$$

where $(iq_0, \mathbf{q})_0 = iq_0$ while $(iq_0, \mathbf{q})_j = -q_j$ for $j = 1, \ldots, d$, and summation over repeated Greek indices is assumed. Reflection invariance implies obviously

$$c^0(\mathbf{q}) = 0 \tag{3.11}$$

Inserting the expression (3.5) for the current operator into the commutator in (3.10), one obtains

$$\mathbf{c}(\mathbf{q}) := \tfrac{1}{V} \langle [j^0(\mathbf{q}), \mathbf{j}(-\mathbf{q})] \rangle = V^{-1} \sum_{\mathbf{k}\sigma} \mathbf{v}_{\mathbf{k}} \langle n_{\mathbf{k}-\mathbf{q}/2,\sigma} - n_{\mathbf{k}+\mathbf{q}/2,\sigma} \rangle \tag{3.12}$$

where $\mathbf{v}_{\mathbf{k}} = \nabla_{\mathbf{k}} \epsilon_{\mathbf{k}}$ is the velocity of non-interacting particles and $n_{\mathbf{k}\sigma} = a^\dagger_{\mathbf{k}\sigma} a_{\mathbf{k}\sigma}$. For models without a cutoff for single-particle momenta[8], the summation variable in (3.12) can be shifted to get

$$\mathbf{c}(\mathbf{q}) = V^{-1} \sum_{\mathbf{k}\sigma} [(\mathbf{q} \cdot \nabla_{\mathbf{k}}) \mathbf{v}_{\mathbf{k}}] \langle n_{\mathbf{k}\sigma} \rangle \tag{3.13}$$

for small $\mathbf{q}$. For continuum systems with $\epsilon_{\mathbf{k}} = \mathbf{k}^2/2m$ one obtains the simpler expression

$$\mathbf{c}(\mathbf{q}) = \frac{n}{m} \mathbf{q} \tag{3.14}$$

with the particle density $n = N/V$, which is valid for any $\mathbf{q}$.

For the charge vertex part, the continuity equation (3.3) implies the Ward identity

$$(iq_0, \mathbf{q})_\mu \Gamma^\mu_\sigma(p; q) = G^{-1}_\sigma(p + q/2) - G^{-1}_\sigma(p - q/2) \tag{3.15}$$

which can be obtained by applying a time-derivative to $\langle j^0 a a^\dagger \rangle$ in time-representation (with respect to the time-variable associated with j^0), and Fourier-transforming the resulting equation of motion. The right-hand-side in (3.15) is due to the derivative of the time-ordering $\mathcal{T}$.

The limit $q \to 0$ is related to the low-energy long-wavelength response of the system, and is thus particularly important. Hence, we define

$$\Gamma^{\mu,\mathbf{r}}_\sigma(p) = \lim_{\substack{q \to 0 \\ \mathbf{q}/q_0 = \mathbf{r}}} \Gamma^\mu_\sigma(p; q) \tag{3.16}$$

In general the limit $q = (q_0, \mathbf{q}) \to 0$ is not unique, but depends on the ratio $\mathbf{r} = \mathbf{q}/q_0$. The Ward identity for $\Gamma^\mu_\sigma(p; q)$ implies

$$\Gamma^{0,0}_\sigma(p) = \lim_{q_0 \to 0} \lim_{\mathbf{q} \to 0} \Gamma^0_\sigma(p; q) = \frac{\partial G^{-1}_\sigma(p)}{\partial(ip_0)} = 1 - \frac{\partial \Sigma_\sigma(p)}{\partial(ip_0)} \tag{3.17}$$

[8] Effective low-energy theories such a the g-ology model for one-dimensional systems may require a cutoff for single-particle momenta in order to be well defined. In this case one must take boundary terms in momentum space into account (see the section on one-dimensional systems).

and

$$\mathbf{\Gamma}^{\infty}_{\sigma}(p) = \lim_{\mathbf{q}\to 0}\lim_{q_0\to 0}\mathbf{\Gamma}_{\sigma}(p;q) = -\frac{\partial G^{-1}_{\sigma}(p)}{\partial \mathbf{p}} = \mathbf{v_p} + \frac{\partial \Sigma_{\sigma}(p)}{\partial \mathbf{p}} \tag{3.18}$$

The Feynman diagrams associated with a perturbation expansion of the vertex part $\Gamma^{\mu}_{\sigma}(p;q)$ may be one-interaction-reducible, i.e. they can be split in two pieces by cutting a single interaction line. The sum over all one-interaction-irreducible Feynman diagrams defines the *irreducible* vertex part

$$\Lambda^{\mu}_{\sigma}(p;q) = \langle j^{\mu}(q)\, a_{p-q/2,\sigma}\, a^{\dagger}_{p+q/2,\sigma}\rangle^{irr}_{tr} \tag{3.19}$$

which will play an important role below. For systems with pure density-density interactions $H_I = \frac{1}{2V}\sum_{\mathbf{q}}\, g(\mathbf{q})\,\rho(\mathbf{q})\rho(-\mathbf{q})$, i.e. for most physically relevant systems, the vertex parts Γ^{μ} and Λ^{μ} are obviously related by the Dyson equation

$$\Gamma^{\mu}_{\sigma}(p;q) = \Lambda^{\mu}_{\sigma}(p;q) + J^{\mu 0}(q)g(\mathbf{q})\Lambda^{0}_{\sigma}(p;q) \tag{3.20}$$

The Ward identities for Γ^{μ} and $J^{\mu\nu}$ imply

$$(iq_0,\mathbf{q})_{\mu}\Lambda^{\mu}_{\sigma}(p;q) = G^{-1}_{\sigma}(p+q/2) - G^{-1}_{\sigma}(p-q/2) \tag{3.21}$$

The inclusion of spin density-density interactions does not modify the above relations. A Ward identity of the form (3.21) holds also for fermions coupled to an abelian gauge-field, where Λ^{μ} is the irreducible fermion-gauge-field vertex (see Sec. 9)).

3.2. SPIN CONSERVATION

We now derive relations following from spin conservation. The logic leading to a continuity equation and Ward identities is the same as that for charge conservation. The spin-density fluctuation operator can be defined as

$$\mathbf{s}(\mathbf{q}) = \sum_{\mathbf{k}}\sum_{\sigma,\sigma'}\mathbf{s}_{\sigma\sigma'}\, a^{\dagger}_{\mathbf{k}-\mathbf{q}/2,\sigma}\, a_{\mathbf{k}+\mathbf{q}/2,\sigma'} \tag{3.22}$$

where $\mathbf{s} = (s^x, s^y, s^z) = \frac{1}{2}(\sigma^x, \sigma^y, \sigma^z)$ with the Pauli matrices σ^a, $a = x, y, z$. The total spin of the system is given by $\mathbf{S} = \mathbf{s}(\mathbf{0})$. If the Hamiltonian conserves spin, i.e. $[\mathbf{S}, H] = 0$, the equation of motion for $\mathbf{s}(\tau,\mathbf{q})$ assumes the form of a continuity equation

$$\partial_{\tau} s^a(\tau,\mathbf{q}) = -\,\mathbf{q}\cdot\mathbf{j}^a(\tau,\mathbf{q}) \tag{3.23}$$

at least for small $\mathbf{q}$. For each given system the spin-current operator $\mathbf{j}^a(\mathbf{q})$ can be calculated explicitly from the commutator $[H, s^a(\mathbf{q})]$. The decomposition $H = H_0 + H_I$ of the Hamiltonian yields a corresponding decomposition $\mathbf{j}^a = \mathbf{j}^a_0 + \mathbf{j}^a_I$ of the current operator.

The part $\mathbf{j}_I^a$ vanishes in most cases of interest (especially for pure charge density-density interactions), while $\mathbf{j}_0^a$ can be written as

$$\mathbf{j}_0^a(\mathbf{q}) = \sum_{\mathbf{k}} \sum_{\sigma,\sigma'} \nabla_{\mathbf{k}} \epsilon_{\mathbf{k}} \; s^a_{\sigma\sigma'} \; a^\dagger_{\mathbf{k}-\mathbf{q}/2,\sigma} \; a_{\mathbf{k}+\mathbf{q}/2,\sigma'} \tag{3.24}$$

For a continuum system with $\epsilon_{\mathbf{k}} = \mathbf{k}^2/2m$, one has $\nabla_{\mathbf{k}}\epsilon_{\mathbf{k}} = \mathbf{k}/m$, and the above equations hold for any $\mathbf{q}$.

Spin-density and spin-current operators are conveniently collected in a (d+1)-dimensional vector $j^{a\mu} = (s^a, \mathbf{j}^a)$, $\mu = 0, 1, \ldots, d$, which will also be referred to as (spin) "current".

The continuity equation (3.23) implies Ward identities for correlation functions involving spin-currents. In close analogy to the section on charge conservation, we consider the (spin) current-current correlation function

$$J^{a\mu,b\nu}(q) = -\tfrac{1}{V} \, \langle j^{a\mu}(q) \, j^{b\nu}(-q) \rangle \tag{3.25}$$

and the (spin) vertex part

$$\Gamma^{a\mu}_{\sigma\sigma'}(p;q) = \langle j^{a\mu}(q) \; a_{p-q/2,\sigma} \; a^\dagger_{p+q/2,\sigma'} \rangle_{tr} \tag{3.26}$$

where truncation of legs ("tr") in (3.7) amounts to dividing the expectation value by $G_\sigma(p-q/2)G_{\sigma'}(p+q/2)$.

The Ward identity for $J^{a\mu,b\nu}(q)$ reads

$$(iq_0, \mathbf{q})_\mu J^{a\mu,b\nu}(q) = \tfrac{1}{V} \, \langle [s^a(\mathbf{q}), j^{b\nu}(-\mathbf{q})] \rangle =: c^{ab\nu}(\mathbf{q}) \tag{3.27}$$

Using the commutation relations $[s^a, s^b] = i \sum_c \epsilon_{abc} \, s^c$, where ϵ_{abc} is the (totally antisymmetric) Levi-Civita tensor, one can reduce the commutator in (3.27) to

$$c^{ab\nu}(\mathbf{q}) = \frac{\delta_{ab}}{4} \, c^\nu(\mathbf{q}) + i \sum_c \epsilon_{abc} \, \tfrac{1}{V} \langle j^{c\nu}(\mathbf{0}) \rangle \tag{3.28}$$

where $c^\nu(\mathbf{q})$, given by (3.11) and (3.13), has already appeared in connection with charge conservation. Note that $\frac{1}{V}\langle s^a(\mathbf{0})\rangle$ vanishes in the absence of a total magnetization, and $\frac{1}{V}\langle \mathbf{j}^a(\mathbf{0})\rangle$ vanishes generally in equilibrium.

For the spin vertex part, the continuity equation implies the Ward identity

$$(iq_0, \mathbf{q})_\mu \Gamma^{a\mu}_{\sigma\sigma'}(p;q) = \; s^a_{\sigma\sigma'} \, [G^{-1}_{\sigma'}(p+q/2) - G^{-1}_{\sigma}(p-q/2)] \tag{3.29}$$

if spin-rotation invariance remains unbroken. In complete analogy to (3.17) and (3.18), this allows one to relate the spin vertex part for $q \to 0$ to derivatives of the self-energy with respect to frequency or momentum. In analogy to (3.19) one can define an *irreducible*

spin vertex part $\Lambda^{a\mu}_{\sigma\sigma'}(p;q)$, which, for systems with pure (charge or spin) density-density interactions also obeys a Ward identity of the form (3.29).

3.3. RENORMALIZATION OF RESPONSE FUNCTIONS

The response to small external fields, coupled to charge or spin densities and currents, belongs to the most important experimentally accessible properties of a system. The combination of renormalization group ideas and conservation laws puts powerful constraints on the structure of the effective low-energy theory for response functions.

Let us consider the response to a field $A = (\phi, \mathbf{A})$ coupled to charge density and current as an example. We consider spinless fermions to keep the number of indices small.

We couple the system to the external field A by adding a term

$$S_A[\psi, \psi^*; A] = -\int_q A_\mu(q) j^\mu(q) = -\int_q [\phi(q)\rho(q) - \mathbf{A}(q)\mathbf{j}(q)] \tag{3.30}$$

to the action, where $j^\mu(q)$ is the current operator constructed from the Grassmann variables ψ, ψ^* (instead of annihilation and creation operators), e.g. $\rho(q) = \int_k \psi^*_{k-q/2}\psi_{k+q/2}$. Adding S_A to the action in the functional integral (2.7), the generating functional $\mathcal{G}[\psi, \psi^*]$ is generalized to a functional $\mathcal{G}[\psi, \psi^*; A]$, whose functional derivatives with respect to A_μ generate expectation values involving current operators j^μ.

In most cases of interest, S_A is quadratic in the fermion fields, because $\rho(q)$ is quadratic and $\mathbf{j}(q)$ is quadratic at least for small $\mathbf{q}$. In all these cases, $j^\mu(q)$ can be written as

$$j^\mu(q) = \int_p \Gamma^\mu_0(p;q)\, \psi^*_{p-q/2}\psi_{p+q/2} \tag{3.31}$$

where $\Gamma^0_0(p;q) = 1$ for all q, and $\mathbf{\Gamma}_0(p;q) = \mathbf{v_p}$ at least for small $\mathbf{q}$. Being quadratic in ψ and ψ^*, S_A can be combined with $S_0[\psi, \psi^*]$ to give

$$S_0[\psi, \psi^*; A] = \int_{k,k'} \psi^*_k \left[\hat{\delta}(k'-k)G_0^{-1}(k) - A_\mu(k'-k)\Gamma^\mu_0[(k+k')/2; k'-k]\right] \psi_{k'} \tag{3.32}$$

where $\hat{\delta}(k'-k) = (2\pi)^{d+1}\delta(k'-k)$.

Integrating out fermionic fields ψ_k, ψ^*_k outside a Λ-shell in momentum space, one obtains an effective action $S^\Lambda[\psi, \psi^*; A]$ which depends on fields ψ_k, ψ^*_k with $d(\mathbf{k}, \partial\mathcal{F}) < \Lambda$, and on A. The effective action can be expanded in powers of ψ, ψ^* and A. Being interested in the linear response to small fields, one can drop all terms beyond quadratic order in A. Furthermore, for small Λ, monomials $(\psi^*\psi)^n A$ are irrelevant for $n \geq 2$, and monomials $(\psi^*\psi)^n A^2$ are irrelevant for all $n \geq 1$. It is easy to see that these terms would yield contributions of order Λ/k_F or smaller to the response functions. Thus, in the low-energy limit, we need only keep terms of the form A^2 and $\psi^*\psi A$ in addition to terms present

already without A. In particular, the quartic term in ψ, ψ^* is not affected by A, and it is sufficient to reconsider the quadratic part of S^Λ, which can be written as

$$S_0^\Lambda[\psi,\psi^*;A] \;=\; \int_k^{<\Lambda} \psi_k^* \,[G_0^{-1}(k) - \Sigma^\Lambda(k)]\, \psi_k \;-\; \int_q A_\mu(q) J^{\mu\nu\Lambda}(q) A_\nu(q)$$
$$- \int_{k,k'}^{<\Lambda} \psi_k^*\, A_\mu(k'-k) \Gamma^{\mu\Lambda}[(k+k')/2; k'-k]\, \psi_{k'} \;+\; \mathcal{O}(\psi^*\psi\, A^2) \tag{3.33}$$

where we have included the contribution of order A^2, generated by the mode elimination. The vertex $\Gamma^{\mu\Lambda}$ is given by the usual Feynman diagrams for the charge vertex part, defined in (3.7), where momenta on internal lines are restricted to values outside the Λ-shell. The kernel $J^{\mu\nu\Lambda}(q)$ is given by diagrams for the current-current correlator, defined in (3.6), again with internal momenta outside the Λ-shell. Expanding the self-energy Σ^Λ and introducing renormalized fields $\bar\psi_k = \psi_k/(Z^\Lambda_{\mathbf{k}_F})^{1/2}$, one obtains the renormalized action

$$\bar S_0^\Lambda[\bar\psi,\bar\psi^*;A] \;=\; \int_k^{<\Lambda} \bar\psi_k^* \,[ik_0 - \bar v^\Lambda_{\mathbf{k}_F} k_r - \bar\Sigma^\Lambda(k)]\, \bar\psi_k \;-\; \int_q A_\mu(q) J^{\mu\nu\Lambda}(q) A_\nu(q)$$
$$- \int_{k,k'}^{<\Lambda} \bar\psi_k^*\, A_\mu(k'-k)\, (Z^\Lambda_{\mathbf{k}_F} Z^\Lambda_{\mathbf{k}'_F})^{1/2}\, \Gamma^{\mu\Lambda}[(k+k')/2; k'-k]\, \bar\psi_{k'} \tag{3.34}$$

We have not assumed isotropy here, i.e. the field renormalization factor $Z^\Lambda_{\mathbf{k}_F}$ and the Fermi velocity $\bar v^\Lambda_{\mathbf{k}_F}$ may depend on $\mathbf{k}_F$.

So far, all steps are valid for arbitrary q, provided that $\Lambda \ll k_F$. We now consider the limit $q \to 0$, i.e. the response to fields with a low frequency and a small wave number. There are two simplifications in that limit, reducing considerably the number of unknown terms in the effective low-energy action.

Firstly, $J^{\mu\nu\Lambda}(q)$ has a *unique* limit $J^{\mu\nu\Lambda}(0) \equiv J^{\mu\nu\Lambda}_{inc}$ for $q \to 0$ (independent of the ratio $\mathbf{q}/q_0$) at $\Lambda > 0$, determined by "incoherent" contributions. Without cutoff $J^{\mu\nu}(q)$ generally has no unique limit for $q \to 0$ due to singular contributions of "coherent" particle-hole excitations with arbitrarily small momentum transfer $\mathbf{q}$ (as is well-known from Fermi liquid theory, see Sec. 4). However, particle-hole excitations across the Λ-shell are not possible, if $|\mathbf{q}|$ is smaller that the width of the shell 2Λ. The Ward identity (3.10) then implies that $J^{00\Lambda}_{inc} = 0$ and $J^{0j\Lambda}_{inc} = J^{j0\Lambda}_{inc} = 0$. Note that the Ward identities derived in 3.1 and 3.2 hold also for $J^{\mu\nu\Lambda}(q)$, $\Gamma^{\mu\Lambda}$, G^Λ and other correlation functions with a cutoff $\Lambda > 0$.[9] In fact these quantities can be viewed as exact correlation functions in a theory where the interaction does not scatter particles inside the Λ-shell (intermediate states corresponding to internal lines in Feynman diagrams inside the Λ-shell are not accessible).

[9]The expression (3.12) for the function $\mathbf{c}(\mathbf{q})$ has to be modified to $\mathbf{c}^\Lambda(\mathbf{q}) := V^{-1}\sum_{\mathbf{k}\sigma}\, \mathbf{v}_{\mathbf{k}}\, \big[\Xi^{>\Lambda}(\mathbf{k}+\mathbf{q}/2)\langle n_{\mathbf{k}-\mathbf{q}/2,\sigma}\rangle - \Xi^{>\Lambda}(\mathbf{k}-\mathbf{q}/2)\langle n_{\mathbf{k}+\mathbf{q}/2,\sigma}\rangle\big]$ if states outside the Λ-shell are excluded, where $\Xi^{>\Lambda}$ is the characteristic function of the allowed momenta.

The second simplification is due to the Ward identity (3.15) for the vertex part Γ^μ. Again, the reason for the non-uniqueness of the limit $q \to 0$ of $\Gamma^\mu(p;q)$, i.e. particle-hole excitations with arbitrarily small momentum transfer $\mathbf{q}$, is eliminated for any $\Lambda > 0$. Hence, the Ward identity (3.15) implies

$$\lim_{q\to 0} \Gamma^{0\Lambda}(p;q) = 1 - \frac{\partial \Sigma^\Lambda(p)}{\partial(ip_0)} \;\longrightarrow\; 1/Z^\Lambda_{\mathbf{p}_F} \tag{3.35}$$

and

$$\lim_{q\to 0} \mathbf{\Gamma}^{\Lambda}(p;q) = \mathbf{v}_{\mathbf{p}} + \frac{\partial \Sigma^\Lambda(p)}{\partial \mathbf{p}} \;\longrightarrow\; \bar{\mathbf{v}}^{c\Lambda}_{\mathbf{p}_F}/Z^\Lambda_{\mathbf{p}_F} \tag{3.36}$$

where the latter limit $p \to (0, \mathbf{p}_F)$ is obtained from the expressions for $Z^\Lambda_{\mathbf{p}_F}$ and $\bar{\mathbf{v}}^{c\Lambda}_{\mathbf{p}_F}$ in terms of Σ^Λ (see (2.24) and (2.26)). Inserting these relations in (3.34), we obtain the following simple result for the quadratic part of the effective low-energy action

$$\bar{S}^\Lambda_0[\bar\psi, \bar\psi^*; A] = \int_k^{<\Lambda} \bar\psi^*_k [ik_0 - \bar v^\Lambda_{\mathbf{k}} k_r - \bar\Sigma^\Lambda(k)] \bar\psi_k - \int_q A_\mu(q) \bar j^{\mu\Lambda}(q) - \int_q A_j(q) J^{jj'\Lambda}_{inc} A_{j'}(q) \tag{3.37}$$

with a renormalized current operator

$$\bar j^{\mu\Lambda}(q) = \int_k^{<\Lambda} (1, \bar{\mathbf{v}}^{c\Lambda}_{\mathbf{k}})^\mu \; \bar\psi^*_{k-q/2} \bar\psi_{k+q/2} \tag{3.38}$$

Note that the *field renormalizations have cancelled out*. We emphasize that (3.37) is valid only for fields $A_\mu(q)$ with small q, and Λ chosen such that $2|\mathbf{q}| < \Lambda \ll k_F$!

In a Fermi liquid, which will be discussed in some detail in Sec. 4, $\bar j^{\mu\Lambda}(q)$ is the quasiparticle current (including backflow), and $\bar S^\Lambda_0[\bar\psi, \bar\psi^*; A]$, supplemented by residual quasiparticle interactions, is nothing but the Landau model for the response of an interacting Fermi system to weak and slowly varying external fields [2]. To describe the response of charged fermions to electromagnetic fields A^μ one must of course add the diamagnetic term ($\propto \int \rho \mathbf{A}^2$) to the microscopic action.

4. FERMI LIQUIDS

In this section we describe a renormalization group picture of microscopic Fermi liquid theory. Renormalization (group) ideas and language have been used to a certain extent already in many early works on Fermi liquid theory. The idea to derive a systematic low-energy expansion by integrating out high-energy degrees of freedom had been worked out for normal and superfluid Fermi systems in the 1980's [77]. As a subject in its own right the rephrasing of Fermi liquid theory in modern renormalization group language has attracted interest only quite recently [19, 82, 93]. Besides providing a faster way of understanding many aspects of Fermi liquid theory (without adding new results) the renormalization group is also a powerful tool for obtaining rigorous results on the Fermi liquid behavior of specific systems.[10]

The following presentation of Fermi liquid theory from a renormalization group point of view absorbs many ideas from the above-mentioned earlier works on this subject. We have clarified several points that we found so far either unclear or incomplete, especially concerning the flow of coupling functions and the derivation of response functions in Fermi liquids. We consider all orders in perturbation theory, but do not address the issue of convergence and non-perturbative effects [17, 34]. For pedagogical reasons we consider only spinless fermions with short-range interactions. Including spin or (sufficiently screened) long-range interactions is not hard.

4.1. MICROSCOPIC FERMI LIQUID THEORY:

A Fermi liquid can be characterized by the following asymptotic behavior of one- and two-particle Green (or vertex) functions in the low-energy limit $k_0 \to 0$ at $T = 0$ [2, 14].

The single-particle propagator behaves as

$$G(k) = \frac{Z_{\mathbf{k}_F}}{ik_0 - v^*_{\mathbf{k}_F} k_r - \bar{\Sigma}(k)} \tag{4.1}$$

where $\bar{\Sigma}(k)$ vanishes faster than linearly for $k \to (0, \mathbf{k}_F)$. Here k_r measures the distance of $\mathbf{k}$ from the Fermi surface of the interacting system, which is defined by the position

[10] Significant rigorous results on the Fermi liquid behavior of interacting Fermi systems have been obtained very recently by Feldman et al. [17, 34]. In particular, it has been shown that two-dimensional Fermi systems without Cooper instability exhibit a finite discontinuity in the momentum distribution function within a finite radius of convergence in coupling space.

of the pole in $G(k)$. This behavior implies the existence of *Landau quasi-particles*, i.e. gapless fermionic single-particle excitations with a linear dispersion relation. The *spectral weight* $Z_{\mathbf{k}_F} \in]0,1]$ of these excitations can be obtained from the self-energy as

$$Z_{\mathbf{k}_F} = \left[1 - \partial\Sigma(k_0,\mathbf{k})/\partial(ik_0)\right]^{-1}\Big|_{(0,\mathbf{k}_F)} \tag{4.2}$$

and the quasi-particle or Fermi *velocity* is given by

$$\mathbf{v}^*_{\mathbf{k}_F} = Z_{\mathbf{k}_F}\left[\mathbf{v}_{\mathbf{k}} + \partial\Sigma(k_0,\mathbf{k})/\partial\mathbf{k}\right]\big|_{(0,\mathbf{k}_F)} \tag{4.3}$$

where $\mathbf{v}_{\mathbf{k}} = \nabla\epsilon_{\mathbf{k}}$ is the bare velocity in the absence of interactions. The momentum distribution function in the ground state has a finite discontinuity $Z_{\mathbf{k}_F}$ across the Fermi surface. For isotropic systems $Z_{\mathbf{k}_F} = Z$ and $v^*_{\mathbf{k}_F} = v^*_F$ are constant all over the Fermi surface.

The two-particle vertex $\Gamma(k_1,k_2;k_1',k_2')$ is finite in the limit $k_i \to (0,\mathbf{k}_{Fi})$, $k_i' \to (0,\mathbf{k}_{Fi}')$.[11] For $\mathbf{k}_{F1}' \neq \mathbf{k}_{F1},\mathbf{k}_{F2}$ this limit is unique and defines the quasi-particle *scattering amplitudes*

$$a(\mathbf{k}_{F1},\mathbf{k}_{F2};\mathbf{k}_{F1}',\mathbf{k}_{F2}') = [Z_{\mathbf{k}_{F1}}Z_{\mathbf{k}_{F2}}Z_{\mathbf{k}_{F1}'}Z_{\mathbf{k}_{F2}'}]^{1/2}\lim_{k_i\to(0,\mathbf{k}_{Fi})}\lim_{k_i'\to(0,\mathbf{k}_{Fi}')}\Gamma(k_1,k_2;k_1',k_2') \tag{4.4}$$

For $\mathbf{k}_{F1}' = \mathbf{k}_{F1}$ or $\mathbf{k}_{F1}' = \mathbf{k}_{F2}$ the above limit is not unique. Due to the antisymmetry of the vertex function under exchange of the outgoing (or incoming) particles it is sufficient to analyze one of these cases. Consider, for example, the vertex function for $k_1' \sim k_1 \sim (0,\mathbf{k}_F)$ and $k_2' \sim k_2 \sim (0,\mathbf{k}_F')$. The non-uniqueness of the limit $q = k_1' - k_1 \to 0$ is due to particle-hole pairs with momentum q as in the second diagram on the right hand side of Fig. 4.1, where the irreducible vertices Γ_{irr} are defined by excluding subdiagrams with such particle-hole pairs. The Feynman diagrams contributing to the irreducible vertex Γ_{irr} are regular in the limit $q \to 0$. The product of propagators $G(p-q/2)G(p+q/2)$ in Fig. 4.1 can be effectively replaced by [2, 14]

$$2\pi\delta(p_0)\, Z^2_{\mathbf{p}_F}\,\frac{\Theta(\xi^*_{\mathbf{p}+\mathbf{q}/2}) - \Theta(\xi^*_{\mathbf{p}-\mathbf{q}/2})}{iq_0 - \mathbf{v}^*_{\mathbf{p}_F}\cdot\mathbf{q}} \;+\; \phi(p) \tag{4.5}$$

where $\xi^*_{\mathbf{k}} = v^*_F k_r$ and $\phi(p)$ is a regular function of p. The first term in (4.5) vanishes for $\mathbf{q} \to \mathbf{0}$ at finite q_0 while it yields finite contributions for $q \to 0$ with $|\mathbf{q}|/q_0 \neq 0$. Two special ways of taking the limit $q \to 0$, i.e.

$$\Gamma^0(\mathbf{k}_F,\mathbf{k}_F') = \lim_{q_0\to 0}\lim_{\mathbf{q}\to\mathbf{0}}\Gamma(\mathbf{k}_F,\mathbf{k}_F';q) \tag{4.6a}$$

[11] Note our notation $\tilde{\Gamma}(k_1,k_2;k_1',k_2') = (2\pi)^{d+1}\delta(k_1'+k_2'-k_1-k_2)\,\Gamma(k_1,k_2;k_1',k_2')$ and $\Gamma(k,k';q) = \Gamma(k-q/2,k'+q/2;k+q/2,k'-q/2)$.

$$\Gamma^{\infty}(\mathbf{k}_F, \mathbf{k}'_F) = \lim_{\mathbf{q}\to\mathbf{0}} \lim_{q_0\to 0} \Gamma(\mathbf{k}_F, \mathbf{k}'_F; q) \tag{4.6b}$$

play a prominent role in Fermi liquid theory (the notation refers to a more general limit $\Gamma^{\mathbf{r}}$ where $\mathbf{q}/q_0 \to \mathbf{r}$ as $q \to 0$). The first one defines Landau's quasi-particle *interaction* $f_{\mathbf{k}_F\mathbf{k}'_F}$ as

$$Z_{\mathbf{k}_F} Z_{\mathbf{k}'_F} \Gamma^0(\mathbf{k}_F, \mathbf{k}'_F) = f_{\mathbf{k}_F\mathbf{k}'_F} \tag{4.7}$$

and the second the *forward scattering amplitude* $a(\mathbf{k}_F, \mathbf{k}'_F)$ via

$$Z_{\mathbf{k}_F} Z_{\mathbf{k}'_F} \Gamma^{\infty}(\mathbf{k}_F, \mathbf{k}'_F) = a(\mathbf{k}_F, \mathbf{k}'_F) = \lim_{\mathbf{k}_{F1},\mathbf{k}'_{F1}\to\mathbf{k}_F} \lim_{\mathbf{k}_{F2},\mathbf{k}'_{F2}\to\mathbf{k}'_F} a(\mathbf{k}_{F1}, \mathbf{k}_{F2}; \mathbf{k}'_{F1}, \mathbf{k}'_{F2}) \tag{4.8}$$

For small q and k, k' close to the Fermi surface, the vertex function $\Gamma(k, k'; q)$ can be reconstucted from the quasi-particle interaction by solving the linear integral equation

$$\bar{\Gamma}(\mathbf{k}_F, \mathbf{k}'_F; q) = f_{\mathbf{k}_F\mathbf{k}'_F} + \int_{\mathbf{k}''_F} f_{\mathbf{k}_F\mathbf{k}''_F} \, \frac{\mathbf{n}_{\mathbf{k}''_F} \cdot \mathbf{q}}{iq_0 - \mathbf{v}^*_{\mathbf{k}''_F} \cdot \mathbf{q}} \, \bar{\Gamma}(\mathbf{k}''_F, \mathbf{k}'_F; q) \tag{4.9}$$

for the renormalized vertex $\bar{\Gamma}(\mathbf{k}_F, \mathbf{k}'_F; q) = Z_{\mathbf{k}_F} Z_{\mathbf{k}'_F} \Gamma(\mathbf{k}_F, \mathbf{k}'_F; q)$, where the integral extends over the Fermi surface.[12]

For isotropic systems, $f_{\mathbf{k}_F\mathbf{k}'_F}$ and $a(\mathbf{k}_F, \mathbf{k}'_F)$ depend only on the angle θ between $\mathbf{k}_F$ and $\mathbf{k}'_F$. Multiplying by the density of quasi-particle excitations at the Fermi level, N^*_F, one obtains dimensionless functions $F(\theta) = N^*_F \, f(\theta)$ and $A(\theta) = N^*_F \, a(\theta)$. Expanding in spherical harmonics, F and A can be expressed in terms of a discrete set of Landau parameters F_l and A_l, respectively, where l labels the various harmonics.

4.2. EFFECTIVE ACTION AND FIXED POINT

The low-energy effective action of a (possibly anisotropic) Fermi system has the form

$$\begin{aligned} \bar{S}^{\Lambda}[\bar{\psi}, \bar{\psi}^*] = \int_k^{<\Lambda} \bar{\psi}^*_k [ik_0 - \bar{v}^{\Lambda}_{\mathbf{k}_F} k_r - \bar{\Sigma}^{\Lambda}(k)] \bar{\psi}_k - \int_k^{<\Lambda} (\delta\bar{\mu}^{\Lambda}_{\mathbf{k}_F} - \delta\bar{v}^{\Lambda}_{\mathbf{k}_F} k_r) \, \bar{\psi}^*_k \bar{\psi}_k \\ - \tfrac{1}{4} \int_{k,k',q}^{<\Lambda} \bar{\Gamma}^{\Lambda}(k, k'; q) \, \bar{\psi}^*_{k-q/2} \bar{\psi}^*_{k'+q/2} \bar{\psi}_{k'-q/2} \bar{\psi}_{k+q/2} \end{aligned} \tag{4.10}$$

for $\Lambda \ll k_F$, provided there are no instabilities with an energy scale outside the Λ-shell (cf. Sec. 2). More-than-two-particle interactions are irrelevant for the low-energy behavior and have already been omitted. The *residual* self-energy $\bar{\Sigma}^{\Lambda}(k)$ vanishes faster than linearly for $k_0, k_r \to 0$ (linear terms have been absorbed in $Z^{\Lambda}_{\mathbf{k}_F}$ and $\bar{v}^{\Lambda}_{\mathbf{k}_F}$).

[12] The notation $\int_{\mathbf{k}_F} \ldots$ is an abbreviation for $(2\pi)^{-d} \int_{\partial\mathcal{F}} dS(\mathbf{k}_F) \ldots$, where $dS(\mathbf{k}_F)$ is the infinitesimal surface element at $\mathbf{k}_F$.

In a Fermi liquid, *all renormalizations are finite* in the limit $\Lambda \to 0$. The field renormalization tends to a finite constant

$$Z^{\Lambda}_{\mathbf{k}_F} \to Z_{\mathbf{k}_F} \in]0,1] \tag{4.11}$$

which is easily identified as the quasi-particle weight. The renormalized Fermi velocity converges to the quasi-particle velocity

$$\bar{v}^{\Lambda}_{\mathbf{k}_F} \to v^*_{\mathbf{k}_F} > 0 \tag{4.12}$$

The counterterm $-\delta\bar{\mu}^{\Lambda}_{\mathbf{k}_F}$ had been introduced to compensate shifts of the Fermi surface induced by the interaction. For $\Lambda \ll k_F$ these shifts are of order Λ, generated by tadpole diagrams (other diagrams lead to shifts of order Λ^2). Hence, this counterterm can be omitted from the effective action, if the latter is normal ordered (with respect to the Fermi sea). Normal ordering also compensates the velocity counterterm $\delta\bar{v}^{\Lambda}_{\mathbf{k}_F} k_r$.

The behavior of the vertex function in the limit $\Lambda \to 0$ is quite subtle. According to (4.4), one has

$$\bar{\Gamma}^{\Lambda}(\mathbf{k}_{F1}, \mathbf{k}_{F2}; \mathbf{k}'_{F1}, \mathbf{k}'_{F2}) \;\to\; a(\mathbf{k}_{F1}, \mathbf{k}_{F2}; \mathbf{k}'_{F1}, \mathbf{k}'_{F2}) \quad \text{for} \quad \mathbf{k}'_{F1} \neq \mathbf{k}_{F1}, \mathbf{k}_{F2} \tag{4.13}$$

On the other hand, if $\mathbf{k}'_{F1}$ coincides with $\mathbf{k}_{F1}$ or $\mathbf{k}_{F2}$, $\bar{\Gamma}^{\Lambda}$ does not tend to the scattering amplitude a, but to the Landau function f:

$$\bar{\Gamma}^{\Lambda}(\mathbf{k}_F, \mathbf{k}'_F; \mathbf{k}_F, \mathbf{k}'_F) = -\bar{\Gamma}^{\Lambda}(\mathbf{k}_F, \mathbf{k}'_F; \mathbf{k}'_F, \mathbf{k}_F) \;\to\; f_{\mathbf{k}_F \mathbf{k}'_F} \tag{4.14}$$

Note that the limits $\Lambda \to 0$ and $\mathbf{k}'_{F1} \to \mathbf{k}_{F1}$ (or $\mathbf{k}_{F2}$) do not commute![13] This singularity has the same origin as the non-uniqueness of the limit $q \to 0$ discussed already in 4.1. To see that, consider the vertex function $\bar{\Gamma}^{\Lambda}(k, k'; q)$ for $k \sim (0, \mathbf{k}_F)$, $k' \sim (0, \mathbf{k}'_F)$ and $q \sim 0$. For $|\mathbf{q}| < 2\Lambda$, particle-hole pairs of propagators as in Fig. 4.1 do not yield any singular contributions to $\bar{\Gamma}^{\Lambda}(k, k'; q)$, because there are no particle-hole excitations across the Λ-shell with $|\mathbf{q}| < 2\Lambda$. Hence, for $\Lambda \ll k_F$ and k, k' close to the Fermi surface (and energy) one has $\bar{\Gamma}^{\Lambda}(k, k'; q) \sim f_{\mathbf{k}_F \mathbf{k}'_F}$ if $|\mathbf{q}| < 2\Lambda$.

The flow of the vertex (or coupling) function is obviously singular for $\mathbf{k}'_{F1} \sim \mathbf{k}_{F1}$ (or $\mathbf{k}_{F2}$). For $|\mathbf{k}'_{F1} - \mathbf{k}_{F1}| < 2\Lambda$, particle-hole pairs do not yield any singular contribution to $\bar{\Gamma}^{\Lambda}$, which therefore seems to flow towards Landau's f-function. But then, reducing 2Λ below $|\mathbf{k}'_{F1} - \mathbf{k}_{F1}|$, the coupling function undergoes a rapid flow towards $a(\mathbf{k}_{F1}, \mathbf{k}_{F2}; \mathbf{k}'_{F1}, \mathbf{k}'_{F2}) \sim$

[13] This problem has led to some confusion in previous works on RG and Fermi liquid theory; for example, in Ref. [19] it seems that the "fixed point coupling $u(\mathbf{k}_{F1}, \mathbf{k}_{F2}; \mathbf{k}'_{F1}, \mathbf{k}'_{F2})$", obtained in the limit $\Lambda \to 0$, converges to the Landau-function in the forward scattering limit, in contrast to (4.8).

$a(\mathbf{k}_{F1}, \mathbf{k}_{F2})$, i.e. the β-functional is strongly Λ-dependent and singular for $|\mathbf{k}'_{F1} - \mathbf{k}_{F1}| \sim \Lambda \ll k_F$! Fixed points are reached only pointwise on the Fermi surface, but not uniformly.

Assuming the above convergence properties, one can express the low-energy behavior of the system in terms of $Z_{\mathbf{k}_F}$, $v^*_{\mathbf{k}_F}$ and a (or f) by expressing physical quantities in terms of $\bar{S}^\Lambda$ and expanding in powers of small energy scales (T, k_0 etc.) and Λ/k_F. The leading and in some cases also the first subleading low-energy behavior can be obtained from a small (i.e. tractable) set of Feynman diagrams [19, 77].

As observed already by Landau, residual scattering processes with finite momentum transfers (i.e. non-forward scattering processes) do not contribute to the *leading* low-energy behavior of thermodynamics, single-particle excitations and long-wavelength response functions (due to the restriction of ingoing and outgoing momenta to the thin Λ-shell). To this leading order one may therefore effectively drop the interactions with momentum transfers $|\mathbf{q}| > 2\Lambda$ from the action $\bar{S}^\Lambda$. In the limit $\Lambda \ll k_F$ one then obtains a scale-invariant or *fixed point* action

$$\bar{S}^*[\bar{\psi}, \bar{\psi}^*] = \int_k^{<\Lambda} \bar{\psi}^*_k (ik_0 - v^*_{\mathbf{k}_F} k_r) \bar{\psi}_k - \tfrac{1}{2} \int_{k,k',q}^{<\Lambda} f_{\mathbf{k}_F \mathbf{k}'_F} \, \bar{\psi}^*_{k-q/2} \bar{\psi}^*_{k'+q/2} \bar{\psi}_{k'-q/2} \bar{\psi}_{k+q/2} \qquad (4.15)$$

where, in addition to the restriction of fermionic momenta to the Λ-shell, the momentum transfers $\mathbf{q}$ are now restricted by the *condition* $|\mathbf{q}| < 2\Lambda$! The prefactor $\frac{1}{2} = \frac{2}{4}$ in the interaction part is the result of a summation of interaction terms with $k'_1 \sim k_1$ and their conjugate under exchange, $k'_1 \sim k_2$. Solving the action $\bar{S}^*[\bar{\psi}, \bar{\psi}^*]$ by Hartree mean-field theory (neglecting fluctuations) leads to Landau's energy functional

$$\delta E[\delta \bar{n}] = \int_{\mathbf{k}} v^*_{\mathbf{k}_F} k_r \, \delta \bar{n}_{\mathbf{k}} + \tfrac{1}{2} \int_{\mathbf{k},\mathbf{k}'} f_{\mathbf{k}_F \mathbf{k}'_F} \, \delta \bar{n}_{\mathbf{k}} \delta \bar{n}_{\mathbf{k}'} \qquad (4.16)$$

where $\delta \bar{n}_{\mathbf{k}}$ is the quasi-particle distribution function. The renormalized single-particle propagator is then determined by the quadratic part (only) of the fixed point action $\bar{S}^*[\bar{\psi}, \bar{\psi}^*]$ as

$$\bar{G}(k) \sim \bar{G}^*_0(k) = \frac{1}{ik_0 - v^*_{\mathbf{k}_F} k_r} \qquad (4.17)$$

This is the quasi-particle propagator, which describes stable quasi-particle excitations with a velocity $v^*_{\mathbf{k}_F}$.

For three-dimensional systems the irrelevance of fluctuations and the ensuing irrelevance of residual interactions for the single particle propagator have been already established by Landau's estimate of the quasi-particle decay-rate. A detailed analysis of the action (4.15) in Sec. 7 will show that these results hold actually in any dimension $d > 1$.

4.3. RESPONSE FUNCTIONS

We now derive the low-energy and long-wavelength density and current response of a Fermi liquid. Adding the renormalized source term from subsection 3.3 to the fixed point action $\bar{S}^*[\bar{\psi},\bar{\psi}^*]$, we immediately find that the linear response functions for perturbations coupling to the density or current in a Fermi liquid are described by the effective action

$$\bar{S}^*[\bar{\psi},\bar{\psi}^*,A] = \bar{S}^*[\bar{\psi},\bar{\psi}^*] - \int_q A_\mu(q)\,\bar{j}^\mu(q) - \int_q A_j(q)\,J^{*jj'}_{inc}\,A_{j'}(q) \tag{4.18}$$

with a *quasi-particle current*

$$\bar{j}^\mu(q) = \int_k^{<\Lambda} (1, \mathbf{v}^{c*}_{\mathbf{k}_F})^\mu\ \bar{\psi}^*_{k-q/2}\bar{\psi}_{k+q/2} \tag{4.19}$$

for $|\mathbf{q}| \ll k_F$ and small q_0, where $\mathbf{v}^{c*}_{\mathbf{k}_F} = \lim_{\Lambda\to 0}\bar{\mathbf{v}}^{c\Lambda}_{\mathbf{k}_F}$ and $J^{*jj'}_{inc} = \lim_{\Lambda\to 0} J^{jj'\Lambda}_{inc}$. We have thus obtained a relatively short microscopic justification of the *Landau model* for the response of a Fermi system to low-energy long-wavelength perturbations.[14] Note that the source field $A_\mu(q)$ remains unrenormalized, as a consequence of charge conservation. Consequently *no renormalization factors appear in the (long wavelength) response functions.* For Galilean invariant systems there is another simplification: $J^{*jj'}_{inc}$ vanishes.

Powercounting [19, 77] shows that a solution of $\bar{S}^*[\psi,\psi^*,A]$ within the random phase approximation (RPA) yields the exact leading low-energy behavior of response functions (expressed in terms of fixed point parameters). The RPA effective interaction D is equal to the renormalized vertex function for small q, i.e.

$$D_{\mathbf{k}\mathbf{k}'}(q) = \bar{\Gamma}(\mathbf{k}_F,\mathbf{k}'_F;q) \tag{4.20}$$

and can be obtained from $f_{\mathbf{k}_F\mathbf{k}'_F}$ by solving the integral equation (4.9). The current-current response function $J^{\mu\nu}(q)$ (including the density-density response as J^{00}) is then given by

$$\begin{aligned} J^{\mu\nu}(q) = J^{*\mu\nu}_{inc} + \bar{J}^{\mu\nu}_0(q) \ + \ \int_{k,k'} &\lambda^\mu(\mathbf{k})\,\bar{G}^*_0(k-q/2)\bar{G}^*_0(k+q/2) \\ &\times D_{\mathbf{k}\mathbf{k}'}(q)\,\bar{G}^*_0(k'-q/2)\bar{G}^*_0(k'+q/2)\,\lambda^\nu(\mathbf{k}') \end{aligned} \tag{4.21}$$

where $\lambda^\mu(\mathbf{k}) = (1,\mathbf{v}^{c*}_{\mathbf{k}_F})^\mu$ and

$$\bar{J}^{\mu\nu}_0(q) = \int_k \lambda^\mu(\mathbf{k})\,\bar{G}^*_0(k-q/2)\bar{G}^*_0(k+q/2)\,\lambda^\nu(\mathbf{k}) \tag{4.22}$$

We recall from Sec. 3.3 that $J^{*\mu\nu}_{inc}$ vanishes, if either μ or ν are zero.

[14] It is instructive to compare with the derivation in Nozières' book [2].

There is no explicit solution of the RPA integral equation for general $f_{\mathbf{k}_F\mathbf{k}'_F}$, but various important generic properties of $J^{\mu\nu}(q)$ have been established [2, 94]. Among those is the existence of propagating collective modes, known as *zero sound* [1]. The dynamical charge conductivity $\sigma(\omega)$ (related to the current-current response by the Kubo formula), has a δ-peak at zero frequency (at $T = 0$), implying a vanishing charge DC-resistivity. A simple expression can be obtained for the compressibility $\kappa = dn/d\mu$, which is related to $J^{00}(q)$ via the limit

$$\kappa = -\lim_{\mathbf{q}\to\mathbf{0}} \lim_{q_0\to 0} J^{00}(q) \tag{4.23}$$

In an isotropic system, κ depends on residual interactions only via the Landau parameter F_0:

$$\kappa = \frac{N_F^*}{1+F_0} \tag{4.24}$$

For further implications of (4.21) the reader is referred to any textbook on Fermi liquid theory [2, 94].

4.4. SUBLEADING CORRECTIONS

So far we have discussed the *leading* low-energy behavior, which was completely determined by the fixed point action $\bar{S}^*$ in (4.15) and (4.18). To this leading order, very simple *scaling* relations hold for all quantities, such as

$$G_{\mathbf{k}_F}(sk_0, sk_r) = s^{-1}\, G_{\mathbf{k}_F}(k_0, k_r) \quad \text{and} \quad J^{\mu\nu}(sq_0, s\mathbf{q}) = J^{\mu\nu}(q_0, \mathbf{q}) \tag{4.25}$$

Here $G_{\mathbf{k}_F}(k_0, k_r) = G(k_0, \mathbf{k})$ where $\mathbf{k}_F$ is a projection of $\mathbf{k}$ on the Fermi surface.

At small but finite energy scales, there are *subleading* corrections to this simple behavior. For example, the single-particle propagator has the form (see (4.1))

$$G(k) = \frac{Z_{\mathbf{k}_F}}{ik_0 - v^*_{\mathbf{k}_F} k_r - \bar{\Sigma}(k)} \tag{4.26}$$

with a residual self-energy $\bar{\Sigma}(k)$ of order $\mathcal{O}(k_0^2, k_r^2)$ in three dimensions, describing quasi-particle decay and a small incoherent background (with a spectral weight that vanishes quadratically in the low-energy limit). For real frequencies ξ close to the quasi-particle pole, one thus obtains

$$G(k) = \frac{Z_{\mathbf{k}_F}}{\xi - \xi^*_{\mathbf{k}} \pm i\gamma_{\mathbf{k}}} \quad \text{for} \quad \xi^*_{\mathbf{k}} \gtrless 0 \tag{4.27}$$

with a quasi-particle dispersion relation $\xi^*_{\mathbf{k}} = v^*_{\mathbf{k}_F} k_r + \mathrm{Re}\bar{\Sigma}(\xi^*_{\mathbf{k}}, \mathbf{k})$ and a decay rate $\gamma_{\mathbf{k}} = |\mathrm{Im}\bar{\Sigma}(\xi^*_{\mathbf{k}}, \mathbf{k})|$ of order k_r^2.

These subleading corrections are partially due to scattering processes with arbitrary momentum transfers, generated by the quasi-particle scattering amplitude a, and also

due to irrelevant terms contained in $\bar{\Sigma}^{\Lambda}$, which had been discarded in $\bar{S}^{*}$. Sometimes subleading corrections can be expressed exactly in terms of fixed point parameters. In particular, in three-dimensional systems the decay rate $\gamma_{\mathbf{k}}$ is given by Fermi's Golden Rule with the scattering amplitude a as matrix elements [94]. This corresponds to a single Feynman diagram for the self-energy [77], i.e. diagram (b) in Fig. 2.3.[15]

Subleading corrections due to residual scattering processes are also responsible for the damping of collective modes and, in particular, for a finite (i.e. non-zero) resistivity $\rho(T)$ at finite temperature (if not already made finite by impurities). Umklapp processes in metals are usually an "irrelevant" perturbation (except for special Fermi surface geometries) in the sense that they die out in the zero energy limit. Nevertheless they are very important at any finite temperature, where, in the absence of impurities and phonons, they are the only way to make the DC resistivity finite. In conventional three-dimensional metals these terms lead to a contribution

$$\rho_u(T) \propto T^2 \tag{4.28}$$

to the electrical DC resistivity [96].

4.5. INSTABILITIES

There are several consistency conditions that a Fermi liquid must satisfy for stability. Two types of instabilities are generic in the sense that they do not require low dimensionality or special Fermi surface geometries: The Pomerantchuk instability and the Cooper instability.

The *Pomerantchuk instability* [97] occurs when the energy of the (putative) Fermi liquid can be lowered by deforming the Fermi surface. This can happen if the Landau function $f_{\mathbf{k}_F\mathbf{k}'_F}$ exceeds certain limits. For three-dimensional isotropic systems, a Pomerantschuk instability occurs if $F_l < -(2l+1)$ for any of the harmonics l [2]. In particular, $F_0 < -1$ would imply a negative compressibility, i.e. an instability with respect to phase separation. A stable isotropic Fermi liquid must therefore satisfy the condition

$$F_l > -(2l+1) \quad \text{for all } l \tag{4.29}$$

The *Cooper instability* [98] occurs when the system can lower its energy by forming bound pairs in the presence of effective attractive interactions. A perturbative signal for this is a run-away flow of the vertex function in the Cooper channel $\bar{g}_C^{\Lambda}(\mathbf{k}_F, \mathbf{k}'_F) = \bar{\Gamma}^{\Lambda}(\mathbf{k}_F, -\mathbf{k}_F; \mathbf{k}'_F, -\mathbf{k}'_F)$. As discussed already in Sec. 2, in isotropic systems this happens if at least one of the partial wave amplitudes $\bar{g}_l^{\Lambda}$ becomes negative. Since the β-function

[15] In two dimensions, more diagrams are needed [95].

for $\bar{g}_C^\Lambda(\mathbf{k}_F, \mathbf{k}'_F)$ vanishes only if $\bar{g}_C^\Lambda(\mathbf{k}_F, \mathbf{k}'_F)$ itself vanishes, the scattering amplitude in a stable Fermi liquid must satisfy the condition

$$a(\mathbf{k}_F, -\mathbf{k}_F; \mathbf{k}'_F, -\mathbf{k}'_F) = 0 \tag{4.30}$$

According to an argument by Kohn and Luttinger [87], all interacting Fermi systems will undergo a Cooper instability (if no other) at sufficiently low energy scales, even if the bare interactions are purely repulsive. Their intuitive argument has been substantiated by systematic low-density expansions [89, 90, 91], as well as by recent explicit calculations of the Cooper flow with the inclusion of irrelevant and higher (than second) orders in perturbation theory [19, 88]. In real systems the energy scale for the Kohn-Luttinger instability seems to be extremely small and thus far below the scale where Fermi liquid low-energy behavior or other (than Kohn-Luttinger) symmetry breaking mechanisms set in.

5. FORWARD SCATTERING

In many systems the low-energy physics is dominated by forward scattering, i.e. scattering processes with small momentum transfers $\mathbf{q}$. Forward scattering obviously plays a prominant role in systems with long-range interactions, where the scattering amplitudes diverge for small momentum transfers. However, forward scattering also determines several low-energy properties of systems with short-range interactions, and may lead to quite subtle effects especially in low dimensions. For example, forward scattering governs the leading low-energy long-wavelength response of a Fermi liquid as well as the breakdown of Fermi liquid theory in one-dimensional systems.

In this section we will derive several important relations for low-energy correlation functions in forward scattering dominated systems. These relations follow from the conservation of the velocities of particles in scattering processes with small momentum transfers, combined with the global conservation of charge and spin. They can also be viewed as *asymptotic Ward identities* associated with the separate conservation of charge and spin near each given point of the Fermi surface in a scattering process with $\mathbf{q} \sim \mathbf{0}$ [35].

An alternative way of treating forward scattering non-perturbatively, by *bosonization*, has been pioneered by Haldane [68]. This method, which leads to the same results as the Ward identity approach, will be reviewed in 5.3.

In the following we consider only scattering processes with small momentum transfers $|\mathbf{q}| \ll k_F$. We assume that the low-energy physics can be described by an effective *forward scattering action* S^F that involves only interactions with small momentum transfers $|\mathbf{q}| \ll k_F$. In cases where the low-energy physics is indeed governed by (residual) forward scattering only, S^F can be constructed from the bare action S as follows. Elimination of high-energy states maps S to an effective action $\bar{S}^\Lambda$, where $\Lambda \ll k_F$ is a band cutoff defining a thin shell around the Fermi surface (see Sec. 2). In forward scattering dominated systems, renormalized interactions with large momentum transfers are negligible compared to forward scattering in the low-energy limit $\Lambda \ll k_F$. We emphasize that in many cases of interest (e.g. most Luttinger liquids) large-$\mathbf{q}$ interactions contribute significantly to the map $S \mapsto \bar{S}^\Lambda$, but become irrelevant in the calculation of correlation functions in terms $\bar{S}^\Lambda$. Interactions with large momentum transfers now being negligible, we can impose an explicit *momentum transfer cutoff* of order Λ and shift the bandwidth cutoff to some larger value $\Lambda_0 \gg \Lambda$, without changing the qualitative physics. This step is

technically convenient because the smallness of momentum transfers can be best exploited if scattering processes are not contrained by a band cutoff. A quantitatively correct way of pushing the band cutoff back to larger values is to choose an effective action S^F with small momentum transfers $|\mathbf{q}| < \Lambda \ll k_F$ (here Λ is generally a *smooth* cutoff) and a band cutoff $\Lambda_0 \gg \Lambda$ such that mode elimination from Λ_0 to Λ would map S^F back to $\bar{S}^\Lambda$.[16] Schematically, S, $\bar{S}^\Lambda$ and S^F are thus related by

$$S \xrightarrow{\text{RG}} \underbrace{\bar{S}^\Lambda}_{|\mathbf{q}|<\Lambda\ll k_F} \xleftarrow{\text{RG}} \underbrace{S^F}_{|\mathbf{q}|<\Lambda\ll\Lambda_0,k_F} \tag{5.1}$$

All these actions describe the same low-energy physics, but only S^F is a suitable starting point for analyzing low-energy scattering processes to all orders in the (renormalized) couplings. An instructive one-dimensional example for (5.1) is: S = repulsive non-half-filled Hubbard model, $\bar{S}^\Lambda$ = g-ology model with band cutoff and back-scattering scaled to zero, S^F = Luttinger model (with momentum transfer cutoff). Note that the band cutoff Λ_0 can be ignored completely if the single-particle dispersion relation of the effective model is bounded from below. If it is unbounded (as in the g-ology model), Λ_0 must be kept to make the model well defined.

Here we consider effective forward scattering actions of the form

$$S^F = \sum_\sigma \int_k \psi^*_{k\sigma}(ik_0 - \xi_{\mathbf{k}})\psi_{k\sigma} \; - \; \tfrac{1}{2}\sum_{\sigma\sigma'}\int_{k,k',q} g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(q)\, \psi^*_{k-q/2,\sigma}\psi_{k+q/2,\sigma}\psi^*_{k'+q/2,\sigma'}\psi_{k'-q/2,\sigma'} \tag{5.2}$$

where $\xi_{\mathbf{k}}$ is a (possibly renormalized) dispersion relation, with the chemical potential subtracted, i.e. $\xi_{\mathbf{k}}$ vanishes on the Fermi surface. The momentum transfers $\mathbf{q}$ are restricted by the condition $|\mathbf{q}| \ll k_F$. We have suppressed bars (indicating renormalization) and explicit cutoffs in S^F to keep the notation readable, but keep of course in mind that $\xi_{\mathbf{k}}$ and $g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(q)$ in (5.2) differ from the bare dispersion and coupling function. The effective coupling $g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(q)$ may depend on momentum and energy transfers (including retarded interactions), but we restrict our analysis to systems where no relevant dependences on the energies k_0 and k'_0 occur. Clearly, S^F can describe the low-energy behavior only if interactions with large momentum transfers do not generate spontaneous symmetry breaking or other dramatic modifications of the effective low-energy action. Most of the subsequent analysis applies also to Fermi systems coupled to a gauge field, to be treated in more detail in Sec. 9.

[16] In practice one rather analyses classes of effective actions (with general coupling functions) and relates the unknown parameters to suitable physical quantities, which are accessible via analytic or numerical solutions (for microscopic models) or by experiment (for real systems).

5.1. GLOBAL CHARGE AND SPIN CONSERVATION

The effective action S^F, (5.2), conserves charge and the z-component of spin; for $g^{\sigma\sigma}_{\mathbf{k}\mathbf{k}'}(q) = g^{\sigma,-\sigma}_{\mathbf{k}\mathbf{k}'}(q)$ it is fully spin-rotation invariant. S_F thus respects the corresponding conservation laws (assumed to be not broken) of the underlying microscopic system. The correlation functions derived from S^F must therefore satisfy the Ward identities associated with these conservation laws. In Sec. 3 we have already derived Ward identities for systems with pure density-density interactions, which apply to most familiar microscopic models. In the following it will be useful, however, to derive Ward identities for (renormalized) correlation functions within the effective low-energy theory S^F. To this end we must now deal with other than density-density interactions, since the effective coupling function $g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(q)$ depends generically on $\mathbf{k}$ and $\mathbf{k}'$. In general, such interactions would lead to complicated currents with quartic (in the Fermi operators) interaction-dependent terms. Fortunately, the current operators associated with the effective forward scattering action S^F involve only a *quadratic* interaction dependent term (known as *"back-flow"* in Fermi liquids).

Let us address *charge* conservation first. Since we derive the Ward identities in a Hamiltonian language (as in Sec. 3), we assume for a moment that $g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(q)$ depends only on $\mathbf{q}$, not on q_0. The resulting identities are clearly not restricted by this assumption, and may be derived more generally via path integrals.

To construct a conserved current, we must evaluate the commutator $[\rho(\mathbf{q}), H]$, where $\rho(\mathbf{q})$ is the density fluctuation operator (cf. Sec. 3.1). The non-interacting part of the Hamiltonian yields a contribution $[\rho(\mathbf{q}), H_0] = \mathbf{q}\cdot\mathbf{j}_0(\mathbf{q})$ with a current operator $\mathbf{j}_0(\mathbf{q}) = \sum_{\mathbf{k},\sigma} \mathbf{v}_{\mathbf{k}}\, a^\dagger_{\mathbf{k}-\mathbf{q}/2,\sigma}\, a_{\mathbf{k}+\mathbf{q}/2,\sigma}$ for small $\mathbf{q}$, while the interaction part reads

$$\begin{aligned}[\rho(\mathbf{q}), H_I] = \tfrac{1}{2V} \sum_{\mathbf{k}\mathbf{k}'} \sum_{\sigma\sigma'} \sum_{\mathbf{q}'} g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(\mathbf{q}') \\ \times \Big[(a^\dagger_{\mathbf{k}-\mathbf{q}-\mathbf{q}'/2,\sigma}\, a_{\mathbf{k}+\mathbf{q}'/2,\sigma} - a^\dagger_{\mathbf{k}-\mathbf{q}'/2,\sigma}\, a_{\mathbf{k}+\mathbf{q}+\mathbf{q}'/2,\sigma})\, a^\dagger_{\mathbf{k}'+\mathbf{q}'/2,\sigma'}\, a_{\mathbf{k}'-\mathbf{q}'/2,\sigma'} \\ + a^\dagger_{\mathbf{k}-\mathbf{q}'/2,\sigma}\, a_{\mathbf{k}+\mathbf{q}'/2,\sigma}\, (a^\dagger_{\mathbf{k}'-\mathbf{q}+\mathbf{q}'/2,\sigma'}\, a_{\mathbf{k}'-\mathbf{q}'/2,\sigma'} - a^\dagger_{\mathbf{k}'+\mathbf{q}'/2,\sigma'}\, a_{\mathbf{k}'+\mathbf{q}-\mathbf{q}'/2,\sigma'}) \Big] \end{aligned} \tag{5.3}$$

For pure density-density interactions, this interaction term vanishes. Arbitrary interactions lead obviously to a rather cumbersome expression, which is quartic in the Fermi operators. For our forward scattering model (5.2), however, the expression can be reduced to a much simpler quadratic form. Firstly, non-diagonal terms in the brackets $(\ldots)$ can be neglected because they give rise to non-diagonal (with four different momenta) quartic contributions to the current operator, which are irrelevant in the low-energy limit. With the remaining diagonal contributions one obtains

$$[\rho(\mathbf{q}), H_I] = \tfrac{1}{2V} \sum_{\mathbf{k}\sigma} \sum_{\mathbf{k}'\sigma'} g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(-\mathbf{q})\, (n_{\mathbf{k}-\mathbf{q}/2,\sigma} - n_{\mathbf{k}+\mathbf{q}/2,\sigma})\, a^\dagger_{\mathbf{k}'-\mathbf{q}/2,\sigma'}\, a_{\mathbf{k}'+\mathbf{q}/2,\sigma'}$$

$$+ \tfrac{1}{2V} \sum_{\mathbf{k}\sigma} \sum_{\mathbf{k}'\sigma'} g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(\mathbf{q})\, a^\dagger_{\mathbf{k}-\mathbf{q}/2,\sigma}\, a_{\mathbf{k}+\mathbf{q}/2,\sigma}\, (n_{\mathbf{k}'-\mathbf{q}/2,\sigma'} - n_{\mathbf{k}'+\mathbf{q}/2,\sigma'}) \tag{5.4}$$

To linear order in $\mathbf{q}$ this can be written as $[\rho(\mathbf{q}), H_I] = \mathbf{q}\cdot\mathbf{j}_I(\mathbf{q})$ with a current operator

$$\mathbf{j}_I(\mathbf{q}) = \sum_{\mathbf{k},\sigma} \mathbf{v}^I_{\mathbf{k}\sigma}(\mathbf{q})\, a^\dagger_{\mathbf{k}-\mathbf{q}/2,\sigma}\, a_{\mathbf{k}+\mathbf{q}/2,\sigma} \tag{5.5}$$

where

$$\mathbf{v}^I_{\mathbf{k}\sigma}(\mathbf{q}) = -V^{-1} \sum_{\mathbf{k}'\sigma'} g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(\mathbf{q})\, \frac{\partial \langle n_{\mathbf{k}'\sigma'}\rangle}{\partial \mathbf{k}'} \tag{5.6}$$

In the last equation, we have replaced the operator $n_{\mathbf{k}\sigma}$ by its expectation value. To justify this step, let us decompose the $\mathbf{k}'$-integral in a radial integral (over k'_r) and a Fermi surface integral (over $\mathbf{k}'_F$). Since S^F involves only small momentum transfers, $\partial n_{\mathbf{k}'\sigma'}/\partial\mathbf{k}'$ is non-zero only close to the Fermi surface (for any low-energy state of the interacting system). A possible dependence of the coupling function $g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(\mathbf{q})$ on k'_r can therefore be neglected (it is clear that such a dependence is irrelevant in the low-energy limit anyway). Furthermore, since $n_{\mathbf{k}'\sigma'} = 1$ deep inside the Fermi sea and $n_{\mathbf{k}'\sigma'} = 0$ far outside, one has $\int dk'_r\, g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(\mathbf{q})\, \partial n_{\mathbf{k}'\sigma'}/\partial\mathbf{k}' = -g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'_F}(\mathbf{q})\, \hat{\mathbf{n}}_{\mathbf{k}'_F}$, where $\hat{\mathbf{n}}_{\mathbf{k}'_F}$ is a normal (with respect to the Fermi surface) unit vector at $\mathbf{k}'_F$. The k'_r integral over the operator expression thus yields a c-number, and the same c-number is obtained when replacing $n_{\mathbf{k}'\sigma'}$ by its expectation value.

We have thus derived a continuity equation of the form (3.3) with an interaction-dependent current operator

$$\mathbf{j}(\mathbf{q}) = \mathbf{j}_0(\mathbf{q}) + \mathbf{j}_I(\mathbf{q}) \tag{5.7}$$

The continuity equation implies Ward identities for the current-current correlator $J^{\mu\nu}(q) = -\frac{1}{V}\,\langle j^\mu(q)\, j^\nu(-q)\rangle$ and the vertex part $\Gamma^\mu_\sigma(p;q) = \langle j^\mu(q)\, a_{p-q/2,\sigma}\, a^\dagger_{p+q/2,\sigma}\rangle_{tr}$ precisely as in Sec. 3, with only one modification: the interaction part $\mathbf{v}^I_{\mathbf{k}\sigma}(\mathbf{q})$ has to be added to the velocity $\mathbf{v}_{\mathbf{k}}$ entering the function $\mathbf{c}(\mathbf{q})$, (3.12).

We now derive a very useful Ward identity for the *irreducible* vertex part

$$\Lambda^\mu_\sigma(p;q) = \langle j^\mu_0(q)\, a_{p-q/2,\sigma}\, a^\dagger_{p+q/2,\sigma}\rangle^{irr}_{tr} \tag{5.8}$$

which is constructed with the *non-interacting* (or "quasi-particle") part of the current operator, in contrast to the function

$$\Lambda'^\mu_\sigma(p;q) = \langle j^\mu(q)\, a_{p-q/2,\sigma}\, a^\dagger_{p+q/2,\sigma}\rangle^{irr}_{tr} \tag{5.9}$$

defined with the full current (in Sec. 3 no such distinction was needed, since there $\mathbf{j}_I$ vanished). The vertex part Γ^μ and its irreducible counterpart Λ'^μ are obviously related

by the Dyson equation (see also Fig. 5.1)

$$\Gamma^{\mu}_{\sigma}(p;q) = \Lambda'^{\mu}_{\sigma}(p;q) + \tfrac{1}{V}\sum_{\mathbf{k}'\sigma'}\sum_{\mathbf{k}''\sigma''} J^{\mu,\mathbf{k}'\sigma'}(q)\, g^{\sigma'\sigma''}_{\mathbf{k}'\mathbf{k}''}(\mathbf{q})\, \Lambda^{\mathbf{k}''\sigma''}_{\sigma}(p;q) \tag{5.10}$$

where

$$J^{\mu,\mathbf{k}\sigma}(q) = -\int \tfrac{dk_0}{2\pi}\, \langle j^{\mu}(q)\, a^{\dagger}_{k+q/2,\sigma}\, a_{k-q/2,\sigma}\rangle = \int \tfrac{dk_0}{2\pi}\, \Gamma^{\mu}_{\sigma}(k;q)\, G_{\sigma}(k{+}q/2)\, G_{\sigma}(k{-}q/2) \tag{5.11}$$

and

$$\Lambda^{\mathbf{k}\sigma'}_{\sigma}(p;q) = \int \tfrac{dk_0}{2\pi}\, \langle a^{\dagger}_{k-q/2,\sigma'}\, a_{k+q/2,\sigma'}\, a_{p-q/2,\sigma}\, a^{\dagger}_{p+q/2,\sigma}\rangle^{irr}_{tr} \tag{5.12}$$

The Ward identity (3.15) for Γ^{μ} yields

$$(iq_0,\mathbf{q})_{\mu} J^{\mu,\mathbf{k}\sigma}(q) = \sum_{\sigma}(\langle n_{\mathbf{k}-\mathbf{q}/2,\sigma}\rangle - \langle n_{\mathbf{k}+\mathbf{q}/2,\sigma}\rangle) \to -\mathbf{q}\cdot\frac{\partial\langle n_{\mathbf{k}\sigma}\rangle}{\partial\mathbf{k}} \tag{5.13}$$

Hence, applying $(iq_0,\mathbf{q})_{\mu}$ to the Dyson equation (5.10) one obtains

$$(iq_0,\mathbf{q})_{\mu}\Gamma^{\mu}_{\sigma}(p;q) = (iq_0,\mathbf{q})_{\mu}\Lambda'^{\mu}_{\sigma}(p;q) + \mathbf{q}\cdot\sum_{\mathbf{k}'\sigma'}\mathbf{v}^{I}_{\mathbf{k}'\sigma'}(\mathbf{q})\Lambda^{\mathbf{k}'\sigma'}_{\sigma}(p;q) = (iq_0,\mathbf{q})_{\mu}\Lambda^{\mu}_{\sigma}(p;q) \tag{5.14}$$

i.e. the "back-flow" terms cancel. The Ward identity for Γ^{μ} thus implies an identity of the same form for Λ^{μ}:

$$(iq_0,\mathbf{q})_{\mu}\Lambda^{\mu}_{\sigma}(p;q) = G^{-1}_{\sigma}(p{+}q/2) - G^{-1}_{\sigma}(p{-}q/2) \tag{5.15}$$

This result generalizes (3.21) to more general coupling functions than just $g(\mathbf{q})$.

From *spin* conservation one can derive analogous relations. In the case of full spin-rotation invariance one obtains

$$(iq_0,\mathbf{q})_{\mu}\Lambda^{a\mu}_{\sigma\sigma'}(p;q) = \; s^{a}_{\sigma\sigma'}\,[G^{-1}_{\sigma'}(p{+}q/2) - G^{-1}_{\sigma}(p{-}q/2)] \tag{5.16}$$

for the irreducible spin vertex part, defined with the non-interacting spin current operator $\mathbf{j}^{a}_{0}(\mathbf{q}) = \sum_{\mathbf{k}}\sum_{\sigma,\sigma'}\; \mathbf{v}_{\mathbf{k}}\, s^{a}_{\sigma\sigma'}\, a^{\dagger}_{\mathbf{k}-\mathbf{q}/2,\sigma}\, a_{\mathbf{k}+\mathbf{q}/2,\sigma'}$ (cf. Sec. 3.2.). If only S^z is conserved, (5.16) holds only for $a = z$.

5.2. VELOCITY CONSERVATION

For small momentum transfers $\mathbf{q}$, the velocity of a particle varies little in a scattering process: $\Delta\mathbf{v}_{\mathbf{k}} = \mathbf{v}_{\mathbf{k}+\mathbf{q}} - \mathbf{v}_{\mathbf{k}} \sim (\mathbf{q}\cdot\nabla_{\mathbf{k}})\mathbf{v}_{\mathbf{k}}$. In particular, for a continuum system with $\epsilon_{\mathbf{k}} = \mathbf{k}^2/2m$ one has $\Delta v_{\mathbf{k}}/v_{\mathbf{k}} \sim |\mathbf{q}|/k_F$ near the Fermi surface. To leading order in the small parameter $|\mathbf{q}|/k_F$, the velocity is conserved in the scattering process. This asymptotic conservation law leads to systematic cancellations and other asymptotic Ward identities, which hold (at least) to leading order in $|\mathbf{q}|/k_F$ and to any order in the coupling constant.

5.2.1. Loop cancellation and response functions

An important consequence of velocity conservation is *loop-cancellation*: For small momentum transfers, *Feynman diagrams involving fermionic loops with more than two insertions cancel each other*, i.e. the sum over permutations of various orderings of the insertions attached to the loop is smaller than what one would expect on the basis of naive power-counting. This cancellation has been noticed long ago in the Luttinger model, where loops cancel completely for any $\mathbf{q}$ [61], while in higher dimensions loop-cancellation has been assumed and exploited at least implicitly in many works on singular forward scattering. An explicit proof in higher dimensions has been constructed recently by Kopietz et al. [62] and by one of us [99]. Here we will provide a slightly different argument, avoiding the decomposition of momentum space in small sectors.

Consider a fermionic loop with N insertions as shown in Fig. 5.2. The insertions may be external density or current vertices, or may be due to interactions that connect the loop to the rest of a bigger Feynman diagram. The value of the diagram can be written as an integral over the function

$$f(q_1, \ldots, q_N) = \int_k h(\mathbf{k}_1, \ldots, \mathbf{k}_N; q_1, \ldots, q_N)\, G_0(k_1) \ldots G_0(k_N) \tag{5.17}$$

where $k_1 := k$, and $k_2, \ldots, k_N$ are given in terms of k and the momentum transfers q_ν by momentum conservation. The fermion lines are associated with the propagators $G_0(k) = [ik_0 - \xi_{\mathbf{k}}]^{-1}$ derived from the quadratic part of the effective action S^F. The function $h(\mathbf{k}_1, \ldots, \mathbf{k}_N; q_1, \ldots, q_N)$ depends in general on the rest of the Feynman diagram; h is independent of $\mathbf{k}_\nu$ if all insertions on the loop are pure density-vertices (external or due to density-density interactions). Eventually the momenta q_ν associated with interaction lines are to be integrated, while those associated with external density or current vertices remain fixed. In addition to the diagram in Fig. 5.2, one has contributions from all possible permutations of the insertions attached to the loop.

For $|\mathbf{q}_\nu| \ll k_F$, all the momenta $\mathbf{k}_\nu$ on the fermion lines are almost equal, i.e. $\mathbf{k}_\nu \approx \mathbf{k}$ for all ν. Hence, all the velocities on the lines in the loop are approximately equal to $\mathbf{v}_{\mathbf{k}} = \nabla \xi_{\mathbf{k}}$ and the function $h(\mathbf{k}_1, \ldots, \mathbf{k}_N; q_1, \ldots, q_N)$ can be replaced by $h(\mathbf{k}, \ldots, \mathbf{k}; q_1, \ldots, q_N)$, to leading order in $|\mathbf{q}_\nu|/k_F$. Corrections are proportional to $\mathbf{q}_\nu$ multiplied by the gradients of $\mathbf{v}_{\mathbf{k}}$ and $h(\mathbf{k}_1, \ldots, \mathbf{k}_N; q_1, \ldots, q_N)$ in $\mathbf{k}_\nu$-space.

From now on, one can simply follow the proof of loop-cancellation in the one-dimensional Luttinger model, where the Fermi velocity is a constant and the function h is independent of $\mathbf{k}_\nu$ [92]. The basic idea is to consider a loop with $N-1$ insertions with momenta $q_1, \ldots, q_{N-1}$ arranged in an arbitrary way, and sum over all possible positions of an additional insertion with momentum q_N (summing thus only a subset of permutations

of all the insertions). Using the basic relation

$$G_0(k+q/2)\,G_0(k-q/2) = \frac{G_0(k-q/2) - G_0(k+q/2)}{iq_0 - \mathbf{v_k}\cdot\mathbf{q} + \mathcal{O}(|\mathbf{q}|^3)} \tag{5.18}$$

for the product of propagators associated with the two lines connected to the extra-insertion, one finds that there is a complete cancellation of terms in the sum over all positions, to leading order in $\mathbf{q}_\nu$. Note that (5.18) is exact without corrections of order $|\mathbf{q}|^3$ in the Luttinger model and also in (unrenormalized) continuum systems with quadratic dispersion $\epsilon_{\mathbf{k}} = \mathbf{k}^2/2m$. In Appendix A we show explicitly how these cancellations work for the case $N = 3$.

Note that the above argument does not always imply that the sum over all permutations vanishes in the limit $q_\nu \to 0$, because the corrections of order $|\mathbf{q}_\nu|$ in the expansion of the velocities and the function h are multiplied by factors which are generally singular for small q_ν. What we have shown, however, is that cancellations reduce the contribution of loops with respect to what one would get from naive power-counting estimates. The naive degree of divergence of the function $f(q_1, \ldots, q_N)$ in the limit $q_\nu \to 0$ is given by $D_f = D_h + N - 2$. Here D_h is the behavior of the function h, each of the N propagators increases D_f by one, while the k-integral reduces the degree of divergence by two (the codimension of the Fermi surface in energy-momentum space). However, by virtue of the above cancellation, at least for systems with short-range interactions (where $g_{\mathbf{k}\mathbf{k}'}(q)$ is finite for $q \to 0$) loops with $N > 2$ and small momentum transfers are irrelevant in any dimension, while according to naive power-counting they would yield finite contributions to the correlation functions. As in one-dimensional systems, the cancellation guarantees that their contribution is suppressed by a positive power of Λ. The situation is not so clear for systems where the coupling function or scattering amplitude is singular for small momentum transfers, especially if strong singularities persist even after screening by polarization effects has been taken into account; this latter problem occurs in Fermi systems coupled to a gauge field (see Sec. 9). In such cases the above argument demonstrates only a cancellation of leading singularities appearing in single Feynman diagrams. A recent explicit low-order calculation [100] confirms in fact this cancellation and suggests the irrelevance of loops with $N > 2$ even in gauge systems. A general proof of irrelevance, however, would require a better control of corrections due to velocity fluctuations.

More detailed results are available for the simple loop

$$l(q_1, \ldots, q_N) = \int_k G_0(k_1) \ldots G_0(k_N) \tag{5.19}$$

which is relevant for cases where the function h depends only on $q_1, \ldots, q_N$, not on $\mathbf{k}$. It

is easy to show that [101]

$$l(0,\ldots,0) \equiv \int_k G_0(k)\ldots G_0(k) = -\frac{1}{(N-1)!}\left.\frac{\partial^{N-2}N(\xi)}{\partial\xi^{N-2}}\right|_{\xi=0} \tag{5.20}$$

where $N(\xi)$ is the density of states per spin (of the non-interacting system). Note that the limit $q_\nu \to 0$ is not unique and $l(0,\ldots,0)$ as defined above reflects the so-called static limit where the frequency components $q_{\nu 0}$ go to zero first. Note also that in this particular limit each single loop with $N > 2$ is already much smaller than the power-counting estimate, and no further cancellations occur in the sum over permutations. In one dimension with a linearized $\epsilon_{\mathbf{k}}$ and in two dimensions with $\epsilon_{\mathbf{k}} = \mathbf{k}^2/2m$ the density of states is a constant and $l(0,\ldots,0)$ thus vanishes for $N > 2$. For two-dimensional systems with $\epsilon_{\mathbf{k}} = \mathbf{k}^2/2m$ the problem of evaluating $l(q_1,\ldots,q_N)$ has been recently reduced to elementary integrals for any N [102]. An explicit (possibly lengthy) evaluation of these integrals with an analysis of the detailed behavior for small q_ν has not yet been done.

As a consequence of loop-cancellation, the effective interaction (including polarization) and response functions for small $\mathbf{q}$ are given by the *random phase approximation* (RPA), even in cases (such as Luttinger liquids) where contributions beyond RPA are not simply suppressed by a small phase space and yield singular corrections to other quantities such as the single-particle Green function. The *effective interaction* D is obtained by summing the diagrams illustrated in Fig. 5.3, which is equivalent to the integral equation

$$D^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(q) = g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(q) + \sum_{\sigma''}\int_{k''} g^{\sigma\sigma''}_{\mathbf{k}\mathbf{k}''}(q) G_0(k''-q/2)G_0(k''+q/2) D^{\sigma''\sigma'}_{\mathbf{k}''\mathbf{k}'}(q) \tag{5.21}$$

In general, the bubbles in Fig. 5.3 must be dressed by self-energy and vertex corrections. However, these corrections involve fermionic loops with more than two insertions and must therefore cancel each other for small $\mathbf{q}$ and Λ, at least to leading order in these small parameters.

The charge current-current response function $J^{\mu\nu}(q)$ for small q can be constructed from the effective interaction in the way illustrated in Fig. 5.4, i.e.

$$\begin{aligned} J^{\mu\nu}(q) = J_0^{\mu\nu}(q) \; + \; \sum_{\sigma\sigma'}\int_{k,k'} & \lambda^\mu(\mathbf{k})\, G_0(k-q/2)G_0(k+q/2) \\ & \times D^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(q)\, G_0(k'-q/2)G_0(k'+q/2)\, \lambda^\nu(\mathbf{k}') \end{aligned} \tag{5.22}$$

where $\lambda^\mu(\mathbf{k}) = (1, \mathbf{v}_{\mathbf{k}})$ and

$$J_0^{\mu\nu}(q) = 2\int_k \lambda^\mu(\mathbf{k})\, G_0(k-q/2)G_0(k+q/2)\, \lambda^\nu(\mathbf{k}) \tag{5.23}$$

The factor 2 is due to the spin sum. Again, vertex and self-energy corrections cancel each other in the loops.

Similarly, the spin current-current response function $J^{a\mu,b\nu}(q)$ can be written as

$$J^{z\mu,z\nu}(q) = J_0^{z\mu,z\nu}(q) \; + \; \sum_{\sigma\sigma'} \int_{k,k'} \lambda^{z\mu}_{\sigma\sigma}(\mathbf{k}) \, G_0(k-q/2) G_0(k+q/2) \\ \times D^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(q) \, G_0(k'-q/2) G_0(k'+q/2) \, \lambda^{z\nu}_{\sigma'\sigma'}(\mathbf{k}') \tag{5.24}$$

for small q, where $\lambda^{z\mu}_{\sigma\sigma'}(\mathbf{k}) = s^z_{\sigma\sigma'}(1, \mathbf{v_k})^\mu$ and

$$J_0^{z\mu,b\nu}(q) = \sum_\sigma \int_k \lambda^{z\mu}_{\sigma\sigma}(\mathbf{k}) \, G_0(k-q/2) G_0(k+q/2) \, \lambda^{z\nu}_{\sigma\sigma}(\mathbf{k}) \tag{5.25}$$

5.2.2. Density and current vertex

Asymptotic velocity conservation implies a simple asymptotic Ward identity for the irreducible charge vertex $\Lambda^\mu_\sigma(p;q)$, defined in eq. (5.8).

The perturbation expansion for Λ^μ can be represented by Feynman diagrams as in Fig. 5.5, where the zeroth-order vertex is $\lambda^\mu(\mathbf{k}) = (1, \mathbf{v_k})^\mu$. We recall that one-interaction reducibe diagrams do not contribute. If the external momentum transfer $\mathbf{q}$ as well as the momentum transfers $\mathbf{q}_\nu$ due to interactions are all small, one has the following simplifications (to leading order in $\mathbf{q}$ and $\mathbf{q}_\nu$). Diagrams containing fermion loops with more than two insertions cancel each other as a consequence of loop cancellation. An example for two such cancelling diagrams is given in Fig. 5.6. In the remaining diagrams the vertex $\lambda^\mu(\mathbf{k})$ must lie on the fermion line connecting the two fermionic external points of $\Lambda^\mu_\sigma(p;q)$ (carrying momenta $p \pm q/2$). The momentum $\mathbf{k}$ passing through the vertex $\lambda^\mu(\mathbf{k})$ differs from $\mathbf{p}$ only by certain linear combinations of the small momentum transfers $\mathbf{q}_\nu$ caused by the interactions. Thus, to leading order in $\mathbf{q}_\nu$ we can replace $\lambda^\mu(\mathbf{k})$ by $\lambda^\mu(\mathbf{p})$ in each diagram contributing to $\Lambda^\mu(p;q)$. This implies that the current vertex $\mathbf{\Lambda}$ is related to the density vertex Λ^0 by the simple asymptotic identity [35, 63]

$$\mathbf{\Lambda}_\sigma(p;q) = \mathbf{v_p} \Lambda^0_\sigma(p;q) \tag{5.26}$$

Combining this with the Ward identity (5.15) associated with charge conservation, one obtains [35, 63]

$$\Lambda^0_\sigma(p;q) = \frac{G^{-1}_\sigma(p+q/2) - G^{-1}_\sigma(p-q/2)}{iq_0 - \mathbf{v_p} \cdot \mathbf{q}} \tag{5.27}$$

These asymptotic Ward identities express the charge vertex Λ^μ uniquely in terms of the propagator G. They are exact in the one-dimensional Luttinger model [61], while in general they hold only asymptotically for small momentum transfers, with the additional proviso that singular interactions do not overpower the loop-cancellation.

For the z-component of the irreducible spin vertex, one obtains essentially the same results, i.e.

$$\mathbf{\Lambda}^z_{\sigma\sigma'}(p;q) = \mathbf{v_p} \Lambda^{z0}_{\sigma\sigma'}(p;q) \tag{5.28}$$

and

$$\Lambda^{z0}_{\sigma\sigma'}(p;q) = s^z_{\sigma\sigma'}\,\frac{G^{-1}_{\sigma'}(p+q/2) - G^{-1}_{\sigma}(p-q/2)}{iq_0 - \mathbf{v_p}\cdot\mathbf{q}} \tag{5.29}$$

Corrections to the asymptotic identities (5.27) and (5.29) can be parametrized by adding a (generally unknown) complex function $Y(p;q)$ of order $\mathcal{O}(\mathbf{q}^2, \mathbf{q}_\nu^2)$ to the denominator $iq_0 - \mathbf{v_p}\cdot\mathbf{q}$ on the right hand side [35, 63].

It is easy to see that the above Ward identities for Λ imply that polarization bubbles are not dressed by interactions (as follows already from loop cancellation). For example, consider the charge density polarization, which can always be written as

$$\Pi(q) = \sum_\sigma \int_p \Lambda^0_\sigma(p;q)\, G_\sigma(p+q/2)\, G_\sigma(p-q/2) \tag{5.30}$$

Inserting (5.27) and performing the p_0-integral, one obtains

$$\Pi(q) = \sum_\sigma \int_{\mathbf{p}} \frac{\langle n_{\mathbf{p}-\mathbf{q}/2,\sigma}\rangle - \langle n_{\mathbf{p}+\mathbf{q}/2,\sigma}\rangle}{iq_0 - \mathbf{v_p}\cdot\mathbf{q}} = \Pi_0(q) \tag{5.31}$$

where $\Pi_0(q)$ is the bubble for the non-interacting system. The last equation holds since $\langle n_{\mathbf{p}-\mathbf{q}/2,\sigma}\rangle - \langle n_{\mathbf{p}+\mathbf{q}/2,\sigma}\rangle = -\mathbf{q}\cdot\partial\langle n_{\mathbf{p}\sigma}\rangle/\partial\mathbf{p}$ for small $\mathbf{q}$ and $\int dp_r\, \partial\langle n_{\mathbf{p}\sigma}\rangle/\partial\mathbf{p} = -\hat{\mathbf{n}}_{\mathbf{p}_F}$ for any momentum distribution in the forward scattering model.

5.2.3. Single-particle propagator

The self-energy correction due to small-$\mathbf{q}$ scattering obeys the Dyson equation represented diagrammatically in Fig. 5.7. Algebraically, this equation reads

$$\Sigma_\sigma(p) = -\sum_{\sigma'} \int_{p'} D^{\sigma\sigma'}_{\mathbf{pp}}(p-p')\, G_\sigma(p')\, \Lambda^0_{\sigma'\sigma}[(p+p')/2; p'-p] \tag{5.32}$$

where

$$\Lambda^0_{\sigma'\sigma}(p;q) = \langle \rho_{\sigma'}(q) a_{p-q/2,\sigma} a^\dagger_{p+q/2,\sigma}\rangle^{irr}_{tr} \tag{5.33}$$

is the irreducible density vertex for fermions with spin orientation σ, with $\rho_\sigma(\mathbf{q}) = \sum_{\mathbf{k}} a^\dagger_{\mathbf{k}-\mathbf{q}/2,\sigma} a_{\mathbf{k}+\mathbf{q}/2,\sigma}$. The effective interaction D is given in terms of the coupling function $g^{\sigma\sigma'}_{\mathbf{kk}'}(q)$ by Eq. (5.21). Note that (5.32) holds only if the contributing momentum transfers $\mathbf{q} = \mathbf{p} - \mathbf{p}'$ are small compared to the scale on which $g^{\sigma\sigma'}_{\mathbf{kk}'}(q)$ varies as a function of $\mathbf{k}$ and $\mathbf{k}'$.

The asymptotic Ward identies (5.27) and (5.29) for the charge and spin vertex imply the identity

$$\Lambda^0_{\sigma'\sigma}(p;q) = \delta_{\sigma'\sigma}\,\frac{G^{-1}_{\sigma}(p+q/2) - G^{-1}_{\sigma}(p-q/2)}{iq_0 - \mathbf{v_p}\cdot\mathbf{q}} \tag{5.34}$$

Note that $\Lambda^0_{\sigma,-\sigma}$ must vanish as a direct consequence of loop cancellation. Now $\Lambda^0_{\sigma'\sigma}(p;q)$ can be eliminated from (5.32) in favor of G_σ. Combining the result of this substitution with the exact Dyson equation $G = G_0 + G_0\Sigma G$, one obtains a linear integral equation that determines G in terms of D [35]:

$$(ip_0 - \xi_{\mathbf{p}})G_\sigma(p) \;=\; 1 \;-\; \int_{p'} \frac{D^{\sigma\sigma}_{\mathbf{pp}}(p-p')}{ip_0 - ip_0' - \mathbf{v}_{(\mathbf{p}+\mathbf{p}')/2}\cdot(\mathbf{p}-\mathbf{p}')}\, G_\sigma(p') \;+\; X_\sigma(p)G_\sigma(p) \quad (5.35)$$

where

$$X_\sigma(p) = \int_{p'} \frac{D^{\sigma\sigma}_{\mathbf{pp}}(p-p')}{ip_0 - ip_0' - \mathbf{v}_{(\mathbf{p}+\mathbf{p}')/2}\cdot(\mathbf{p}-\mathbf{p}')} \quad (5.36)$$

Let us now specialize to rotationally invariant systems, to arrive at a more explicit solution. In this case the propagator depends only on two variables, the energy p_0 and the radial momentum component $p_r = |\mathbf{p}| - k_F$, i.e. we can write $G(p) = G(p_0, p_r)$. Furthermore $|\mathbf{v}_\mathbf{k}| = v_F$ is now constant, and for small $\mathbf{q} = \mathbf{p} - \mathbf{p}'$ we can use the relation

$$\mathbf{v}_{(\mathbf{p}+\mathbf{p}')/2}\cdot(\mathbf{p}-\mathbf{p}') \;=\; v_F(p_r - p_r') + \mathcal{O}(\mathbf{q}^2) \quad (5.37)$$

For each momentum $\mathbf{p}$ near the Fermi surface, the momentum transfer $\mathbf{q}$ can be decomposed in a radial (or normal) component $\mathbf{q}_r$ and a tangential component $\mathbf{q}_t$, via

$$\mathbf{q}_r = q_r\hat{\mathbf{p}}\,, \quad q_r = \mathbf{q}\cdot\hat{\mathbf{p}}\,, \quad \mathbf{q}_t = \mathbf{q} - \mathbf{q}_r \quad (5.38)$$

where $\hat{\mathbf{p}}$ is a unit vector parallel to $\mathbf{p}$ (i.e. normal to the Fermi surface near $\mathbf{p}$). For small $\mathbf{q}_t$, we have $p_r - p_r' = q_r + \mathcal{O}(\mathbf{q}_t^2)$, and $|\mathbf{q}_t|/k_F$ is the angle between $\mathbf{p}$ and $\mathbf{p}'$. For isotropic systems and small $\mathbf{q}$, the effective interaction $D^{\sigma\sigma}_{\mathbf{pp}}(q_0, \mathbf{q})$ can be parametrized by q_0, q_r and $q_t = |\mathbf{q}_t|$, i.e.

$$D^{\sigma\sigma}_{\mathbf{pp}}(q_0, \mathbf{q}) = D(q_0, q_r, q_t) \quad \text{for} \quad \mathbf{q} \sim \mathbf{0} \quad (5.39)$$

Note that $q_t \geq 0$ by definition, while q_r can also be negative. The crucial point is that small tangential momentum transfers enter the integrand in Eqs. (5.35) and (5.36) only via the effective interaction D. We therefore define a q_t-averaged effective interaction

$$\bar{D}^{\Lambda_t}(q_0, q_r) = \frac{S_{d-1}}{(2\pi)^{d-1}} \int_0^{\Lambda_t} dq_t\, q_t^{d-2}\, D(q_0, q_r, q_t) \quad (5.40)$$

where $S_{d-1} = 2\pi^{d/2}/\Gamma(d/2)$ is the surface of the d-dimensional unit-sphere, and Λ_t is a cutoff for tangential momentum transfers. The factor q_t^{d-2} is due to the phase space in the $(d-1)$-dimensional tangential space spanned by $\mathbf{q}_t$. We will see that leading low-energy terms are often independent of the cutoff Λ_t, confirming the dominance of forward scattering. In the following we will usually suppress the Λ_t-dependence of $\bar{D}$ in our

notation. Note that in one dimension, where no tangential degrees of freedom exist, eq. (5.40) reduces to $\bar{D} = D$.

In terms of $\bar{D}$, the function $X_\sigma(p)$ in (5.36) can be written as

$$X(p) = \int_{q_0,q_r} \frac{\bar{D}(q_0,q_r)}{iq_0 - v_F q_r} \tag{5.41}$$

where $q_0 = p_0 - p_0'$ and $q_r = p_r - p_r'$. This is obviously a constant (p-independent), which can been absorbed by shifting the chemical potential in $\xi_{\mathbf{k}} = \epsilon_{\mathbf{k}} - \mu$ (keeping thus the density and the Fermi surface fixed). The integral equation for the propagator now simplifies to [35]

$$(ip_0 - v_F p_r)\, G(p_0,p_r) \;=\; 1 \;-\; \int_{p_0',p_r'} \frac{\bar{D}(p_0-p_0',p_r-p_r')}{ip_0 - ip_0' - v_F(p_r-p_r')}\, G(p_0',p_r') \tag{5.42}$$

where we have omitted the (now unimportant) spin variables. This equation has precisely the same form as the equation obtained by Dzyaloshinkii and Larkin [61] for the propagator of the one-dimensional Luttinger model, with D substituted by $\bar{D}$. Thus, to leading order in small momentum transfers, the single-particle propagator is given by a *one-dimensional* equation of motion corresponding to a ficticious Luttinger model with an effective interaction given by $\bar{D}(q_0,q_r)$.

The solution of the integral equation (5.42) proceeds as in one dimension [61]. While Eq. (5.42) holds also at small finite temperatures (with p_0' to be summed over the Matsubara frequencies), we present the solution only at zero temperature. Continuing (5.42) to real frequencies yields

$$(\xi - v_F p_r)\, G(\xi,p_r) \;=\; 1 \;+\; \int_{\xi',p_r'} \frac{i\bar{D}(\xi-\xi',p_r-p_r')}{\xi - \xi' - v_F(p_r-p_r')}\, G(\xi',p_r') \tag{5.43}$$

We transform this equation to real space-time by defining

$$G(t,r) = \int \frac{d\xi}{2\pi} \int \frac{dp_r}{2\pi}\, G(\xi,p_r)\, e^{ip_r r - i\xi t} \tag{5.44}$$

Note that $G(t,r)$ is *not* the $(d+1)$-dimensional Fourier transform of $G(\xi,\mathbf{p})$. The integral equation (5.43) transfroms to a partial differential equation for $G(t,r)$, namely

$$(\partial_t + v_F\partial_r)G(t,r) = \delta(t)\delta(r) + K(t,r)G(t,r) \tag{5.45}$$

where $K(t,r)$ is the Fouriertransform of $\bar{D}(\omega,q_r)/(\omega - v_F q_r)$. This differential equation is solved by the ansatz

$$G(t,r) = e^{L(t,r) - L_0(r-v_F t)} G_0(t,r) \tag{5.46}$$

where $L(t,r)$ is a solution of the inhomogeneous linear differential equation

$$(\partial_t + v_F\partial_r)L(t,r) = K(t,r) \tag{5.47}$$

i.e. the Fouriertransform of $L(\omega,q_r) = i\bar{D}(\omega,q_r)/(\omega - v_F q_r)^2$. More explicitly, one obtains

$$L(t,r) = \int \frac{d\omega}{2\pi}\int \frac{dq_r}{2\pi}\,\frac{i\bar{D}(\omega,q_r)}{[\omega - v_F q_r + i0^+ s(\omega)]^2}\,e^{iq_r r - i\omega t} \tag{5.48}$$

where $s(.)$ is the sign-function. The function $L_0(r-v_F t)$ is a solution of the corresponding homogeneous equation, which is chosen such that boundary conditions in the complex t-plane are satisfied: Analyticity of $G(t,r)$ in the second and forth quadrant of the t-plane and

$$G(0^+,r) - G(0^-,r) = -i\delta(r) \tag{5.49}$$

The non-interacting propagator in (t,r)-representation reads

$$G_0(t,r) = \frac{1}{2\pi}\,\frac{1}{r - v_F t + i0^+ s(t)} \tag{5.50}$$

and satisfies of course the boundary conditions. Analyticity of $G(t,r)$ is automatically guaranteed (even for $L_0(r-v_F t) = 0$), since $\bar{D}$ is a time-ordered function, while the condition (5.49) can be satisfied by choosing

$$L_0(r-v_F t) = L(0,0) =: L_0 \tag{5.51}$$

i.e. a constant. The extension of the above solution to anisotropic systems is straightforward: one simply has to replace v_F by $v_{\mathbf{k}_F}$ and add a label $\mathbf{k}_F$ to $\bar{D}$, L and G.

The *momentum distribution* $n_{\mathbf{p}}$ is obtained by Fourier transforming $G(0^-,r)$, i.e.

$$n_{\mathbf{p}} = -iG(0^-,\mathbf{p}) = -i\int_{-\infty}^{\infty} dr\, G(0^-,r)\,e^{-ip_r r} \tag{5.52}$$

The *density of states* (per spin) can be calculated by Fourier transforming $G(t,0)$, i.e.

$$N(\xi) = \pi^{-1}\,\left|\,\mathrm{Im}\int_{-\infty}^{\infty} dt\, G(t,0)\,e^{i\xi t}\,\right| \tag{5.53}$$

In Appendix B we derive an expression for $L(t,r)$ in terms of the spectral function $\bar{\Delta}$ associated with $\bar{D}$. With that representation one can check general analytic properties such as the reality of $n_{\mathbf{p}}$.

In $d > 1$ the above result for the propagator G is plagued by unphysical exponential singularities in the special limit $r,t \to \infty$ with $r/t \to v_F$, corresponding to the quasiparticle regime $p_r, \xi \to 0$ with $\xi/p_r \to v_F$. These singularities are associated with the

double pole in $\omega = v_F q_r$ in the expression (5.48) for the exponent $L(t,r)$. In the one-dimensional Luttinger model this double pole is reduced to a simple pole by the effective interaction in the numerator, which vanishes linearly for $\omega \to v_F q_r$ in that case. In general, however, $\bar{D}(\omega, q_r)$ is finite for $\omega = v_F q_r$, but the double pole should then be cut off by neglected terms of order $\mathbf{q}^2$ in the denominator, especially in the Ward identities (5.27) and (5.29). Although these quadratic corrections are usually subleading, they become important when the leading term $\omega - v_F q_r$ in the denominator vanishes. Indeed, for $r, t \to \infty$ with $r/v_F t = 1$, the function $L(t,r)$ is dominated by Fourier components $L(\omega, q_r)$ with $\omega \sim v_F q_r$, which is precisely the situation where corrections should be important, and would cut off the pole in the Ward identities. In all other cases one integrates over many values for the ratio $\omega / v_F q_r$, and quadratic corrections are thus negligible. In particular, the results for the momentum distribution and the density of states, obtained from $G(0,r)$ and $G(t,0)$, respectively, are stable.

Asymptotic Ward identities and a solution for the fermion propagator in Fermi systems with gauge-fields will be discussed in Sec. 9.

5.3. BOSONIZATION

An alternative way of treating systems whose low-energy physics is dominated by forward scattering is *bosonization*, i.e. the representation of the Hamiltonian and of fermionic creation and annihilation operators in terms of (bosonic) density fluctuation operators. This technique has been originally invented for the analysis of one-dimensional Fermi systems [22, 65, 66], where it proved to be an extremely valuable non-perturbative tool [20, 25, 23]. An early generalization of bosonization to higher dimensions by Luther [67] could deal only with very special scattering processes, and therefore remained essentially unnoticed. Recently, however, Haldane [68] invented a different bosonization approach in dimensions $d \geq 1$, which turned out to be more successful. Haldane's idea of d-dimensional bosonization has been elaborated in detail by Houghton and Marston [69], and subsequently by many others [70, 71, 72]. A formulation of Haldane's bosonization in terms of functional integrals instead of operators and Hamiltonians has been proposed by Kopietz and Schönhammer [73], and further elaborated by Kopietz et al. [62]. Mathematically oriented readers are also referred to a paper by Fröhlich et al. [103] on a bosonization approach in terms of gauge forms.

We will now briefly review the main ideas of the bosonization approach to d-dimensional Fermi systems. As already pointed out by several authors [38, 70, 62], the results obtained by bosonization are equivalent to those obtained from asymptotic Ward identities. Since spin plays no important role in the following, we consider spinless fermions to make the notation as simple as possible.

Common to all versions of d-dimensional bosonization is a decomposition of momentum space in disjoint sectors K_α, $\alpha = 1, \ldots, M$, which yields a corresponding partition of the Fermi surface $\partial\mathcal{F}$ in "patches" $\partial\mathcal{F}_\alpha = K_\alpha \cap \partial\mathcal{F}$. Let Λ_α be a length in momentum space such that Λ_α^{d-1} is the ($(d-1)$-dimensional) area of the patch $\partial\mathcal{F}_\alpha$. The size of the patches must be choosen small enough to make sure that the Fermi surface is almost flat and the Fermi velocity almost constant within each patch; for most systems one thus has to require $\Lambda_\alpha \ll k_F$. The condition of constant velocities is equivalent to the velocity conservation underlying the Ward identity approach. The central objects in the bosonization method are the *density fluctuation operators on sectors*

$$\rho_\alpha(\mathbf{q}) = \sum_{\mathbf{k}} \chi_\alpha(\mathbf{k}-\mathbf{q}/2)\, \chi_\alpha(\mathbf{k}+\mathbf{q}/2)\, a^\dagger_{\mathbf{k}-\mathbf{q}/2}\, a_{\mathbf{k}+\mathbf{q}/2} \tag{5.54}$$

where $\chi_\alpha(\mathbf{k})$ is the characteristic function of the sector K_α (i.e. one if $\mathbf{k} \in K_\alpha$ and zero else). For $|\mathbf{q}| \ll \Lambda_\alpha$, the operators $\rho_\alpha(\mathbf{q})$ obey the commutation relation [68, 69]

$$[\rho_\alpha(\mathbf{q}), \rho_{\alpha'}(\mathbf{q}')] \;\sim\; \delta_{\alpha\alpha'}\delta_{\mathbf{q},-\mathbf{q}'}\, \frac{V}{(2\pi)^d}\, \Lambda_\alpha^{d-1}\, \mathbf{n}_\alpha\cdot\mathbf{q} \tag{5.55}$$

on a (restricted) Hilbert space of states that differ from the Fermi sea only in the vicinity of the Fermi surface within a distance $\lambda_\alpha \ll \Lambda_\alpha$. Here $\mathbf{n}_\alpha$ is a unit vector normal to the patch $\partial\mathcal{F}_\alpha$ and V is the volume of the system. Possibly existing band cutoffs must be much larger than $|\mathbf{q}|$. In one-dimensional systems one needs only two sectors corresponding to left-moving and right-moving particles, respectively; in that case the two "patches" are simply the two Fermi points, and the "area" Λ_α^{d-1} in (5.55) has to be substituted by a factor one. Notice that in $d = 1$ eq. (5.55) is an exact operator identity. In $d > 1$, instead, it holds only asymptotically for states with a support $\lambda_\alpha \ll \Lambda_\alpha \ll k_F$.

The density fluctuation operators $\rho_\alpha(\mathbf{q})$ can be related to canonically normalized boson annihililation and creation operators $b_\alpha(\mathbf{q})$ and $b^\dagger_\alpha(\mathbf{q})$ with $\mathbf{n}_\alpha\cdot\mathbf{q} > 0$ via

$$\rho_\alpha(\mathbf{q}) = \sqrt{\Omega_\alpha|\mathbf{n}_\alpha\cdot\mathbf{q}|}\, \Big[\Theta(\mathbf{n}_\alpha\cdot\mathbf{q})\, b_\alpha(\mathbf{q}) + \Theta(-\mathbf{n}_\alpha\cdot\mathbf{q})\, b^\dagger_\alpha(-\mathbf{q})\Big] \tag{5.56}$$

where $\Omega_\alpha \equiv V\tilde{\Omega}_\alpha = (V/(2\pi)^d)\, \Lambda_\alpha^{d-1}$. The bosonic commutation relation

$$[b_\alpha(\mathbf{q}), b^\dagger_{\alpha'}(\mathbf{q}')] = \delta_{\alpha\alpha'}\delta_{\mathbf{q}\mathbf{q}'} \tag{5.57}$$

is equivalent to (5.55).

Correlation functions involving fermionic creation and annihilation operators (especially the single-particle propagator) can be expressed as expectation values of the operators [69]

$$\psi_\alpha(t, \mathbf{r}) = \sqrt{\lambda\tilde{\Omega}_\alpha}\, e^{i\mathbf{k}_{F\alpha}\cdot\mathbf{r}}\, e^{i\tilde{\Omega}_\alpha^{-1}\sqrt{4\pi}\phi_\alpha(t,\mathbf{r})}\, O_\alpha \tag{5.58}$$

and the corresponding conjugate $\psi^\dagger_\alpha(t,\mathbf{r})$, where $\phi_\alpha(\mathbf{r})$ is the Fourier transform of

$$\phi_\alpha(\mathbf{q}) = \rho_\alpha(\mathbf{q})/[i\sqrt{4\pi}\,\mathbf{n}_\alpha\cdot\mathbf{q}] \tag{5.59}$$

and λ is an ultraviolet cutoff; $\mathbf{k}_{F\alpha}$ is a suitably chosen Fermi momentum belonging to the patch $\partial\mathcal{F}_\alpha$. The ordering operator O_α is chosen such that $\psi_\alpha(\mathbf{r})$ and $\psi_{\alpha'}(\mathbf{r})$ obey fermionic anticommutation relations also for $\alpha \neq \alpha'$.

The main point of the bosonization idea is the observation that the Hamiltonian (or action) of systems with scattering processes restricted to small momentum transfers can be expressed as a *quadratic form* in $\rho_\alpha(\mathbf{q})$, to leading order in $\mathbf{q}$. Combining this with the bosonization (5.58) of fermionic operators, any correlation function can be calculated explicitly.

Consider the spinless forward scattering Hamiltonian

$$H^F = H_0 + H_I = \sum_{\mathbf{k}} \xi_{\mathbf{k}} a^\dagger_{\mathbf{k}} a_{\mathbf{k}} + \tfrac{1}{2V}\sum_{\mathbf{k},\mathbf{k}',\mathbf{q}} g_{\mathbf{k}\mathbf{k}'}(\mathbf{q})\, a^\dagger_{\mathbf{k}-\mathbf{q}/2}\, a_{\mathbf{k}+\mathbf{q}/2}\, a^\dagger_{\mathbf{k}'+\mathbf{q}/2}\, a_{\mathbf{k}'-\mathbf{q}/2} \tag{5.60}$$

which is the Hamiltonian corresponding to a spinless version of the forward scattering action S^F in Eq. (5.2), with a non-retarded coupling function $g_{\mathbf{k}\mathbf{k}'}(\mathbf{q})$. Momentum transfers are assumed to satisfy the condition $|\mathbf{q}| \ll k_F$.

For $\Lambda_\alpha \ll k_F$ and excitations restricted to the vicinity of the Fermi surface, the kinetic part of the Hamiltonian can be written as [68, 69]

$$H_0 = \tfrac{1}{2}\sum_\alpha \sum_{\mathbf{q}} \frac{\mathbf{v}_\alpha\cdot\mathbf{q}}{\Omega_\alpha|\mathbf{n}_\alpha\cdot\mathbf{q}|}\ :\rho_\alpha(\mathbf{q})\rho_\alpha(-\mathbf{q}): = \sum_\alpha\sum_{\mathbf{q}} \Theta(\mathbf{n}_\alpha\cdot\mathbf{q})\,\mathbf{v}_\alpha\cdot\mathbf{q}\, b^\dagger_\alpha(\mathbf{q}) b_\alpha(\mathbf{q}) \tag{5.61}$$

where $\mathbf{v}_\alpha$ is the velocity $\mathbf{v}_{\mathbf{k}} = \nabla\xi_{\mathbf{k}}$ for states near the patch $\partial\mathcal{F}_\alpha$, and the double dots indicate normal ordering with respect to the Fermi sea. The condition $\Lambda_\alpha \ll k_F$ makes sure that the velocity $\mathbf{v}_{\mathbf{k}}$ can be replaced by a constant $\mathbf{v}_\alpha$ within each sector K_α.

For interactions with momentum transfers restricted by $|\mathbf{q}| \ll \Lambda_\alpha$, the interaction part of the Hamiltonian can be written as

$$H_I = \tfrac{1}{2V}\sum_{\alpha,\alpha'}\sum_{\mathbf{q}} g_{\alpha\alpha'}(\mathbf{q})\, :\rho_\alpha(\mathbf{q})\,\rho_{\alpha'}(-\mathbf{q}): \tag{5.62}$$

where $g_{\alpha\alpha'}(\mathbf{q})$ is obtained from $g_{\mathbf{k}\mathbf{k}'}(\mathbf{q})$ by averaging over $\mathbf{k} \in \partial\mathcal{F}_\alpha$ and $\mathbf{k}' \in \partial\mathcal{F}_{\alpha'}$. Via (5.56) this can be also be expressed as a quadratic form of the canonical Bose creation and annihilation operators $b^\dagger_\alpha(\mathbf{q})$ and $b_\alpha(\mathbf{q})$. The condition $|\mathbf{q}| \ll \Lambda_\alpha$ is necessary to make scattering of particles from one sector to another (not taken into account in (5.42)) rare.

For interactions satisfying $|\mathbf{q}| \ll k_F$ one can choose the size of the sectors K_α such $|\mathbf{q}| \ll \Lambda_\alpha \ll k_F$. Using (5.61) and (5.62), the whole Hamiltonian $H_0 + H_I$ can then be

written as a quadratic form in $\rho_\alpha(\mathbf{q})$, i.e.

$$H^F = \sum_{\alpha,\alpha'} \sum_{\mathbf{q}} \left[\tfrac{\mathbf{v}_\alpha \cdot \mathbf{q}}{2\Omega_\alpha |\mathbf{n}_\alpha \cdot \mathbf{q}|} \delta_{\alpha\alpha'} + \tfrac{1}{2V} g_{\alpha\alpha'}(\mathbf{q}) \right] :\rho_\alpha(\mathbf{q}) \rho_{\alpha'}(-\mathbf{q}): \tag{5.63}$$

or, equivalently, in $b^\dagger_\alpha(\mathbf{q})$ and $b_\alpha(\mathbf{q})$. Clearly these results hold only to leading order in $\mathbf{q}$, since corrections due to scattering into different sectors have been neglected.

Due to the quadratic structure of the bosonized representation of the Hamiltonian H^F, the dynamics of the operators $\rho_\alpha(\mathbf{q})$ can be calculated exactly. Since the (total) density fluctuation operator $\rho(\mathbf{q})$ is given by the sum $\sum_\alpha \rho_\alpha(\mathbf{q})$ for $|\mathbf{q}| \ll \Lambda_\alpha$, the dynamical density-density response for small $\mathbf{q}$ is thus easily obtained [68, 69]. The result is the same as that obtained in subsection 5.2.1 as a corollary of loop cancellation: RPA.

Using the bosonic representation of fermionic operators, the single-particle propagator can be expressed in terms of the expectation value

$$\langle b_\alpha(q) b^\dagger_\alpha(q) \rangle = \frac{i}{\omega - \mathbf{v}_\alpha \cdot \mathbf{q} + i0^+ s(\omega)} + \frac{\Lambda_\alpha^{d-1}}{(2\pi)^d} \, \mathbf{n}_\alpha \cdot \mathbf{q} \, \frac{i D_{\alpha\alpha}(q)}{[\omega - \mathbf{v}_\alpha \cdot \mathbf{q} + i0^+ s(\omega)}]^2 \tag{5.64}$$

where $D_{\alpha\alpha'}(q)$ is obtained from the RPA effective interaction $D_{\mathbf{k}\mathbf{k}'}(q)$, eq. (5.21), by averaging over $\mathbf{k} \in \partial\mathcal{F}_\alpha$ and $\mathbf{k}' \in \partial\mathcal{F}_{\alpha'}$. The propagator $G(p)$ is then obtained [69, 70] as the Fourier transform of

$$G_\alpha(t, \mathbf{r}) = \frac{\delta_{\Lambda_\alpha^{-1}}(\mathbf{r}_t) \, e^{i \mathbf{k}_{F\alpha} \cdot \mathbf{r}}}{\mathbf{n}_\alpha \cdot \mathbf{r} - v_\alpha t + i0^+ s(t)} \, \exp\left[\int_q [e^{i(\mathbf{q}\cdot\mathbf{r} - \omega t)} - 1] \, \frac{i D_{\alpha\alpha}(q)}{[\omega - \mathbf{v}_\alpha \cdot \mathbf{q} + i0^+ s(\omega)]^2} \right] \tag{5.65}$$

for $\mathbf{p} \in K_\alpha$, where $\mathbf{r}_t$ is the component of $\mathbf{r}$ orthogonal to $\mathbf{n}_\alpha$ and $\delta_{\Lambda^{-1}}(.)$ is a broadened δ-function with width Λ^{-1}. The behavior of $G(p)$ close to the Fermi surface is determined by $G_\alpha(t, \mathbf{r})$ for large $\mathbf{r}$; the tangential component $\mathbf{r}_t$, being limited by Λ_α^{-1}, can thus be set to zero [70]. Decomposing $\mathbf{q}$ in normal and tangential components as in 5.2.3, one thus recovers the result (5.46)-(5.51) for G obtained from the asymptotic Ward identities. An advantage of the Ward identity approach is that it avoids the patch construction with its somewhat artificial complications due to overcompleteness of states and inter-patch scattering processes.

An alternative formulation of d-dimensional bosonization via functional integrals has been proposed by Kopietz and Schönhammer [73], who generalized earlier work on functional bosonization of the Luttinger model [104, 105] to higher dimensions. Unlike the Hamiltonian formulation the path integral language is applicable also to retarded interactions. In addition, the path integral approach seems to be also more suitable for a discussion of corrections to the leading (free boson) results.

Starting point is an action for interacting fermions written as

$$S[\psi, \psi^*] = \int_k \psi^*_k (i k_0 - \xi_{\mathbf{k}}) \psi_k - \tfrac{1}{2} \sum_{\alpha\alpha'} \int_q g_{\alpha\alpha'}(q) \, \rho_\alpha(q) \rho_{\alpha'}(-q) \tag{5.66}$$

where

$$\rho_\alpha(q) = \int_k \chi_\alpha(\mathbf{k})\, \psi^*_{k-q/2}\psi_{k+q/2} \tag{5.67}$$

and $g_{\alpha\alpha'}(q)$ is an average of $g_{\mathbf{k}\mathbf{k}'}(q)$ over $\mathbf{k} \in \partial\mathcal{F}_\alpha$ and $\mathbf{k}' \in \partial\mathcal{F}_{\alpha'}$. As long as the variation of $g_{\mathbf{k}\mathbf{k}'}(q)$ as a function of $\mathbf{k}$ and $\mathbf{k}'$ is small on the scale set by Λ_α, this averaging does not introduce significant errors. Note that in (5.67) only $\mathbf{k}$ is restricted to the sector K_α, while in (5.54) both fermionic momenta $\mathbf{k} \pm \mathbf{q}/2$ had to be in K_α.

Introducing an auxiliary field $\phi_\alpha(q)$ that mediates the interaction, the partition function can be written as a functional integral over ψ_k, ψ^*_k and $\phi_\alpha(q)$, with an action [62, 73]

$$S[\psi,\psi^*,\phi] = \int_k \psi^*_k(ik_0 - \xi_{\mathbf{k}})\psi_k - i\sum_\alpha \int_q \phi_\alpha(-q)\rho_\alpha(q) - \tfrac{1}{2}\sum_{\alpha\alpha'}\int_q [g^{-1}(q)]_{\alpha\alpha'}\phi_\alpha(q)\phi_{\alpha'}(-q) \tag{5.68}$$

where $g^{-1}(q)$ is the inverse of the matrix $g(q)$ defined by the matrix elements $g_{\alpha\alpha'}(q)$.

The integration over the fermionic fields ψ and ψ^* is now Gaussian and can thus be carried out exactly. This yields an effective action [62, 73]

$$S[\phi] = S_{kin}[\phi] - \tfrac{1}{2}\sum_{\alpha\alpha'}\int_q [g^{-1}(q)]_{\alpha\alpha'}\phi_\alpha(q)\phi_{\alpha'}(-q) \tag{5.69}$$

where

$$S_{kin}[\phi] = \mathrm{tr}\log[\hat{1} - \hat{G}_0\hat{\Phi}] \tag{5.70}$$

Here the trace is over all momenta and frequencies, and $\hat{G}_0$ and $\hat{\Phi}$ are infinite matrices in momentum-frequency space, with matrix elements given by $[\hat{G}_0]_{kk'} = \delta_{kk'}G_0(k)$ and $[\hat{\Phi}]_{kk'} = i\sum_\alpha \chi_\alpha(\mathbf{k})\, \phi_\alpha(k-k')$. Expanding the logarithm, the kinetic part of the action can also be written as an infinite sum $S_{kin}[\phi] = \sum_{n=1}^\infty S_{kin,n}[\phi]$ where

$$S_{kin,n}[\phi] = -\frac{1}{n}\mathrm{tr}[\hat{G}_0\hat{\Phi}]^n =: \frac{1}{n}\sum_{\alpha_1,\ldots,\alpha_n}\int_{q_1,\ldots,q_n} U_n^{\alpha_1,\ldots,\alpha_n}(q_1,\ldots,q_n)\,\phi_{\alpha_1}(q_1)\ldots\phi_{\alpha_n}(q_n) \tag{5.71}$$

Diagrammatically, the n-th order term corresponds to a fermionic loop with n insertions due to fields $\phi_{\alpha_1}(q_1),\ldots,\phi_{\alpha_n}(q_n)$ summed over all possible permutations.

In general, $S_{kin}[\phi]$ is a complicated functional, containing arbitrarily high powers in ϕ. Significant simplifications occur however in systems dominated by forward scattering, where only interactions with small momentum transfers are important. In that case the contribution from fields $\phi_\alpha(q)$ with large $\mathbf{q}$ is negligible, since $\phi_\alpha(q)$ mediates interactions with momentum transfer $\mathbf{q}$. As a consequence of the *loop cancellation* discussed already in 5.2.1, loops with more than two insertions are irrelevant for small $\mathbf{q}$ to leading order in $\mathbf{q}$, and the series expansion of $S_{kin}[\phi]$ can be truncated at second order. The first order term vanishes for all $q \neq 0$, while the quadratic part yields a kinetic term [62, 73]

$$S_{kin}[\phi] = \tfrac{1}{2}\sum_{\alpha\alpha'}\int_q \Pi_0^\alpha(q)\,\phi_\alpha(q)\phi_\alpha(-q) \tag{5.72}$$

where $\Pi_0^\alpha(q)$ is the bare polarisation bubble with momenta on the fermion lines restricted to the sector K_α, i.e.

$$\Pi_0^\alpha(q) = \frac{\Lambda_\alpha^{d-1}}{(2\pi)^d} \frac{\mathbf{n}_\alpha \cdot \mathbf{q}}{iq_0 - \mathbf{v}_\alpha \cdot \mathbf{q}} \tag{5.73}$$

The q-integral in (5.72) and in subsequent results is restricted by a cutoff function $\chi_\alpha(\mathbf{k}_{F\alpha} + \mathbf{q})$, which is chosen such that the degrees of freedom are correctly counted [106]. For the one-dimensional Luttinger model the truncation leading to (5.72) is exact to all orders in $\mathbf{q}$ [105].

Due to the loop cancellation, the action $S[\phi]$ is now quadratic in the field ϕ, i.e. its partition function can be calculated exactly.

Instead of ϕ, one may also choose its conjugate field, the density fluctuation field $\tilde{\rho}$ as dynamical variable, especially to make contact with the Hamiltonian version of bosonization. One then obtains an effective action [62, 73]

$$\tilde{S}[\tilde{\rho}] = \tfrac{1}{2} \sum_{\alpha\alpha'} \int_q \left\{ [\Pi_0^\alpha(q)]^{-1} \delta_{\alpha\alpha'} - g_{\alpha\alpha'}(q) \right\} \tilde{\rho}_\alpha(q) \tilde{\rho}_{\alpha'}(-q) \tag{5.74}$$

which is a quadratic form in the bosonic collective field $\tilde{\rho}_\alpha(q)$. For frequency-independent interactions the action $\tilde{S}[\tilde{\rho}]$ is equivalent to the bosonized Hamiltonian in Eq. (5.63). For the small-$\mathbf{q}$ density-density response, which can be expressed as an expectation value $\sum_{\alpha\alpha'} \langle \tilde{\rho}_\alpha(q) \tilde{\rho}_{\alpha'}(-q) \rangle$ with the action $\tilde{S}[\tilde{\rho}]$, one obviously recovers the RPA-result.

The single-particle propagator G has been calculated by Kopietz et al. [73, 106] directly from $S[\phi]$, without introducing the $\tilde{\rho}$-field. The expectation value $\langle \psi \psi^* \rangle$ can be written as a functional integral over ψ, ψ^* and ϕ with the action $S[\psi, \psi^*, \phi]$ in (5.68). Performing the Gaussian integral over ψ and ψ^*, the propagator $G(k)$ is obtained as an average

$$G(k) = \langle [\hat{G}]_{kk} \rangle_{S[\phi]} \tag{5.75}$$

Here $\hat{G}$ is an infinite matrix in momentum-frequency space, defined by the Dyson equation $[\hat{G}^{-1}]_{kk'} = [\hat{G}_0^{-1}]_{kk'} - [\hat{\Phi}]_{kk'}$, where $[\hat{G}_0^{-1}]_{kk'} = (2\pi)^{d+1} \delta(k - k') G_0(k)$, and $S[\phi]$ is given by (5.69). The matrix $\hat{G}$ is the propagator of a non-interacting system in a dynamical random field ϕ. If $\phi_\alpha(q)$ is finite only for small momenta $\mathbf{q}$, the diagonal elements $[\hat{G}]_{kk}$ can be expressed explicitly as [106]

$$[\hat{G}]_{kk} = \sum_\alpha \chi_\alpha(\mathbf{k}) \int d\mathbf{r} \int_0^\beta d\tau \, e^{-i(\mathbf{k} - \mathbf{k}_{F\alpha})\mathbf{r} + ik_0\tau} \exp\left[i \int_q \frac{e^{i(\mathbf{q}\cdot\mathbf{r} - q_0\tau)} - 1}{iq_0 - \mathbf{v}_\alpha \cdot \mathbf{q}} \phi_\alpha(q) \right] G_0^\alpha(\tau, \mathbf{r}) \tag{5.76}$$

where $G_0^\alpha(\tau, \mathbf{r})$ is the Fourier transform of $(ik_0 - \mathbf{v}_\alpha \cdot \mathbf{k})^{-1}$. Approximating $S_{kin}[\phi]$ by the quadratic form in (5.72), the calculation of $G(k)$ from (5.75) has thus been reduced to a Gaussian integral over ϕ, that can be carried out explicitly [73, 106]. The result is

equivalent to the result (5.65) from the Hamiltonian version of bosonization and also to the one obtained from asymptotic Ward identities.

We emphasize that d-dimensional bosonization, like the asymptotic Ward identity method, can treat only scattering processes with small momentum transfers $\mathbf{q}$. The results are meaningful only if scattering processes with large momentum transfers are negligible with respect to forward scattering. The calculation of corrections to the leading order result is difficult. Kopietz et al. [62] have analyzed certain corrections to the RPA result for the density-density response function by including non-quadratic terms in the expansion of the effective kinetic action $S_{kin}[\phi]$. However, scattering processes into other sectors K_α seem to be hard to treat, and thus only a partial contribution to the subleading corrections in a small-$\mathbf{q}$ expansion could be calculated explicitly. Expressed as a correction to the polarization insertion Π, this contribution turned out to be linear in the RPA effective interaction D. Corrections to loop cancellation (and other asymptotic Ward identities) to finite order in D can of course also be calculated from the corresponding Feynman diagrams.

6. ONE-DIMENSIONAL LUTTINGER LIQUID

In one-dimensional interacting Fermi systems Fermi liquid theory is not valid. Its breakdown is signalled already in second order perturbation theory. The perturbatively calculated reduction of the quasi-particle weight by interactions diverges logarithmically at the Fermi surface. The problem of treating these divergencies has been first solved by a weak coupling renormalization group method applied to an effective low-energy theory known as *"g-ology"* model [20]. Assuming a scaling ansatz for the vertex functions, one approaches the low-energy regime by rescaling the fields and the coupling constants (a small number of "g's"). The consistency of the ansatz is verified order by order in perturbation theory. Depending on the values of the bare couplings the renormalized couplings flow either to strong coupling, and hence out of the perturbatively controlled regime, or to a fixed point Hamiltonian, the exactly soluble Luttinger model [22, 21]. In the latter case the system is a *"Luttinger liquid"*, i.e. a normal (not symmetry-broken) metallic phase characterized by i) a continuous momentum distribution with a power-law singularity at the Fermi surface, the exponent η being non-universal; ii) a single-particle density of states which vanishes as ω^η near the Fermi energy, i.e. absence of fermionic quasi-particles; iii) finite charge and spin density response for long wavelengths and the existence of collective bosonic charge and spin density modes; iv) power-law singularities in various superconducting and short wave length (e.g. $|\mathbf{q}| = 2k_F$) density correlation functions; v) separation of spin and charge degrees of freedom [20, 23, 24].

Luttinger liquid behavior is not confined to special weak-coupling models but may also govern strongly interacting systems. Although in the latter case a perturbative renormalization group calculation of the correlation functions is not adequate, the low-energy properties are still uniquely characterized by a small number of parameters, which are directly related to simple physical quantities. This leap beyond weak coupling, which follows the spirit of Fermi liquid theory, has been pioneered by Haldane [25], who also introduced the suggestive term "Luttinger liquid". The one-dimensional Hubbard model, which is exactly soluble by the Bethe ansatz method [107], is most probably a Luttinger liquid for *any* coupling strength, except for half-filling [108, 109, 110, 111]. This conjecture is well established for weak coupling, while for strong coupling it is supported by the structure of the low lying excitations [108, 109], the absence of non-analyticities in the exact ground state energy at finite coupling [111], by studies of the infinite coupling limit

[112], as well as by numerical evaluations of finite systems [113]. The Luttinger liquid parameters (and hence all critical exponents) for the one-dimensional Hubbard model have been obtained exactly for arbitrary coupling strength [108, 109, 110, 111].[17]

Fermi liquid theory and one-dimensional Luttinger liquid theory are commonly formulated in two distinct languages. Fermi liquid theory (i.e. its microscopic basis) is derived in terms of Feynman diagrams and Ward identities [2, 14], while Luttinger liquid theory is usually worked out using the bosonization technique [22, 25, 65, 66]. Nevertheless, it has often been emphasized that both liquids are in many respects very similar [115]. In fact Luttinger liquid theory can also be constructed by standard many-body techniques, without bosonization, by exploiting the Ward identities associated with the peculiar conservation laws in one-dimensional Fermi systems [61, 116, 86]. The choice of a common language for Fermi and Luttinger liquid theory allows for a direct comparison of both theories, making common features and differences more clear.

The use of Ward identities makes evident the crucial role played by conservation laws in one-dimensional Fermi systems (especially in Luttinger liquids). In addition to the usual charge and spin conservation, the discrete structure of the Fermi surface in one dimension allows for a more stringent conservation law: separate charge (and possibly spin) conservation in low-energy-scattering processes for particles near the left and right Fermi points, respectively. Only the *separate* conservation of charge (spin) on each Fermi point guarantees the requirement of finite charge (spin) density response in "normal" one-dimensional metals. The velocities associated with the corresponding conserved currents provide a complete parametrization of the low energy physics [86]. The conservation of charge and spin on each Fermi point in a one-dimensional Luttinger liquid implies velocity conservation as defined and discussed for d-dimensional systems in Sec. 5.

There are several good reviews on one-dimensional Fermi systems: The understanding reached by the end of the seventies has been reviewed by Solyom [20] and Emery [117]. Recent up-to-date reviews have been presented by Voit [23] and Schulz [24]. The following is not a comprehensive review on one-dimensional Luttinger liquids. Our main goal here is to show how Luttinger liquid theory fits in with the more general framework constructed for d-dimensional systems in Secs. 2,3 and 5.

6.1. THE G-OLOGY MODEL

The generic low-energy physics of a one-dimensional Fermi system with short-range interactions is incorporated in the *g-ology model* [20], a continuum model with a linear

[17] Note, however, that the low-energy scale controlled by Luttinger liquid behavior may shrink for increasing coupling strength; this has been demonstrated very clearly for the Hubbard model by Penc, Mila and Shiba [114].

dispersion relation and two-particle interactions. The kinetic term

$$H_0 = \sum_{\mathbf{k},\sigma}^{<\Lambda} v_F k_r \, a^\dagger_{\mathbf{k}\sigma} a_{\mathbf{k}\sigma} \tag{6.1}$$

describes a linear band as illustrated in Fig. 6.1. Here $k_r = |\mathbf{k}| - k_F$ measures the distance from the Fermi "surface" (consisting of two Fermi points in one dimension) and the momentum sum is restricted by $|k_r| < \Lambda$. We keep vector notation (bold face) for momenta even in one dimension to distinguish them from bi-vectors $k = (k_0, \mathbf{k})$ which include the energy variable k_0. The only parameter characterizing H_0 is the Fermi velocity v_F, i.e. the slope of the band.

The interaction H_I is given by

$$H_I = H_1 + H_2 + H_3 + H_4 \tag{6.2}$$

where

$$H_1 = \tfrac{1}{V} \sum_{\mathbf{q}} \sum_{\sigma\sigma'} g_1^{\sigma\sigma'} \rho^+_\sigma(\mathbf{q}) \rho^-_{\sigma'}(-\mathbf{q}) \tag{6.3a}$$

$$H_2 = \tfrac{1}{V} \sum_{\mathbf{q}} \sum_{\sigma\sigma'} g_2^{\sigma\sigma'} \rho_{+,\sigma}(\mathbf{q}) \rho_{-,\sigma'}(-\mathbf{q}) \tag{6.3b}$$

$$H_3 = \tfrac{1}{2V} \sum_{\mathbf{q}} \sum_{\sigma\sigma'} g_3^{\sigma\sigma'} [\rho^+_\sigma(\mathbf{q}) \rho^+_{\sigma'}(-\mathbf{q}) + \rho^-_\sigma(\mathbf{q}) \rho^-_{\sigma'}(-\mathbf{q})] \tag{6.3c}$$

$$H_4 = \tfrac{1}{2V} \sum_{\mathbf{q}} \sum_{\sigma\sigma'} g_4^{\sigma\sigma'} [\rho_{+,\sigma}(\mathbf{q}) \rho_{+,\sigma'}(-\mathbf{q}) + \rho_{-,\sigma}(\mathbf{q}) \rho_{-,\sigma'}(-\mathbf{q})] \tag{6.3d}$$

where V is the volume of the system and

$$\rho_{\alpha\sigma}(\mathbf{q}) = \sum_{\mathbf{k}}^{<\Lambda} \chi_\alpha(\mathbf{k}-\mathbf{q}/2) \, \chi_\alpha(\mathbf{k}+\mathbf{q}/2) \, a^\dagger_{\mathbf{k}-\mathbf{q}/2,\sigma} a_{\mathbf{k}+\mathbf{q}/2,\sigma} \tag{6.4a}$$

$$\rho^\pm_\sigma(\mathbf{q}) = \sum_{\mathbf{k}}^{<\Lambda} \chi_\pm(\mathbf{k}-\mathbf{q}/2) \, \chi_\mp(\mathbf{k}+\mathbf{q}/2) \, a^\dagger_{\mathbf{k}-\mathbf{q}/2,\sigma} a_{\mathbf{k}+\mathbf{q}/2,\sigma} \tag{6.4b}$$

Here $\chi_\alpha(\mathbf{k})$ is the characteristic function of the two *sectors* K_α in momentum space corresponding to *right*-moving ($\alpha = +$) and *left*-moving ($\alpha = -$) particles (cf. Sec. 5.4). Below we will often use the spin symmetric and antisymmetric linear combinations

$$\rho_\alpha(\mathbf{q}) = \rho_{\alpha\uparrow}(\mathbf{q}) + \rho_{\alpha\downarrow}(\mathbf{q}) \quad \text{and} \quad s^z_\alpha(\mathbf{q}) = [\rho_{\alpha\uparrow}(\mathbf{q}) - \rho_{\alpha\downarrow}(\mathbf{q})]/2 \tag{6.5}$$

All parts of the Hamiltonian are understood to be normal ordered with respect to the ground state of H_0, i.e. the Fermi sea. The interaction H_I generates the various types of scattering processes listed in Fig. 6.2: small momentum transfer processes (H_2 and

H_4), back-scattering (H_1) and umklapp-scattering (H_3).[18] The coupling constants may be spin-dependent:

$$g_i^{\sigma\sigma'} = g_{i\parallel}\delta_{\sigma\sigma'} + g_{i\perp}\delta_{\sigma,-\sigma'} , \quad i = 1, \ldots, 4 \tag{6.6}$$

i.e. $g_{i\parallel}$ refers to parallel and $g_{i\perp}$ to opposite spins, respectively. For later reference we define symmetric and antisymmetric linear combinations

$$g_i^c = (g_{i\parallel} + g_{i\perp})/2 \quad \text{and} \quad g_i^s = (g_{i\parallel} - g_{i\perp})/2 \tag{6.7}$$

where the indices "c" and "s" are abbreviations for "charge" and "spin", respectively.

The terms $H_{1\parallel}$ and $H_{2\parallel}$ describe obviously the same process (since fermions with equal spin projection are indistinguishable). Hence, we can (and will) set $g_{1\parallel} = 0$ without loss of generality. Due to momentum conservation, umklapp-processes (H_3) can be relevant only if $4k_F$ is equal to a reciprocal lattice vector (so that all scattering particles can be near the Fermi points), as is the case, for example, for a Hubbard model at half-filling.

6.2. RENORMALIZATION GROUP AND CUTOFFS

The g-ology model is the generic effective low-energy theory for one-dimensional interacting Fermi systems. A g-ology Hamiltonian (or action) is the general outcome of a Wilson renormalization procedure (integrate out far-from-Fermi-surface momentum states) applied to a microscopic one-dimensional model, such as the one-dimensional Hubbard model. Non-linear dependences in the band structure and momentum-dependences in the couplings $g_1 \ldots g_4$ are irrelevant for the leading low-energy behavior. The model parameters are related to the effective low-energy action $\bar{S}^\Lambda$ from Sec. 2 as follows:

$$\begin{aligned} v_F &= v_F^\Lambda \\ g_1 &= \bar{\Gamma}^\Lambda(\mathbf{k}_F, -\mathbf{k}_F; -\mathbf{k}_F, \mathbf{k}_F) \\ g_2 &= \bar{\Gamma}^\Lambda(\mathbf{k}_F, -\mathbf{k}_F; \mathbf{k}_F, -\mathbf{k}_F) \\ g_3 &= \bar{\Gamma}^\Lambda(-\mathbf{k}_F, -\mathbf{k}_F; \mathbf{k}_F, \mathbf{k}_F) \\ g_4 &= \bar{\Gamma}^\Lambda(\mathbf{k}_F, \mathbf{k}_F; \mathbf{k}_F, \mathbf{k}_F) \end{aligned} \tag{6.8}$$

Here we have suppressed spin variables and energy variables (which are all set equal to zero) in the vertex functions.

Perturbation theory for one-dimensional Fermi systems is plagued by several logarithmic infrared divergencies in one- and two-particle correlation functions, which cannot be treated by a simple resummation of certain subsets of Feynman diagrams. A first understanding and successful treatment of these divergencies has been achieved in the 1970's

[18] Note that in H_1 and H_3 the momentum transfer is $|\mathbf{q}| \sim 2k_F$.

by Solyom and coworkers [20] by applying perturbative renormalization group methods to the g-ology model. Making a scaling ansatz for the vertex functions, he approached the low-energy limit by rescaling fields and coupling constants. The consistency of the ansatz has been verified a posteriori order by order in perturbation theory. One may view this approach as a slightly modified version of the field-theoretic renormalization group [118, 119]; consistency of the scaling ansatz is guaranteed by the renormalizability of the g-ology model [86, 120]. Depending on the values (and signs) of the bare couplings, and on $4k_F$ being equal or different from a reciprocal lattice vector, the renormalized couplings may either follow a run-away trajectory or reach a fixed point (at weak coupling for weak bare interactions). The former behavior signals a dramatic change of the effective low-energy Hamiltonian (or action) compared to the bare one, usually associated with dynamical generation of gaps [121, 20]. In the latter one has reached a fixed point Hamiltonian given by the Luttinger model [21], where only interactions with small momentum transfers survive, i.e. $g_1^* = g_3^* = 0$. The Luttinger model is exactly soluble [22], and the low-energy physics associated with this fixed point can be calculated without further use of perturbative renormalization group methods (see Section 6.4).

It is however important to realize that all the peculiarities of Luttinger liquid behavior have clear perturbative signals. An instructive example is the anomalous scaling behavior of the single-particle propagator

$$G(sk_0, sk_r) = s^{\eta-1} G(k_0, k_r) \tag{6.9}$$

where $\eta > 0$ is a (non-universal) constant, which implies a power-law singularity of the momentum distribution function near $\mathbf{k}_F$ and a density of states vanishing as ω^η. This anomalous scaling is signalled by a logarithmic singularity in the perturbatively calculated quasi-particle weight $Z = [1 - \partial\Sigma/\partial(ik_0)]^{-1}$ at the Fermi level (cf. eq. (2.24)) in second order perturbation theory. Within a perturbative renormalization group scheme one obtains a wave function renormalization obeying

$$\frac{d \log Z^\Lambda}{d \log \Lambda} \;=\; \frac{(\bar{g}_{2\|}^*)^2 + (\bar{g}_{2\perp}^*)^2}{8\pi^2 v_F^2} + \mathcal{O}(\bar{g}^{*3}) \tag{6.10}$$

for the Luttinger model (where $g_1 = g_3 = 0$), where the quadratic terms are obtained from 2-loop contributions to the self-energy. In the low-energy limit, one has thus

$$Z^\Lambda \to 0 \quad \text{for} \quad \Lambda \to 0 \tag{6.11}$$

Integrating the flow for Z^Λ one obtains anomalous scaling behavior as in (6.9) with an anomalous dimension

$$\eta \;=\; \frac{(\bar{g}_{2\|}^*)^2 + (\bar{g}_{2\perp}^*)^2}{8\pi^2 v_F^2} + \mathcal{O}(\bar{g}^{*3}) \tag{6.12}$$

By constrast, in a Fermi liquid $Z^\Lambda \to Z > 0$, and there is no anomalous scaling dimension (i.e. $\eta = 0$). In both (Fermi and Luttinger) liquids the long wavelength response-functions do not acquire anomalous scaling dimensions and both describe a normal metallic phase.

Note that concrete perturbative renormalization group calculations for the Luttinger or g-ology model are easiest within the field-theoretic version of the RG or, what is almost the same, by using Solyom's scaling ansatz. Wilson's RG version becomes relatively cumbersome in practical calculations beyond one-loop order, while characteristic differences between Luttinger and Fermi liquid behavior show up only at two-loop order.

The effective g-ology model obtained by integrating out high-energy states has a *band-width cutoff* Λ, and no explicit momentum transfer cutoff. Momentum transfers are restricted only indirectly by the finite band-width cutoff. For a non-perturbative analysis it is however more convenient to restrict the momentum transfers $\mathbf{q}$ in the interaction part (6.3) to values $|\mathbf{q}| < \Lambda_q$ in H_2 and H_4 and $||\mathbf{q}| - 2k_F| < \Lambda_q$ in H_1 and H_3, with a *"momentum transfer cutoff"* $\Lambda_q \ll \Lambda$ [20]. This can be done either ad hoc, or via the mapping described in the introductory part of Sec. 5. The qualitative structure of the infrared asymptotics is of course not altered by different choices of the cutoff procedure. In practice not even quantitative results depend on the cutoff procedure, if the low-energy physics is expressed in terms of parameters which are directly related to physical observables.

Note that the scale k_F does not appear in the g-ology model. Important is only the *existence* of the two Fermi points, not their *distance* in momentum space. Thus the inverse cutoff Λ_q^{-1} is the only length scale in the model; Λ_q replaces the natural cutoff given by the Brillouin zone boundary or by non-linear terms in $\epsilon_{\mathbf{k}}$ in the underlying microscopic system.

6.3. CHARGE/SPIN CONSERVATION AND WARD IDENTITIES

6.3.1. Global charge and spin conservation

The g-ology Hamiltonian conserves charge and the spin component in z-direction (the quantization axis). Spin components in any direction are conserved only if the coupling constants satisfy the relation $g_{2\perp}$–$g_{1\perp} = g_{2\parallel}$–$g_{1\parallel}$, which can be obtained from the condition $[H, \mathbf{S}] = 0$. As discussed in detail in Secs. 3 and 5, such conservation laws give rise to important Ward identities.

In general, the g-ology interactions are not of density-density type. The current operators depend therefore explicitly on the coupling constants. We begin with charge (or particle) conservation and its consequences, and address spin conservation briefly afterwards. Calculating the commutator of the g-ology Hamiltonian with the charge density

fluctuation operator at small $\mathbf{q}$, i.e. $\rho(\mathbf{q}) = \rho_+(\mathbf{q}) + \rho_-(\mathbf{q})$, one obtains a continuity equation with a current operator given by [86]

$$\mathbf{j}(\mathbf{q}) = v_c \left[\rho_+(\mathbf{q}) - \rho_-(\mathbf{q})\right] \tag{6.13}$$

with a coupling-dependent velocity

$$v_c = v_F + (g_4^c - g_2^c)/\pi \tag{6.14}$$

The current-operator depends only on *forward* scattering couplings, because $\rho(\mathbf{q})$ commutes with H_1 (with $g_{1\|} = 0$) and H_3. Hence, we can now simply apply the Ward identities derived from global charge conservation for the d-dimensional forward scattering action in Sec. 5.1 to the one-dimensional case. In particular, the correlation function $J^{\mu\nu} = -\frac{1}{V}\langle j^\mu j^\nu\rangle$ and the current vertex $\Gamma^\mu = \langle j^\mu a a^\dagger\rangle_{tr}$, obey the Ward identities (3.10) and (3.15), respectively [86], with a function $\mathbf{c}(\mathbf{q})$ given by

$$\mathbf{c}(\mathbf{q}) = \frac{2}{\pi} v_c \, \mathbf{q} \tag{6.15}$$

for the g-ology model. Note that the formula (3.13) for $\mathbf{c}(\mathbf{q})$ does not apply here, as a consequence of the band-width cutoff; the more general expression (3.12) yields a finite result due to this cutoff, not as a consequence of a finite gradient of $\mathbf{v_k}$ in momentum space. The irreducible current vertex $\Lambda^\mu_\sigma(p;q) = \langle j_0^\mu(q)\, a_{p-q/2,\sigma}\, a^\dagger_{p+q/2,\sigma}\rangle^{irr}_{tr}$ with the non-interacting current operator

$$\mathbf{j}_0(\mathbf{q}) = v_F \left[\rho_+(\mathbf{q}) - \rho_-(\mathbf{q})\right] \tag{6.16}$$

obeys the Ward identity (5.15). Note that $\mathbf{j}_0$ has been constructed with v_F, i.e. the bare velocity in the g-ology model.

Conservation of the spin component in z-direction has analogous consequences. A continuity equation relates the z-component of the spin density $s^z(\mathbf{q}) = s^z_+(\mathbf{q}) + s^z_-(\mathbf{q})$ to a spin current given by

$$\mathbf{j}^z(\mathbf{q}) = v_s \left[s^z_+(\mathbf{q}) - s^z_-(\mathbf{q})\right] \tag{6.17}$$

with a velocity

$$v_s = v_F + (g_4^s - g_2^s)/\pi \tag{6.18}$$

Again only forward scattering couplings are involved. The spin correlation function $J^{z\mu,z\nu} = -\frac{1}{V}\langle j^{z\mu} j^{z\nu}\rangle$ and the spin current vertex $\Gamma^{z\mu} = \langle j^{z\mu} a a^\dagger\rangle_{tr}$, obey the Ward identities (3.27) and (3.29), respectively [86], with $\mathbf{c}(\mathbf{q})$ given by (6.15). The irreducible spin current vertex $\Lambda^{z\mu}_{\sigma\sigma'} = \langle j_0^{z\mu} a_\sigma a^\dagger_{\sigma'}\rangle^{irr}_{tr}$ with the non-interacting spin current operator

$$\mathbf{j}^z_0(\mathbf{q}) = v_F \left[s^z_+(\mathbf{q}) - s^z_-(\mathbf{q})\right] \tag{6.19}$$

obeys the Ward identity (5.16) for $a = z$.

6.3.2. Separate left/right conservation laws

In the absence of umklapp processes, there is an additional conservation law: *charge near the left and right Fermi point is conserved separately.* This case is by no means academic since the presence of umklapp processes, being subject to the matching condition $4\mathbf{k}_F = \mathbf{Q}$ ($\mathbf{Q}$ a reciprocal lattice vector), is the exception rather than the rule. For example, the g-ology describing the low-energy physics of the Hubbard model involves umklapp terms only at half-filling. The separate left/right charge conservation yields additional (besides those implied by usual charge conservation) significant constraints on the structure of correlation functions and the renormalization group [86, 120].

Separate *left/right spin conservation* is spoiled by the back-scattering process $H_{1\perp}$, which is generically present in a model of spin-$\frac{1}{2}$ fermions. However, in many cases of interest (in particular in the one-dimensional Hubbard model with repulsive interaction [24]) the back-scattering amplitude scales to zero at low energies, *restoring* thus separate left/right spin conservation asymptotically, and yielding further constraints on the asymptotic low-energy theory. This is indeed the case of the Luttinger liquid discussed in the next Section 6.4.

Let us start by exploring the consequences of separate conservation of charge on each Fermi point. In addition to conservation of the global charge, i.e. left and right summed, separate left/right conservation implies also the conservation of the charge *difference*, i.e. right minus left. Correspondingly, the operator

$$\tilde{\rho}(\mathbf{q}) = \rho_+(\mathbf{q}) - \rho_-(\mathbf{q}) \tag{6.20}$$

obeys a continuity equation $\partial_\tau \tilde{\rho}(\tau, \mathbf{q}) = [H, \tilde{\rho}(\tau, \mathbf{q})] = -\mathbf{q}\cdot\tilde{\mathbf{j}}(\tau, \mathbf{q})$ with a current operator given by [86]

$$\tilde{\mathbf{j}}(\mathbf{q}) = \tilde{v}_c \left[\rho_+(\mathbf{q}) + \rho_-(\mathbf{q})\right] \tag{6.21}$$

and a velocity

$$\tilde{v}_c = v_F + (g_4^c + g_2^c)/\pi \tag{6.22}$$

In analogy to the Thirring model one may refer to $\tilde{j}^\mu = (\tilde{\rho}, \tilde{\mathbf{j}})$ as the *axial charge* current [122].

The continuity equation for $\tilde{j}^\mu$ implies the Ward identity

$$(iq_0, \mathbf{q})_\mu \tilde{J}^{\mu\nu}(q) = \tilde{c}^\nu(\mathbf{q}) = \delta_{\nu 1} \frac{2}{\pi} \tilde{v}_c \, \mathbf{q} \tag{6.23}$$

for $\tilde{J}^{\mu\nu}(q) = -\frac{1}{V}\langle \tilde{j}^\mu \tilde{j}^\nu \rangle$ and

$$(iq_0, \mathbf{q})_\mu \tilde{\Gamma}^\mu_\sigma(p; q) = s(\mathbf{p}) \left[G_\sigma^{-1}(p+q/2) - G_\sigma^{-1}(p-q/2) \right] \tag{6.24}$$

for $\tilde{\Gamma}^\mu = \langle \tilde{j}^\mu a a^\dagger \rangle_{tr}$, where $s(\mathbf{p})$ is the sign of $\mathbf{p}$ (i.e. $s(\mathbf{p}) = 1$ for right-moving particles and $s(\mathbf{p}) = -1$ for left-moving particles).

Combining the Ward identities from global and axial conservation laws determines $J^{\mu\nu}$ (and $\tilde{J}^{\mu\nu}$) completely. Results will be discussed in 6.4. Note that the velocities v_c and $\tilde{v}_c$ appearing in the two conserved currents correspond to Haldane's [25] velocities v_J and v_N, which control the current (J) and density (N) excitations (cf. eqs. (6.32) and (6.34)).

In the absence of backscattering, the z-component of spin is conserved separately on each Fermi point. In that case one has an additional continuity equation for an *axial spin* current $\tilde{j}^{z\mu} = (\tilde{s}^z, \tilde{\mathbf{j}}^z)$ where

$$\tilde{s}^z(\mathbf{q}) = s^z_+(\mathbf{q}) - s^z_-(\mathbf{q}) \tag{6.25}$$

and

$$\tilde{\mathbf{j}}^z(\mathbf{q}) = \tilde{v}_s \left[s^z_+(\mathbf{q}) + s^z_-(\mathbf{q}) \right] \tag{6.26}$$

with a velocity

$$\tilde{v}_s = v_F + (g^s_4 + g^s_2)/\pi \tag{6.27}$$

This yields Ward identities for correlation and vertex functions in complete analogy to those from axial charge conservation. The identities from global and axial spin conservation determine $J^{z\mu,z\nu}$. Results will be discussed in 6.4.

6.4. LUTTINGER LIQUID FIXED POINT

In many one-dimensional Fermi systems both umklapp- and backscattering are irrelevant in the low-energy limit.[19] In these cases only *forward scattering*, i.e. scattering with small momentum transfers survives. The corresponding couplings g_2 and g_4 reach finite non-universal fixed points g_2^* and g_4^* in the low-energy limit [20]. The β-function at the fixed points vanishes due to cancellations imposed by the peculiar left/right conservation laws [86, 120], not simply as a consequence of vanishing phase space as in a Fermi liquid!

The fixed point Hamiltonian can be written as

$$H^* = H_0 + \tfrac{1}{2V} \sum_{\sigma,\sigma'} \sum_{\alpha,\alpha'=\pm 1} \sum_{\mathbf{q}} f^{\sigma\sigma'}_{\alpha\alpha'} \, \rho_{\alpha\sigma}(\mathbf{q}) \rho_{\alpha'\sigma'}(-\mathbf{q}) \tag{6.28}$$

with H_0 as in (6.1) and a *Landau function*

$$f^{\sigma\sigma'}_{\alpha\alpha} = g_4^{*\sigma\sigma'} \quad \text{and} \quad f^{\sigma\sigma'}_{\alpha,-\alpha} = g_2^{*\sigma\sigma'} \tag{6.29}$$

This is the Hamiltonian of the *Luttinger model* (actually a simple generalization of the original Tomonaga-Luttinger model [21] obtained by including spin). As proposed by

[19] Backscattering is always marginal at tree (0-loop) level, but usually (marginally) irrelevant due to fluctuations (1-loop or higher) for systems with repulsive interactions.

Haldane [25], one-dimensional systems whose leading low-energy physics is governed by a Luttinger model fixed point are called *Luttinger liquids*. Note that the fixed point action $\bar{S}^*$ corresponding to H^* is identical to the fixed point action (4.15) of a Fermi liquid, adapted to the one-dimensional case with only two Fermi points $\mathbf{k}_F = \pm k_F$! Differences between Fermi and Luttinger liquid behavior arise only due to the enhanced phase space for forward scattering in one dimension.

With a momentum transfer cutoff $\Lambda_q \ll \Lambda$, the Luttinger model can be solved exactly and completely in the sense that all correlation functions can be obtained [20]. This can be achieved either by exploiting the Ward identities associated with the peculiar conservation laws of the system [61, 86, 116], or by bosonization [22, 25, 65]. The Luttinger model Hamiltonian H^* conserves charge and the z-component of spin separately on each Fermi point. The Ward identities obtained from the continuity equations associated with these conservation laws yield a complete system of equations for any correlation function.

The solution of the Luttinger model can also be obtained as a special case of the (more general) results for d-dimensional systems in Sec. 5. The properties derived there to leading order in a small-momentum-transfer expansion for systems dominated by forward scattering are exact to all orders in $\mathbf{q}$ in the Luttinger model. The reason for this is that here the velocity of the fermions is conserved *exactly* in each scattering process.[20] In particular, the loop cancellation derived in 5.1 and the density-current relations $\mathbf{\Lambda}(p;q) = \mathbf{v_p}\Lambda^0(p;q)$ obtained in 5.2 hold exactly in the Luttinger model.

As a consequence of loop cancellation, polarization insertions remain undressed and RPA results for the long-wavelength charge and spin response are exact [61, 20]. Let us discuss the results for the density-density response and the dynamical conductivity.

Continuing analytically from Matsubara frequencies q_0 to real frequencies ω, RPA (or the Ward identities) yield the well-known result for the charge density-density response [20], which can be expressed in terms of the velocities v_c and $\tilde{v}_c$ associated with global and axial charge conservation as [86]

$$N(\omega, \mathbf{q}) := J^{00}(\omega, \mathbf{q}) = \frac{2}{\pi} \frac{v_c \mathbf{q}^2}{\omega^2 - (u_c \mathbf{q})^2} \tag{6.30}$$

where

$$u_c = \sqrt{v_c \tilde{v}_c} \tag{6.31}$$

This response function has poles in $\omega = \pm u_c \mathbf{q}$, implying the existence of charge density modes (zero sound) with velocity u_c in the system. The existence of these modes is a

[20] Velocity conservation is equivalent to left/right conservation laws for charge and spin together with the linearity of the dispersion relation in the Luttinger model.

direct consequence of left/right charge conservation: Noting that $\mathbf{j} = v_c\tilde{\rho}$ and $\tilde{\mathbf{j}} = \tilde{v}_c\rho$, the continuity equations $i\partial_t\rho = \mathbf{q}\cdot\mathbf{j}$ and $i\partial_t\tilde{\rho} = \mathbf{q}\cdot\tilde{\mathbf{j}}$ can be combined to a harmonic oscillator equation for $\rho(t,\mathbf{q})$, namely $\partial_t^2\rho(t,\mathbf{q}) + v_c\tilde{v}_c\,\mathbf{q}^2\rho(t,\mathbf{q}) = 0$, describing undamped harmonic oscillations with frequency $u_c|\mathbf{q}|$ where $u_c = \sqrt{v_c\tilde{v}_c}$.

The *compressibility* $\kappa = \partial n/\partial\mu$ is obtained from (6.30) as

$$\kappa = \frac{2}{\pi\tilde{v}_c} \tag{6.32}$$

Using the relation $\sigma(q) = (i\omega/\mathbf{q}^2)\,N(q)$ for the dynamical *conductivity*, one finds

$$\sigma(\omega,\mathbf{q}) = \frac{2}{\pi}v_c\frac{i\omega}{\omega^2 - (u_c\mathbf{q})^2} \tag{6.33}$$

The absorptive part in a homogeneous field is therefore

$$Re\,\sigma(\omega) = 2v_c\delta(\omega) \tag{6.34}$$

i.e. a delta-peak with weight $2v_c$. Note that the velocities v_c and $\tilde{v}_c$ are in a one-to-one correspondence to directly observable physical quantities: v_c determines the weight of the Drude peak in the conductivity and $\tilde{v}_c$ the compressibility of the system.

The compressibility and the weight of the Drude peak are both finite, indicating that the g-ology model without (or with irrelevant) Umklapp terms describes a *stable metal*. By contrast, if Umklapp terms are relevant the compressibility is singular and the system becomes insulating. This latter behavior occurs in the one-dimensional Hubbard model with repulsive interactions at half-filling [24].

The spin density-density response is obtained as [20, 86]

$$S(\omega,\mathbf{q}) := J^{z0,z0}(\omega,\mathbf{q}) = \frac{1}{2\pi}\,\frac{v_s\mathbf{q}^2}{\omega^2 - (u_s\mathbf{q})^2} \tag{6.35}$$

with

$$u_s = \sqrt{v_s\tilde{v}_s} \tag{6.36}$$

where v_s and $\tilde{v}_s$ are the velocities associated with the conserved global and axial spin currents. The poles in $S(\omega,\mathbf{q})$ imply the existence of spin density modes with a velocity u_s. The spin *susceptibility* obtained from (6.35) is simply

$$\chi = \frac{2\mu_B^2}{\pi\tilde{v}_s} \tag{6.37}$$

where μ_B is the Bohr magneton.

Note that the above results for the charge response remain valid even in the presence of backscattering, while the results for the spin response are not affected by umklapp scattering.

The single-particle propagator can be obtained from the solution for G derived for d-dimensional forward scattering systems in Sec. 5. This solution is exact for the Luttinger model. Notice that the Ward identities (5.27) and (5.29) are indeed exact in the Luttinger model, and could be derived, without invoking loop cancellation, by directly using the results of Section 6.3.2 obtained from separate left and right conservation laws (which imply exact loop cancellation) [86]. Hence, G is determined by the integral equation (cf. (5.42))

$$(p_0 - v_F p_r)\, G(p_0, p_r) \;=\; 1 \;-\; \int_{p_0', p_r'} \frac{D(p_0 - p_0', p_r - p_r')}{i(p_0 - p_0') - v_F(p_r - p_r')}\, G(p_0', p_r') \tag{6.38}$$

where $D(q_0, q_r) = D^{\sigma\sigma}_{\alpha\alpha}(q_0, \mathbf{q})$ is the RPA effective interaction between particles with parallel spin near the same Fermi point, and $q_r = \mathbf{n}_\alpha \cdot \mathbf{q}$. In one dimension, Eq. (5.21) for D can be written as

$$D^{\sigma\sigma'}_{\alpha\alpha'}(q) = f^{\sigma\sigma'}_{\alpha\alpha'} + \sum_{\sigma''} \sum_{\alpha''} f^{\sigma\sigma''}_{\alpha\alpha''}\, \Pi_0^{\alpha''}(q)\, D^{\sigma''\sigma'}_{\alpha''\alpha'}(q) \tag{6.39}$$

where

$$\Pi_0^\alpha(q) = \frac{1}{2\pi}\, \frac{q_r}{iq_0 - v_F q_r} \tag{6.40}$$

The above equation for G has been first derived by Dzyaloshinskii and Larkin [61]. Equation (6.39) for the matrix $D^{\sigma\sigma'}_{\alpha\alpha'}(q)$ can be easily solved; the "diagonal" element can be written explicitly as

$$D(q_0, q_r) = (iq_0 - v_F q_r)\pi \sum_{\nu=c,s} \left[\frac{(2-\eta_\nu)(u_\nu - v_F)}{iq_0 - u_\nu q_r} + \frac{\eta_\nu (u_\nu + v_F)}{iq_0 + u_\nu q_r} \right] \tag{6.41}$$

where $u_\nu = (v_\nu \tilde{v}_\nu)^{1/2}$ is the velocity of the collective modes ($\nu = c, s$), and

$$\eta_\nu = (K_\nu + K_\nu^{-1} - 2)/4 \quad \text{where} \quad K_\nu = (v_\nu/\tilde{v}_\nu)^{1/2} \tag{6.42}$$

Inserting the analytic continuation of D to real frequencies into the expression (5.48) for L, one obtains

$$\begin{aligned} L(t, r) \;&=\; \log(r - v_F t + is(t)/\Lambda_q) \\ &- \sum_{\nu=c,s} \Big[(1/2 + \eta_\nu/2) \log(r - u_\nu t + is(t)/\Lambda_q) + (\eta_\nu/2) \log(r + u_\nu t + is(t)/\Lambda_q) \Big] \end{aligned} \tag{6.43}$$

and $L_0 = L(0, 0) = \eta \log \Lambda_q$, where $\eta = \eta_c + \eta_s$. For large r and/or large t one thus finds [20]

$$G(t, r) = \frac{1}{2\pi \Lambda_q^\eta} \prod_{\nu=c,s} \frac{1}{(r - u_\nu t + is(t)/\Lambda_q)^{1/2 + \eta_\nu/2}} \frac{1}{(r + u_\nu t - is(t)/\Lambda_q)^{\eta_\nu/2}} \tag{6.44}$$

Note that the exponents are uniquely determined by K_ν, i.e. by the ratios $v_\nu/\tilde{v}_\nu$. For non-interacting particles $K_c = K_s = 1$. The value $K_s = 1$ is maintained for spin-rotation invariant interactions, while $K_c < 1$ (> 1) for repulsive (attractive) forces. The Fourier-transform $G(\omega, k_r)$ of $G(t, r)$ is not an elementary function. The result (6.44) implies that the density of single-particle excitations vanishes as ω^η at low energy, and the momentum distribution function near k_F obeys a power law

$$n_{\mathbf{k}} - n_{\mathbf{k}_F} \;\propto\; -s(k_r)|k_r|^\eta \tag{6.45}$$

Spin-rotationally invariant microscopic models such as the Hubbard model lead generically to $\eta_s = 0$, $\eta_c > 0$ and $u_c \neq u_s$. In this case, for $\mathbf{k}$ outside the Fermi-surface and $u_c > u_s$, say, the $\mathbf{k}$-resolved spectral function $\rho(\omega, \mathbf{k})$ for single particle excitations, has power-law divergences for $\omega \to u_c k_r$ and for $\omega \to u_s k_r$ with $\omega > u_s k_r$, vanishes for $-u_c k_r < \omega < u_s k_r$, and has finite values also for $\omega < -u_c k_r$ [123, 124]. The two peaks in the spectral function at $\omega = u_c k_r$ and $\omega = u_s k_r$ indicate that the extra charge and spin associated with an additional electron inserted into the system propagate with different velocities, a phenomenon called "spin-charge-separation". This is in striking contrast to the behavior of a Fermi liquid, where the spectral function has the form $\rho(\omega, \mathbf{k}) \propto \delta(\omega - \xi_{\mathbf{k}})$ in the low-energy limit. Landau quasi-particle excitations are absent in the Luttinger liquid.

The velocity ratios $v_\nu/\tilde{v}_\nu$ (i.e. K_ν) determine also the anomalous scaling dimensions of $2k_F$-density correlations and Cooper pair correlation functions in a Luttinger liquid [23, 24].

The (bosonic) charge and spin density modes corresponding to the poles in the charge and spin response functions are the only low-energy excitations in a Luttinger liquid. They determine the leading low-temperature contribution to the specific heat

$$c_V = \frac{\pi}{6}(u_c^{-1} + u_s^{-1})\, T \tag{6.46}$$

A computation of the free energy associated with $\bar{S}^*$ (given by RPA due to loop cancellation) shows that there are no other contributions.

In the Luttinger model, the charge and spin density modes are exact eigenstates (i.e. undamped), and any excited state of the model is a superposition of these elementary excitations. This important fact becomes particularly explicit in the bosonized form of the Luttinger model obtained first by Mattis and Lieb [22]. The bosonization procedure described in Sec. 5 is exact for the Luttinger model. The expression (5.63) for the Hamiltonian as a quadratic form in density fluctuation operators turns into

$$H^* = \tfrac{1}{2V} \sum_{\sigma,\sigma'} \sum_{\alpha,\alpha'} \sum_{\mathbf{q}} \left[2\pi \mathbf{v}_\alpha \cdot \mathbf{q}\, \delta_{\sigma\sigma'} \delta_{\alpha\alpha'} + f^{\sigma\sigma'}_{\alpha\alpha'} \right] \rho_{\alpha\sigma}(\mathbf{q}) \rho_{\alpha'\sigma'}(-\mathbf{q}) \tag{6.47}$$

for fermions with spin in one dimension. Transforming to canonical boson annihilation and creation operators $b_{\alpha\sigma}(\mathbf{q})$ and $b^\dagger_{\alpha\sigma}(\mathbf{q})$ with $\mathbf{n}_\alpha\cdot\mathbf{q} > 0$ via (cf. (5.56))

$$\rho_{\alpha\sigma}(\mathbf{q}) = \sqrt{V/2\pi}\left[\Theta(\mathbf{n}_\alpha\cdot\mathbf{q})b_{\alpha\sigma}(\mathbf{q}) + \Theta(-\mathbf{n}_\alpha\cdot\mathbf{q})b^\dagger_{\alpha\sigma}(-\mathbf{q})\right] \tag{6.48}$$

one obtains a quadratic form in b and $b^\dagger$. Off-diagonal terms are easily eliminated by a linear (Bogoliubov) transformation from b and $b^\dagger$ to new boson operators β and $\beta^\dagger$. In each subspace with fixed particle number and total momentum, the Luttinger model Hamiltonian can thus be transformed into [22, 25, 20]

$$H^* = \sum_{\nu=c,s}\sum_{\mathbf{q}} u_\nu|\mathbf{q}|\beta^\dagger_\nu(\mathbf{q})\beta_\nu(\mathbf{q}) \tag{6.49}$$

Any excited state is thus obviously a superposition of non-interacting elementary bosonic excitations corresponding to charge or spin density modes in the underlying interacting Fermi system. In particular, one has "spin-charge separation" in the sense that charge and spin excitations are independent from each other.

Let us briefly compare the above results to those of Fermi liquid theory. Concerning the leading low-energy long-wavelength response functions there is no difference between Fermi and Luttinger liquids. In both liquids the response is governed by the same fixed point action $\bar{S}^*$ and the residual forward scattering can be treated in RPA. However, in a Luttinger liquid RPA works due to more subtle reasons than in a (two- or three-dimensional) Fermi liquid. In the latter corrections to RPA are suppressed simply by the reduced phase space for residual scattering processes, while in the former RPA emerges due to cancellations (of self-energy and vertex corrections) imposed by the (asymptotic) conservation laws!

Marked differences between Fermi and Luttinger liquids appear in the single-particle propagator G, which determines the momentum distribution function and the spectral density for single-particle excitations. In a Fermi liquid residual interactions modify the propagator only on a subleading level (leading to a small quasi-particle decay etc.), while in a Luttinger liquid forward scattering affects the leading low energy behavior. Another distinctive feature of Luttinger liquids is the singular behavior of density correlations with momenta near $2k_F$ [20, 23].

The similarity of the fixed point Hamiltonians in Fermi and Luttinger liquids (cf. (4.15) and (6.28)) as well as the common RPA structure of response functions suggests the identification of $f^{\sigma\sigma'}_{\alpha\alpha} = g_4^{\sigma\sigma'}$ and $f^{\sigma\sigma'}_{\alpha,-\alpha} = g_2^{\sigma\sigma'}$ as a Landau function in one-dimensional systems. One should, however, be aware of the following ambiguity: In a Luttinger liquid, the Landau function is not uniquely determined from the low-energy behavior of the correlation functions, in constrast to the situation in a Fermi liquid, where (4.7) yields

a unique identification. In fact, all correlation functions in the one-dimensional Luttinger model are uniquely parametrized by the four velocities v_ν, $\tilde{v}_\nu$ ($\nu = c, s$) in the low-energy limit, where, in terms of v_F and g's (see Sec. 6.3)

$$v_\nu = v_F + (g_4^\nu - g_2^\nu)/\pi \quad , \quad \tilde{v}_\nu = v_F + (g_4^\nu + g_2^\nu)/\pi \tag{6.50}$$

An equivalent parametrization can be given in terms of K_ν and the sound velocities u_ν. Obviously all the velocities v_ν, and hence all correlation functions are invariant under the shift

$$v_F \mapsto v_F + \delta v_F \quad , \quad g_4^{\sigma\sigma} \mapsto g_4^{\sigma\sigma} - \pi \delta v_F \tag{6.51}$$

Auxiliary functions such as the effective interaction $D(\omega, q_r)$ depend on the choice of v_F. The ambiguity in the identification of a Landau function in $d = 1$ can be removed by imposing the Landau sum-rule for the scattering amplitudes [2, 94], which is equivalent to the condition $D(0, q_r) = 0$ in eq. (6.41). In general, one finds

$$D(0, q_r) = 2\pi v_F - \frac{\pi v_F^2}{2} \sum_\nu \left(v_\nu^{-1} + \tilde{v}_\nu^{-1} \right) \tag{6.52}$$

Hence, only for the special and unique choice

$$v_F = \left[\tfrac{1}{4} \sum_\nu \left(v_\nu^{-1} + \tilde{v}_\nu^{-1} \right) \right]^{-1} \tag{6.53}$$

D satisfies the condition $D(0, q_r) = 0$ associated with the Landau sum-rule. These remarks will play a role in the analysis of the crossover from Luttinger to Fermi liquid behavior as a function of dimensionality in Sec. 7.

6.5. INSTABILITIES

Instabilities of the Luttinger liquid closely parallel those of a Fermi liquid. Stability with respect to deformations of the Fermi "surface" requires

$$(g_4^\nu \pm g_2^\nu)/\pi v_F \; > \; -1 \tag{6.54}$$

or (equivalently) $\tilde{v}_\nu > 0$ and $v_\nu > 0$. Violation of one of these conditions leads to a one-dimensional analogue of the *Pomerantchuk instabilities* known for Fermi liquids (cf. (4.29)). For example, for $(g_4^c + g_2^c)/\pi v_F < -1$ the compressibility becomes negative and the liquid will undergo phase-separation.

Depending on the signs of the coupling constants, backscattering processes may or may not generate an instability of the Luttinger liquid towards a state with a spin-gap and enhanced charge-density wave or Cooper-pair correlations (both may be enhanced; the sign of $K_c - 1$ determines which correlation is stronger) [20]. This behavior may signal a

Spin-Peierls instability or a one-dimensional analogue of the *Cooper instability*, although genuine off-diagonal long-range order is prevented here by the strong order-parameter fluctuations in one dimension. Similarly, umklapp scattering may lead to the formation of a charge gap. This latter instability is an analogue of *density-wave instabilities* (or Mott transition) in higher dimensions.

7. SHORT-RANGE INTERACTIONS IN D DIMENSIONS

This section is devoted to a quantitative analysis of residual low-energy scattering processes in normal d-dimensional Fermi systems with *short-range* interactions. Our starting point is an effective low-energy action of the form

$$\bar{S}^{\Lambda} = \sum_{\sigma} \int_{k} \psi^{*}_{k\sigma}(ik_0 - \xi_{\mathbf{k}})\psi_{k\sigma} \; - \; \tfrac{1}{2} \sum_{\sigma\sigma'} \int_{k,k',q} g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(\mathbf{q}) \, \psi^{*}_{k-q/2,\sigma}\psi_{k+q/2,\sigma}\psi^{*}_{k'+q/2,\sigma'}\psi_{k'-q/2,\sigma'} \quad (7.1)$$

where $\xi_{\mathbf{k}} = v_{\mathbf{k}_F} k_r$ is a linearized dispersion relation and $g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(\mathbf{q})$ a renormalized coupling function. In a system with short-range interactions, g is assumed to be a *regular* function of $\mathbf{k}$, $\mathbf{k}'$ and $\mathbf{q}$. To keep the notation readable, we have suppressed bars indicating renormalization and also the cutoff-dependence on the right hand side of (7.1). Of course we have to keep in mind that a field renormalization Z^{Λ} has been performed in passing from some microscopic model to the effective low-energy action. We assume that Λ is much smaller than k_F, and will analyze the effect of the residual two-particle scattering processes. Explicit results will be derived mainly for rotationally invariant continuum systems (without umklapp processes) with a spherical Fermi surface and a constant Fermi velocity $v_{\mathbf{k}_F} = v_F$.

Let us assume that the residual interactions in the Cooper channel are purely repulsive such that we need not bother about the Cooper instability (or backscattering instability in $d = 1$). We will refer to $\bar{S}^{\Lambda}$ in (7.1) with finite (non-singular) coupling functions and vanishing or repulsive Cooper couplings as the *regular normal model*. This model describes the low-energy physics of all those Fermi systems with short-range interactions where high-energy degrees of freedom do not generate an instability (such as symmetry breaking, pair-formation, etc.).

In two or three dimensions the regular normal model leads to Fermi liquid behavior with fermionic quasi-particle excitations (at least to each finite order in perturbation theory), while in one dimension Luttinger liquid behavior is found. In any dimension the leading low-energy long-wavelength response is given exactly by the random phase approximation in the limit $\Lambda \to 0$. The momentum distribution function and the structure of single-particle excitations are however dramatically different in $d = 1$, compared to $d = 2$ or $d = 3$.

In this situation it is interesting to analyze the crossover from one-dimensional Luttinger liquid behavior to Fermi liquid behavior in higher dimensions as a function of

continuous dimensionality.[21] In particular, one would like to know which *critical dimension* d_c separates Fermi from Luttinger liquid behavior, and whether new non-Fermi liquid fixed points can be found within a suitable ϵ-expansion. It turns out that there are in fact *two* characteristic dimensions: In any dimension d above the critical dimension $d_c = 1$ the *leading* low-energy behavior is of Fermi liquid type, but for $d \leq d_c' = 2$ the *subleading* corrections differ from the simple behavior known for three-dimensional Fermi systems. In systems with short-range interactions there are no non-Fermi liquid fixed points in $d = 1 + \epsilon$ dimensions.

We will reach a detailed understanding of the influence of residual interactions on the low-energy physics of d-dimensional systems in three steps: i) perturbation theory, ii) RPA, and iii) a resummation of forward scattering along the lines of Sec. 5. Forward scattering is favored in dimensions $1 \leq d < 2$ by an angle-dependent phase-space factor (see the angle-integration in eq. (7.2)). As a byproduct we obtain some technical material such as analytic results for the d-dimensional particle-hole bubble, that can be useful in future investigations of Fermi systems in continuous dimensionality (especially ϵ-expansions).

A short account of the main results of this section has been published in Ref. [35]. In the following we work exclusively with real frequencies at zero temperature.

7.1. CONTINUATION TO NON-INTEGER DIMENSIONS

The continuation of the theory to non-integer dimensions is obtained as usual by analytic continuation of Feynman diagrams, defined for general integer d, in the complex d-plane. For our purposes it will be sufficient to continue momentum integrals of functions $f(\mathbf{k})$ which depend on $\mathbf{k}$ only via $|\mathbf{k}|$ and an angle θ between $\mathbf{k}$ and another momentum which is fixed. In these cases one can use

$$\int d^d\mathbf{k} \ldots \;=\; S_{d-1} \int_0^\infty d|\mathbf{k}|\, |\mathbf{k}|^{d-1} \int_0^\pi d\theta\, (\sin\theta)^{d-2} \ldots \tag{7.2}$$

where

$$S_d = 2\pi^{d/2}/\Gamma(d/2) \tag{7.3}$$

is the surface of the d-dimensional unit sphere. In the limit $d \to 1$ one has $S_{d-1} \sim d - 1$, and thus

$$S_{d-1}(\sin\theta)^{d-2} \;\to\; \delta(\theta) + \delta(\theta - \pi) \quad \text{for} \quad d \to 1 \tag{7.4}$$

as expected. For fermionic momenta (not momentum transfers) near the Fermi surface it is convenient to decompose

$$|\mathbf{k}| = k_F + k_r \tag{7.5}$$

[21] Analytic continuation of dimensionality is a common technique in statistical mechanics and field-theory, especially as a means to treat strong coupling problems by performing an ϵ-expansion around a critical dimension of the system [118, 119].

where k_r (the oriented distance from the Fermi surface) is much smaller than k_F. Approximating $|\mathbf{k}|^{d-1} \sim k_F^{d-1}$, (7.2) may then be simplified to

$$\int d^d\mathbf{k} \ldots = S_{d-1} k_F^{d-1} \int dk_r \int_0^\pi d\theta \, (\sin\theta)^{d-2} \ldots \tag{7.6}$$

where the k_r-integration is restricted either by a cutoff Λ or by small external momentum or energy variables. Corrections are suppressed by a factor of order $(d-1)k_r/k_F$ in the integral (7.6).

7.2. PERTURBATIVE RESULTS

We will first present explicit results for the particle-hole and particle-particle bubble in d dimensions, and then for the second order self-energy.

7.2.1. Particle-hole bubble

The *particle-hole bubble* (or bare polarization insertion)

$$\Pi_0(q) = -i \int_k G_0(k) G_0(k+q) \tag{7.7}$$

is essentially the dynamical density-density correlation function of the non-interacting system, and will be important for subsequent explicit calculations of the self-energy Σ and the effective interaction D. At zero temperature, the particle-hole bubble can be written as

$$\Pi_0(\omega, \mathbf{q}) = \int \frac{d^d\mathbf{k}}{(2\pi)^d} \frac{\Theta(\xi_{\mathbf{k}+\mathbf{q}}) - \Theta(\xi_{\mathbf{k}})}{\omega - (\xi_{\mathbf{k}+\mathbf{q}} - \xi_{\mathbf{k}}) + i0^+ s(\omega)} \tag{7.8}$$

where $\xi_{\mathbf{k}} = \epsilon_{\mathbf{k}} - \mu$. The spectral density of particle-hole excitations is given by

$$\Delta_0^{ph}(\omega, \mathbf{q}) = -\frac{1}{\pi} s(\omega) \, \mathrm{Im}\Pi_0(\omega, \mathbf{q}) \tag{7.9}$$

This is a positive quantity for $\omega > 0$ and negative for $\omega < 0$ (as usual for spectral functions of bosons). For positive energies, Δ_0^{ph} can be written as

$$\Delta_0^{ph}(\omega, \mathbf{q}) = \int \frac{d^d\mathbf{k}}{(2\pi)^d} \, \Xi(\xi_{\mathbf{k}} < 0 < \xi_{\mathbf{k}+\mathbf{q}}) \, \delta[\omega - (\xi_{\mathbf{k}+\mathbf{q}} - \xi_{\mathbf{k}})] \tag{7.10}$$

while $\Delta_0^{ph}(-\omega, \mathbf{q}) = -\Delta_0^{ph}(\omega, \mathbf{q})$. Here $\Xi(\mathit{condition}) = 1$ if the condition is satisfied, and $\Xi(\mathit{condition}) = 0$ otherwise.

For a linear dispersion, $\xi_{\mathbf{k}} = v_F k_r$, several analytic results can be obtained for the low energy behavior (small ω) of $\Pi_0(\omega, \mathbf{q})$, in any dimension d. Using

$$\xi_{\mathbf{k}+\mathbf{q}} = v_F(|\mathbf{k}+\mathbf{q}| - k_F) = v_F \left[(|\mathbf{k}|^2 + |\mathbf{q}|^2 + 2|\mathbf{k}||\mathbf{q}|\cos\theta)^{1/2} - k_F\right] \tag{7.11}$$

where θ is the angle spanned by $\mathbf{k}$ and $\mathbf{q}$, and introducing spherical coodinates, $\Pi_0(\omega, \mathbf{q})$ can be written as a two-fold integral over the variables $|\mathbf{k}|$ and θ.

For the *imaginary* part Δ_0^{ph} the integration is simplified by the presence of the δ-function. For small ω (compared to $v_F k_F$) but general $\mathbf{q}$ one obtains

$$\Delta_0^{ph}(\omega, \mathbf{q}) \sim \frac{S_{d-1}}{(2\pi)^d} \frac{2}{d-1} \frac{k_F^d}{v_F} \frac{1}{|\mathbf{q}|} (1 + |\mathbf{q}|/2k_F)^{\frac{d-3}{2}} [1 - (\omega/v_F|\mathbf{q}|)^2]^{\frac{d-3}{2}}$$

$$\Big[(1 - |\mathbf{q}|/2k_F + \omega/2v_F k_F)^{\frac{d-1}{2}} - (1 - |\mathbf{q}|/2k_F - \omega/2v_F k_F)^{\frac{d-1}{2}} \Xi(\omega/v_F < 2k_F - |\mathbf{q}|)\Big] \tag{7.12}$$

for $\max(0, |\mathbf{q}| - 2k_F) < \omega/v_F < |\mathbf{q}|$, while $\Delta_0^{ph}(\omega, \mathbf{q}) = 0$ for other $\omega > 0$, and $\Delta_0^{ph}(-\omega, \mathbf{q}) = -\Delta_0^{ph}(\omega, \mathbf{q})$. For $|\mathbf{q}| < 2k_F$ and not close to $2k_F$, this simplifies to

$$\Delta_0^{ph}(\omega, \mathbf{q}) \sim \frac{S_{d-1}}{(2\pi)^d} \frac{k_F^{d-1}}{v_F^2} \frac{\omega}{|\mathbf{q}|} [1 - \omega^2/(v_F|\mathbf{q}|)^2]^{\frac{d-3}{2}} [1 - (|\mathbf{q}|/2k_F)^2]^{\frac{d-3}{2}} \Xi(|\omega| < v_F|\mathbf{q}|) \tag{7.13}$$

In general (non-integer) dimensions, the *real* part of $\Pi_0(\omega, \mathbf{q})$ cannot be expressed in terms of elementary functions, even if ω is small. Hence we will only present some useful results obtained for various limits. The behavior of $\Pi_0(\omega, \mathbf{q})$ for small ω and $\mathbf{q}$ is particularly important. Since for $|\mathbf{q}| \sim 0$ the excitation energies $\xi_{\mathbf{k}+\mathbf{q}} - \xi_{\mathbf{k}} \sim |\mathbf{q}| \cos\theta$ depend only on $|\mathbf{q}|$ and θ, the integration over $|\mathbf{k}|$ is easily carried out. For $\cos\theta > 0$ one has contributions from $k_r \in [-|\mathbf{q}| \cos\theta, 0]$, while for $\cos\theta < 0$ values $k_r \in [0, -|\mathbf{q}| \cos\theta]$ contribute, yielding

$$\Pi_0(\omega, \mathbf{q}) \sim \frac{S_{d-1}}{(2\pi)^d} k_F^{d-1} \int_0^\pi d\theta \frac{|\mathbf{q}| \cos\theta}{\omega - v_F|\mathbf{q}| \cos\theta + i0^+ s(\omega)} (\sin\theta)^{d-2} \tag{7.14}$$

This depends on ω and $\mathbf{q}$ only via the ratio $\omega/|\mathbf{q}|$. Results from a numerical evaluation of (7.14) in various dimensions are shown in Fig. 7.1. For $|\mathbf{q}|/\omega \to 0$, $\Pi_0(\omega, \mathbf{q})$ vanishes, while in the opposite limit $\omega/|\mathbf{q}| \to 0$ it assumes the negative real value

$$\Pi_0(0, \mathbf{q}) \sim -\frac{S_d}{(2\pi)^d} \frac{k_F^{d-1}}{v_F} = -N_F \tag{7.15}$$

where N_F is the density of states per spin at the Fermi level. For a linear dispersion, $\Pi_0(\omega, \mathbf{q})$ diverges in the limit $\omega/v_F|\mathbf{q}| \to 1$ in any dimension $d \leq 3$. In $d < 3$, the leading singularity is given by

$$\Pi_0(\omega, \mathbf{q}) \sim \frac{S_{d-1}}{(2\pi)^d} 2^{\frac{d-3}{2}} (B_d^\pm + i\pi) k_F^{d-1} v_F^{-1} |\omega/v_F|\mathbf{q}| - 1|^{\frac{d-3}{2}} \quad \text{for} \quad \frac{\omega}{v_F|\mathbf{q}|} \to 1 + 0^\pm \tag{7.16}$$

where

$$B_d^+ = B[(d-1)/2), (3-d)/2]\,, \quad B_d^- = B_d^+ \cos[(3-d)/2] \tag{7.17}$$

where $B(x,y) = \Gamma(x)\Gamma(y)/\Gamma(x+y)$ is Euler's Beta-function (a special function, not to be confused with the β-function from the renormalization group). Note that B_d^- is positive in $d > 2$ and negative in $d < 2$. An exception is the real part of $\Pi_0(\omega, \mathbf{q})$ for $\omega < v_F|\mathbf{q}|$ in $d = 2$, since $B_2^- = 0$; in this case one finds

$$\mathrm{Re}\Pi_0(\omega, \mathbf{q}) \sim -k_F/2\pi v_F \quad \text{for all} \quad \omega < v_F|\mathbf{q}| \quad \text{in } d = 2 \tag{7.18}$$

Note that the cutoff Λ does not appear in the above results, since the $\mathbf{k}$-integral is already limited by other small variables such as ω/v_F or $|\mathbf{q}|$.

In Appendix C we list results for the particle-hole bubble in d dimensions obtained with a quadratic (not linearized) dispersion relation $\epsilon_{\mathbf{k}} = \mathbf{k}^2/2m$.

7.2.2. Particle-particle bubble

The particle-particle bubble is defined by

$$K_0(p) = i \int_k G_0(k) G_0(p-k) \tag{7.19}$$

This quantity plays a role in the T-matrix approximation for systems with local interactions [30, 31]. It also enters into a 1-loop calculation of the β-function for Cooper couplings.

At zero temperature, the particle-particle bubble can be written as

$$K_0(\omega, \mathbf{p}) = \int \frac{d^d\mathbf{k}}{(2\pi)^d} \frac{\Theta(\xi_{\mathbf{p}-\mathbf{k}}) - \Theta(-\xi_{\mathbf{k}})}{\omega - (\xi_{\mathbf{p}-\mathbf{k}} + \xi_{\mathbf{k}}) + i0^+ s(\omega)} \tag{7.20}$$

The spectral density for two-particle (two-hole) excitations is given by

$$\Delta_0^{pp}(\omega, \mathbf{p}) = -\frac{1}{\pi} s(\omega) \mathrm{Im} K_0(\omega, \mathbf{p}) \tag{7.21}$$

This quantity is positive for $\omega > 0$ (two particles) and negative for $\omega < 0$ (two holes). Another way of writing Δ_0^{pp} is

$$\Delta_0^{pp}(\omega, \mathbf{p}) = \int \frac{d^d\mathbf{k}}{(2\pi)^d} [\Theta(\xi_{\mathbf{k}})\Theta(\xi_{\mathbf{p}-\mathbf{k}}) - \Theta(-\xi_{\mathbf{k}})\Theta(-\xi_{\mathbf{p}-\mathbf{k}})] \delta(\omega - \xi_{\mathbf{k}} - \xi_{\mathbf{p}-\mathbf{k}}) \tag{7.22}$$

We now present analytic results for $\Delta_0^{pp}(\omega, \mathbf{p})$ for small ω and a linear dispersion relation $\xi_{\mathbf{k}} = v_F k_r$. For $\omega < v_F(|\mathbf{p}| - 2k_F)$, $\Delta_0^{pp}(\omega, \mathbf{p})$ vanishes. For $\omega > v_F(|\mathbf{p}| - 2k_F)$, one finds

$$\Delta_0^{pp}(\omega, \mathbf{p}) \sim s(\omega) \frac{S_d}{2(2\pi)^d} \frac{k_F^{d-1}}{v_F} \tag{7.23a}$$

if $|\omega| > |\mathbf{p}|$, while

$$\Delta_0^{pp}(\omega, \mathbf{p}) \sim \frac{S_{d-1}}{(2\pi)^d} \frac{k_F^{d-1}}{v_F} \left[(1 + \omega/2v_F k_F)^2 - (|\mathbf{p}|/2k_F)^2\right]^{\frac{d-3}{2}} \int_0^{\omega/v_F|\mathbf{p}|} dx\, (1-x^2)^{(d-3)/2} \tag{7.23b}$$

if $|\omega| < |\mathbf{p}|$. The latter result simplifies in particular cases, namely to

$$\Delta_0^{pp}(\omega, \mathbf{p}) \sim \frac{S_{d-1}}{(2\pi)^d} \frac{k_F^{d-1}}{v_F} \int_0^{\omega/v_F|\mathbf{p}|} (1-x^2)^{(d-3)/2} \tag{7.24a}$$

for small $|\mathbf{p}|$,

$$\Delta_0^{pp}(\omega, \mathbf{p}) \sim \frac{S_{d-1}}{(2\pi)^d} \frac{k_F^{d-1}}{v_F} \left[(1+\omega/2v_F k_F)^2 - (|\mathbf{p}|/2k_F)^2\right]^{\frac{d-3}{2}} \frac{\omega}{|\mathbf{p}|} \tag{7.24b}$$

for $|\mathbf{p}| \sim 2k_F$, and to

$$\Delta_0^{pp}(\omega, \mathbf{p}) \sim \frac{S_{d-1}}{(2\pi)^d} \frac{k_F^{d-1}}{v_F} [(1-(|\mathbf{p}|/2k_F)^2]^{\frac{d-3}{2}} \frac{\omega}{|\mathbf{p}|} \tag{7.24c}$$

if $|\mathbf{p}|$ is neither close to 0 nor close to $2k_F$. We emphasize that these results hold only for ω small compared to $v_F k_F$.

Note that (7.23a) implies that $|\Delta_0^{pp}(\omega, \mathbf{0})|$ is a constant at small ω in any dimension. Hence (by Kramers-Kronig relations), the real part of the particle-particle bubble $\mathrm{Re}K_0(\omega, \mathbf{0})$ is logarithmically divergent for $\omega \to 0$:

$$\mathrm{Re}K_0(\omega, \mathbf{0}) \sim -\frac{S_d}{(2\pi)^d} \frac{k_F^{d-1}}{v_F} \log(\omega/v_F\Lambda) \tag{7.25}$$

where Λ is a cutoff. This leads to a finite contribution to the β-function for Cooper couplings, as long as (renormalized) Cooper couplings are non-zero (see Sec. 2).

In Appendix C we list results for the particle-particle bubble in d dimensions obtained with a quadratic (not linearized) dispersion relation $\epsilon_{\mathbf{k}} = \mathbf{k}^2/2m$.

7.2.3. Second order self-energy

The *first order* self-energy does not yield any many-body effects, but merely shifts the chemical potential μ, and possibly the Fermi velocity v_F. Hence we concentrate immediately on the *second order* self-energy, which will be calculated for a *constant coupling* g acting between particles with *opposite spins*. The restriction to a momentum-independent coupling corresponds to a local interaction in real space, and is made to provide us with explicit results for one concrete case.[22] This case is however a generic representative for short-range interactions, yielding all the typical dynamical many-body effects described by a self-energy in a normal Fermi system: quasi-particle decay, reduction of quasi-particle weight, and complete destruction of the quasi-particle in one dimension.

[22] Obviously, a constant coupling between parallel spins would yield no contribution, as a consequence of the Pauli principle.

In a system with interactions between opposite spins (only), there is only one contribution to the second order self-energy, represented diagrammatically in Fig. 7.2. Algebraically, this corresponds to

$$\Sigma(p) = g^2 \int_{k,q} G_0(p-q) G_0(k) G_0(k+q) \tag{7.26}$$

We concentrate on the imaginary part, $\mathrm{Im}\Sigma(p)$, from which the real part can be constructed via the Kramers-Kronig (or spectral) representation

$$\Sigma(\xi, \mathbf{p}) = -\pi^{-1} \int_{-\infty}^{\infty} d\xi' \, \frac{s(\xi') \mathrm{Im}\Sigma(\xi', \mathbf{p})}{\xi - \xi' + i0^+ s(\xi)} \tag{7.27}$$

In general the imaginary part determines the real part only up to an energy-independent constant, which however in our case must vanish, since $\Sigma(p)$ in (7.26) obviously vanishes for $\xi \to \infty$. For constant couplings, $\mathrm{Im}\Sigma$ can be conveniently expressed in terms of the spectral density of particle-hole excitations as

$$\mathrm{Im}\Sigma(\xi, \mathbf{p}) = \begin{cases} -\pi g^2 \int \frac{d^d\mathbf{k}}{(2\pi)^d} \, \Delta_0^{ph}(\xi - \xi_{\mathbf{k}}, \mathbf{p} - \mathbf{k}) \, \Xi(0 < \xi_{\mathbf{k}} < \xi) & \text{for} \quad \xi > 0 \\ -\pi g^2 \int \frac{d^d\mathbf{k}}{(2\pi)^d} \, \Delta_0^{ph}(\xi - \xi_{\mathbf{k}}, \mathbf{p} - \mathbf{k}) \, \Xi(\xi < \xi_{\mathbf{k}} < 0) & \text{for} \quad \xi < 0 \end{cases} \tag{7.28}$$

or, alternatively, in terms of the spectral density of two-particle (or two-hole) excitations

$$\mathrm{Im}\Sigma(\xi, \mathbf{p}) = \begin{cases} -\pi g^2 \int \frac{d^d\mathbf{k}}{(2\pi)^d} \, \Delta_0^{pp}(\xi + \xi_{\mathbf{k}}, \mathbf{p} + \mathbf{k}) \, \Xi(\xi + \xi_{\mathbf{k}} > 0 > \xi_{\mathbf{k}}) & \text{for} \quad \xi > 0 \\ -\pi g^2 \int \frac{d^d\mathbf{k}}{(2\pi)^d} \, \Delta_0^{pp}(\xi + \xi_{\mathbf{k}}, \mathbf{p} + \mathbf{k}) \, \Xi(\xi + \xi_{\mathbf{k}} < 0 < \xi_{\mathbf{k}}) & \text{for} \quad \xi < 0 \end{cases} \tag{7.29}$$

A general analytic evaluation of these integrals is difficult. Hence, we have calculated $\mathrm{Im}\Sigma$ numerically, and confirmed only the most striking properties by analytic derivations. Typical results for $\mathrm{Im}\Sigma(\xi, \mathbf{p})$ are shown in Fig. 7.3. In three dimensions we observe the well-known quadratic energy dependence, $\mathrm{Im}\Sigma(\xi, \mathbf{p}) \propto \xi^2$, without any special feature at $\xi = \xi_{\mathbf{p}} = v_F p_r$. For $1 < d \leq 2$ contributions of order ξ^2 are superposed by larger $\xi_{\mathbf{p}}$-dependent terms. In $d < 2$, for small $p_r = |\mathbf{p}| - k_F$ and ξ, the self energy scales as

$$\mathrm{Im}\Sigma(\xi, \mathbf{p}) = |p_r|^d \, \mathrm{Im}\tilde{\Sigma}(\xi / p_r) \tag{7.30}$$

and diverges in $\xi = \xi_{\mathbf{p}}$ as [35]

$$\mathrm{Im}\Sigma(\xi, \mathbf{p}) \;\sim\; -C_d \, g^2 k_F^{d-1} v_F^{-(d+1)} s(\xi_{\mathbf{p}}) \, \xi^2 |\xi - \xi_{\mathbf{p}}|^{d-2} \tag{7.31}$$

where $C_d = 2^{-d-4} \pi^{1-2d} S_{d-1}^2 B[2-d, (d-1)/2]$ is a constant depending only on dimensionality. In $d = 2$ there is a weak logarithmic divergence for $\xi \to \xi_{\mathbf{p}}$. This singularity is exclusively due to *forward scattering* (i.e. small momentum transfers $\mathbf{q} \sim 0$) of particles with *opposite spin* and almost *parallel momenta*. Hence, for more general spin- and

momentum-dependent couplings $g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(\mathbf{q})$ one would obtain the same singularity, with a coupling $g^{\sigma,-\sigma}_{\mathbf{k}\mathbf{k}}(\mathbf{0}) = g_F^{\sigma,-\sigma}(\theta = 0)$ replacing g. Forward scattering of particles with *antiparallel momenta* yields a contribution proportional to $g^2|\xi - \xi_{\mathbf{p}}|^d\,\theta(|\xi| - |\xi_{\mathbf{p}}|)$ in $d < 2$, with g to be replaced by $g^{\sigma\sigma'}_{\mathbf{k},-\mathbf{k}}(\mathbf{0}) = g_F^{\sigma\sigma'}(\theta = \pi)$ for more general couplings.[23]

In $d < 2$, the single-particle propagator G is drastically affected by the singular contributions of forward scattering to the second order self-energy.

Let us consider the one-dimensional case first. In $d = 1$, forward scattering between particles with opposite momenta (H_2, in g-ology notation) yields a contribution proportional to $g^2(\xi - \xi_{\mathbf{p}})\theta(|\xi| - |\xi_{\mathbf{p}}|)$ to $\mathrm{Im}\Sigma$, which, via Kramers-Kronig, yields a real part proportional to $g^2\xi\log|\xi|$ for $\mathbf{p}$ on the Fermi surface. This leads to a wave function renormalization $Z = (1 - \partial\Sigma/\partial\xi)^{-1} \propto g^2/\log|\xi| \to 0$ as $\xi \to \xi_{\mathbf{p}} = 0$, which is a well-known perturbative signal for the breakdown of Fermi liquid theory in one dimension [20], i.e. a power-law behavior of the wave function renormalization with a non-universal exponent η. Forward scattering between particles with parallel momenta (H_4, in g-ology notation) does not contribute to the wave function renormalization, but nevertheless destroys the quasiparticle pole in the propagator, leading to *spin-charge separation* [23, 24]. In contrast to common wisdom this latter effect also has a clear perturbative signal [35]: In $d \to 1$, (7.31) reduces to

$$\mathrm{Im}\Sigma(\xi, \mathbf{p}) \;\sim\; -\frac{g^2}{8\pi v_F^2} s(\xi_{\mathbf{p}})\,\xi^2\delta(\xi - \xi_{\mathbf{p}}) \tag{7.32}$$

yielding, by Kramers-Kronig, a real part

$$\mathrm{Re}\Sigma(\xi, \mathbf{p}) \;\sim\; \frac{g^2}{8\pi^2 v_F^2}\frac{\xi_{\mathbf{p}}^2}{\xi - \xi_{\mathbf{p}}} \tag{7.33}$$

Inserting this into $G = (\xi - \xi_{\mathbf{p}} - \Sigma)^{-1}$ one obtains a propagator which has two poles instead of one, i.e. the spectral function becomes a sum of two δ-functions with weight $1/2$ each:

$$\rho(\xi, \mathbf{p}) = \frac{1}{2}\delta(\xi - \xi_{\mathbf{p}}^+) + \frac{1}{2}\delta(\xi - \xi_{\mathbf{p}}^-) \tag{7.34}$$

where $\xi_{\mathbf{p}}^- < \xi_{\mathbf{p}} < \xi_{\mathbf{p}}^+$.

In $1 < d < 2$ the perturbatively calculated wave function renormalization is finite, but the forward scattering processes of particles with almost parallel momenta still have dramatic consequences: $\Sigma(\xi, \mathbf{p})$ has an algebraic divergence proportional to $(\xi - \xi_{\mathbf{p}})^{d-2}$ for $\xi \to \xi_{\mathbf{p}}$, leading to two well-separated peaks of comparable weight in the spectral function, as shown for $d = 1.5$ in Fig. 7.4. The width of the peaks is finite in $d > 1$, but smaller than

[23] Analytic results for the second order self-energy within a one-dimensional g-ology model are given in Appendix D.

the distance between them.[24] Hence, second order perturbation theory seems to indicate destruction of the quasiparticle pole due to forward scattering of particles with almost parallel momenta in any dimension below two! However, the divergence found in $\Sigma(\xi, \mathbf{p})$ clearly forces us to go beyond perturbation theory even for weak coupling constants.

7.3. RANDOM PHASE APPROXIMATION

Our next step is the calculation of the small-q density-density response and the self-energy in random phase approximation (RPA). To leading order in q, the response function obtained by this "approximation" is actually the exact result for the regular normal model in any dimension, if the cutoff Λ is so small that the forward scattering interactions $f^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'} = g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(\mathbf{0})$ have already reached their fixed point values.

7.3.1. Effective interaction

The RPA *effective interaction* D is an important auxiliary quantity, which will play a central role not only in RPA calculations. It can be defined by the sum of diagrams illustrated already in Fig. 5.2, which is equivalent to the linear integral equation (5.21). For $\mathbf{k}$ and $\mathbf{k}'$ close to the Fermi surface and small q this can be simplified to

$$D^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(q) = g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(\mathbf{0}) + \sum_{\sigma''} \int_{\mathbf{k}''\in\partial\mathcal{F}} g^{\sigma\sigma''}_{\mathbf{k}\mathbf{k}''}(\mathbf{0}) \, \frac{\mathbf{n}_{\mathbf{k}''}\cdot\mathbf{q}}{\omega - \mathbf{v}_{\mathbf{k}''}\cdot\mathbf{q} + i0^+ s(\omega)} \, D^{\sigma''\sigma'}_{\mathbf{k}''\mathbf{k}'}(q) \tag{7.35}$$

with a surface integral extending over the Fermi surface $\partial\mathcal{F}$, where $\mathbf{n}_{\mathbf{k}}$ is a normal unit vector on the Fermi surface in $\mathbf{k}$. Obviously, for small q the effective action depends only via the ratio $\mathbf{q}/\omega$ on $q = (\omega, \mathbf{q})$. Equation (7.35) cannot be solved in general. There are however instructive special cases with drastic simplifications. In particular,

$$D^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(q) = D(q) = \frac{g}{1 - 2g\Pi_0(q)} \quad \text{for} \quad g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(\mathbf{q}) = g \tag{7.36}$$

where the factor 2 is due to the spin degeneracy, while

$$D^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(q) = D^{\sigma\sigma'}(q) = \begin{cases} g^2\Pi_0(q)/[1 - (g\Pi_0(q))^2] & \text{for} \quad \sigma = \sigma' \\ g/[1 - (g\Pi_0(q))^2] & \text{for} \quad \sigma = -\sigma' \end{cases}$$
$$\text{for} \quad g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(\mathbf{q}) = g\delta_{\sigma,-\sigma'} \tag{7.37}$$

The latter case, constant coupling between opposite spins, has been chosen for the perturbative evaluation of the self-energy in the preceding section. In both special cases the effective interaction depends only via the modulus $|\mathbf{q}|$ on $\mathbf{q}$. If in addition q is small, the effective interactions depend only on a single variable, e.g. the ratio $\tilde{\omega} = \omega/|\mathbf{q}|$.

[24] A similar behavior has been found recently near zero-curvature points on special anisotropic Fermi surfaces in two dimensions [126].

Numerical results for $D^{\sigma\sigma}(q)$ for small q as a function of $\tilde{\omega}$ in various dimensions are shown in Fig. 7.5, for the case (7.37) with $g = 2$ and $v_F = 1 = k_F$. In any dimension there is a singularity of the form (for $\tilde{\omega} > 0$)

$$D_c(q) = \frac{Z_c}{\tilde{\omega} - u_c + i0^+} \tag{7.38}$$

with $u_c > v_F$ associated with a propagating density mode. Apart from a δ-function in $\omega = u_c|\mathbf{q}|$, one has $\mathrm{Im}D^{\sigma\sigma}(q) = 0$ for $\omega > v_F|\mathbf{q}|$, as follows directly from the absence of particle-hole excitations with energies larger than $v_F|\mathbf{q}|$. For $\tilde{\omega} \to 0$, $\mathrm{Im}D^{\sigma\sigma}(q)$ vanishes linearly, and in $1 < d < 3$ as $(v_F - \tilde{\omega})^{(3-d)/2}$ for $\tilde{\omega} \to v_F - 0^+$. Close to one dimension, the effective interaction has a damped singularity at $\omega = u_s|\mathbf{q}|$ with a velocity $u_s < v_F$. For $d \to 1$ the damping vanishes, and one is left with an additional pole as in (7.38), but now in $\tilde{\omega} = u_s$, while all the spectral weight at other energies ($\omega \neq u_c|\mathbf{q}|, u_s|\mathbf{q}|$) has disappeared. Qualitatively the same behavior is found for other positive coupling strengths, too. The only difference in the case (7.36) is the absence of the second singularity at $\omega = u_s|\mathbf{q}|$.

7.3.2. Density-density response

The RPA charge density-density response (or correlation function) $N(q)$ can be constructed from the effective interaction in the way illustrated in Fig. 5.3, i.e.

$$N(q) = 2\Pi_0(q) + \sum_{\sigma\sigma'} \int_{k,k'} D^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(q)\, G_0(k - q/2)G_0(k + q/2)G_0(k' - q/2)G_0(k' + q/2) \tag{7.39}$$

For small q the energy-momentum integrals can be reduced to Fermi surface averages as before. For the two simple special cases introduced in 7.3.1 one obtains

$$N(q) = \frac{2\Pi_0(q)}{1 - 2g\Pi_0(q)} \quad \text{for} \quad g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(\mathbf{q}) = g \tag{7.40}$$

and

$$N(q) = \frac{2\Pi_0(q)}{1 - g\Pi_0(q)} \quad \text{for} \quad g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(\mathbf{q}) = g\delta_{\sigma,-\sigma'} \tag{7.41}$$

For $g > 0$ these functions have a pole in $\omega = u_c|\mathbf{q}|$, describing a propagating charge density mode, which is Landau's [1] zero-sound mode in our special system.

Similar results are easily obtained for the spin density-density response. At least one propagating (charge or spin) density mode exists in any Fermi liquid with short-range interactions [125].

7.3.3. RPA self-energy

We now calculate the self-energy in random phase approximation, concentrating in particular on the leading contributions from scattering with *small* momentum transfers,

which led to striking singularities in perturbation theory in dimensions $d < 2$. Diagrammatically, the RPA self-energy is described by Fig. 7.6, representing a propagator dressed once by the effective interation D. Algebraically, this reads

$$\Sigma(p) = i \int_{p'} D^{\sigma\sigma}_{\mathbf{kk}}(p-p')\, G_0(p') \quad \text{where} \quad \mathbf{k} = \frac{\mathbf{p}+\mathbf{p}'}{2} \tag{7.42}$$

In particular, the imaginary part can be expressed as

$$\mathrm{Im}\Sigma(\xi,\mathbf{p}) = \begin{cases} \int \frac{d^d\mathbf{p}'}{(2\pi)^d}\, \mathrm{Im}D^{\sigma\sigma}_{\mathbf{kk}}(\xi-\xi_{\mathbf{p}'},\mathbf{p}-\mathbf{p}')\, \Xi(0<\xi_{\mathbf{p}'}<\xi) & \text{for} \quad \xi>0 \\ -\int \frac{d^d\mathbf{p}'}{(2\pi)^d}\, \mathrm{Im}D^{\sigma\sigma}_{\mathbf{kk}}(\xi-\xi_{\mathbf{p}'},\mathbf{p}-\mathbf{p}')\, \Xi(\xi<\xi_{\mathbf{p}'}<0) & \text{for} \quad \xi<0 \end{cases} \tag{7.43}$$

We are interested in the contributions from small momentum transfers $\mathbf{q} = \mathbf{p}-\mathbf{p}'$ to $\mathrm{Im}\Sigma(p)$. A small $\mathbf{q}$ can be decomposed in normal and tangential components with respect to the Fermi surface in $\mathbf{p} \sim \mathbf{p}' \sim \mathbf{k}$, and the effective interaction can be parametrized as (see Eq. (5.21))

$$D^{\sigma\sigma}_{\mathbf{kk}}(\omega,\mathbf{q}) \;\sim\; D(\omega,q_r,q_t) \quad \text{for} \quad \mathbf{q} = \mathbf{p}-\mathbf{p}' \sim \mathbf{0} \tag{7.44}$$

We now consider contributions from scattering processes with tangential momentum transfers restricted by $q_t < \Lambda_t$ to the self-energy, where $\Lambda_t \ll k_F$. Decomposing the $\mathbf{p}'$-integral in a radial and an angular integral as in (7.6) and using $\sin\theta \sim \theta \sim q_t/k_F$, one obtains

$$\mathrm{Im}\Sigma^{\Lambda_t}(\xi,\mathbf{p}) = \int_0^{\xi/v_F} \frac{dp'_r}{2\pi}\, \mathrm{Im}\bar{D}^{\Lambda_t}(\xi - v_F p'_r, p_r - p'_r) \tag{7.45}$$

where $\bar{D}^{\Lambda_t}$ is the q_t-averaged effective interaction, defined by (cf. (5.22))

$$\bar{D}^{\Lambda_t}(\omega,q_r) = \frac{S_{d-1}}{(2\pi)^{d-1}} \int_0^{\Lambda_t} dq_t\, q_t^{d-2}\, D(\omega,q_r,q_t) \tag{7.46}$$

We will now show that for $p \to (0,\mathbf{k}_F)$ the leading contribution to $\mathrm{Im}\Sigma^{\Lambda_t}(p)$ obeys a scaling law with a scaling function $\mathrm{Im}\tilde{\Sigma}(\xi/p_r)$ that is independent of Λ_t in $d < 2$. To see this, we note that for short-range interactions the effective interaction depends on $(\omega,\mathbf{q})$ only via the ratio $\mathbf{q}/\omega$, if q is small, i.e. (7.44) reduces to

$$D^{\sigma\sigma}_{\mathbf{kk}}(\omega,\mathbf{q}) \sim D(q_r/\omega, q_t/|\omega|) \quad \text{for} \quad \mathbf{q} \sim \mathbf{0} \tag{7.47}$$

Introducing a rescaled variable $\tilde{q}_t = q_t/|\omega|$, one thus obtains

$$\begin{aligned} \mathrm{Im}\bar{D}^{\Lambda_t}(\omega,q_r) \;&= \frac{S_{d-1}}{(2\pi)^{d-1}}|\omega|^{d-1} \int_0^{\Lambda_t/|\omega|} d\tilde{q}_t\, \tilde{q}_t^{d-2}\, \mathrm{Im}D(q_r/\omega,\tilde{q}_t) \\ &\overset{\omega,q_r\to 0}{\longrightarrow}\; |\omega|^{d-1}\mathrm{Im}\tilde{D}(q_r/\omega) \end{aligned} \tag{7.48}$$

in $d < 2$, where

$$\mathrm{Im}\tilde{D}(q_r/\omega) = \frac{S_{d-1}}{(2\pi)^{d-1}} \int_0^{\infty} d\tilde{q}_t\, \tilde{q}_t^{d-2}\, \mathrm{Im}D(q_r/\omega,\tilde{q}_t) \tag{7.49}$$

Note that this integral is convergent in $d < 2$ (but not in $d \geq 2$), since $\mathrm{Im}D(q_r/\omega, q_t/|\omega|)$ vanishes linearly for $\omega \to 0$ at fixed $\mathbf{q}$, and thus the integrand is proportional to $\tilde{q}_t^{d-3}$ for large $\tilde{q}_t$. Inserting (7.48) in (7.45), one obtains

$$\mathrm{Im}\Sigma^{\Lambda_t}(\xi, \mathbf{p}) \sim s(p_r)|p_r|^d \, \mathrm{Im}\tilde{\Sigma}(\xi/p_r) \quad \text{for} \quad p \to (0, \mathbf{k}_F) \tag{7.50}$$

where

$$\mathrm{Im}\tilde{\Sigma}(\tilde{\xi}) = \int_0^{\tilde{\xi}/v_F} d\tilde{p}'_r \, |\tilde{\xi} - v_F\tilde{p}'_r|^{d-1} \, \mathrm{Im}\tilde{D}[(1-\tilde{p}'_r)/(\tilde{\xi}-v_F\tilde{p}'_r)] \tag{7.51}$$

The fact that the tangential cutoff Λ_t does not appear in the leading small-p scaling function implies in particular that contributions from momentum transfers with $q_t > \Lambda_t$ scale to zero more rapidly than $|p_r|^d$ in $d < 2$, however small Λ_t may be. The closer $(\xi, \mathbf{p})$ is to the Fermi surface, the smaller is the typical size of q_t contributing to the self-energy. Note that the radial momentum transfer q_r is also confined to a small interval, e.g. to $[p_r - \xi, p_r]$ for $\xi > 0$. In this sense the low-energy behavior of the RPA self-energy is dominated by small momentum transfers $|\mathbf{q}|$ in $d < 2$. We note that special Cooper processes with momentum transfers $|\mathbf{q}| \sim 2k_F$ would give rise to a contribution of the same order. However, we ignore this contribution, since their scattering amplitudes must vanish in the low-energy limit in a normal phase.

In Fig. 7.7 we show $\mathrm{Im}\Sigma(\xi, \mathbf{p})$ at $p_r = 0.1k_F$ for a constant coupling $g = 2$ between opposite spins (as in Fig. 7.3), in $d = 1.5$ dimensions. The divergence in $\xi = \xi_{\mathbf{p}}$ found in second order perturbation theory has disappeared. Instead one finds two finite peaks above and below $\xi_{\mathbf{p}}$, which are obviously due to low energy charge and spin density fluctuations (signalled by the corresponding peaks in the effective interaction D). In contrast to the perturbative result, $\tilde{\Sigma}(\tilde{\xi})$ is now a bounded function. Hence, in $1 < d < 2$ dimensions and for $\mathbf{p}$ sufficiently close to the Fermi surface, the RPA self-energy does not destroy the quasi-particle pole in the propagator, but leads only to a damping proportional to $|p_r|^d$, which is larger than the quadratic behavior known for three-dimensional Fermi liquids, but smaller than the quasi-particle energy $\xi_{\mathbf{p}} = v_F p_r$.

7.4. RESUMMATION OF FORWARD SCATTERING

We have learned from the perturbative and RPA analysis that in the absence of Cooper scattering (or "back-scattering", in $d = 1$) the self-energy of the regular normal model in $d < 2$ dimensions is dominated by scattering processes with small momentum transfers $\mathbf{q} \sim \mathbf{0}$, i.e. forward scattering. We now sum these dominant processes to all orders in the (residual) interaction by applying the results derived in Sec. 5. In one dimension this resummation is necessary to obtain the correct Luttinger liquid behavior of the single-particle propagator. Hence, such a resummation is also required to get a quantitative

control of the crossover from one-dimensional Luttinger liquid behavior to Fermi liquid behavior in dimensions $d > 1$.

Focussing on the role of forward scattering, we can introduce a momentum transfer cutoff $\Lambda_q \ll \Lambda, k_F$, and restrict momentum transfers by $|\mathbf{q}| < \Lambda_q$ (as done already in Secs. 5 and 6). Actually we will frequently work with two distinct cutoffs Λ_r and Λ_t for radial and tangential momentum transfers, respectively, and discuss their role separately. In explicit calculations we will usually cut off smoothly with an exponential factor, e.g. $e^{-|\mathbf{q}|/\Lambda_q}$, which is technically more convenient than a sharp cutoff.

In Sec. 5 we have shown that a resummation of forward scattering to all orders in the coupling leads to the effective one-dimensional equation of motion

$$(\xi - v_F p_r)\, G(\xi, p_r) \;=\; 1 \;+\; \int_{\xi', p_r'} \frac{i\bar{D}(\xi - \xi', p_r - p_r')}{\xi - \xi' - v_F(p_r - p_r')}\, G(\xi', p_r') \tag{7.52}$$

for the propagator G, which is valid to leading order in Λ_q (see (5.25)). Here the q_t-averaged effective interaction $\bar{D}$ is constructed from the RPA effective interaction D as in (7.46). As shown in (5.28)-(5.33), the solution for $G(\xi, p_r)$ is given by the Fourier transform of

$$G(t, r) = e^{L(t,r) - L_0}\, G_0(t, r) \tag{7.53}$$

where $L(t, r)$ is the Fourier transform of $L(\omega, q_r) = i\bar{D}(\omega, q_r)/(\omega - v_F q_r)^2$ and $G_0(t, r)$ the Fourier transform of $G_0(\xi, p_r)$; the constant L_0 is given by $L(t = 0, r = 0)$.

In the RPA calculation in 7.3 we have shown that the *imaginary* part $\mathrm{Im}\bar{D}$ is asymptotically independent of the tangential cutoff Λ_t for $q_r, \omega \ll \Lambda_t$ in $d < 2$ dimensions, since $\mathrm{Im}D(\omega, q_r, q_t)$ vanishes for $v_F|\mathbf{q}| \gg \omega$. By contrast, for the *real* part an arbitrary choice of effective couplings $g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(\mathbf{q})$ will generally lead to a finite limit for $v_F|\mathbf{q}| \gg \omega$, and the dependence on Λ_t does not disappear (except, of course, in one dimension). However, the effective low-energy couplings are not really independent. For a Fermi liquid it has been shown that the antisymmetry of the two-particle vertex poses a constraint on the Landau function $f^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'} = g^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(\mathbf{0})$, which is equivalent to the Landau sum-rule for Landau parameters [94]. As a consequence, effective interactions constructed from Landau functions as couplings satisfy [94]

$$D^{\sigma\sigma}_{\mathbf{k}\mathbf{k}}(\omega, \mathbf{q}) \to 0 \quad \text{for} \quad (\omega, \mathbf{q}) \to (0, \mathbf{0}) \quad \text{with} \quad |\mathbf{q}/\omega| \to \infty \tag{7.54}$$

For the behavior of D in a one-dimensional Luttinger liquid in that limit see the final paragraph of Sec. 6.4. Equation (7.54) implies that the real part of the effective interaction vanishes, too, for $v_F|\mathbf{q}| \gg \omega$ (recall that for short-range interactions D depends only on the ratio $\mathbf{q}/\omega$). Hence, the asymptotic behavior of $\bar{D}$ is given by the Λ_t-independent

expression

$$\bar{D}(\omega, q_r) \sim |\omega|^{d-1} \tilde{D}(q_r/\omega) = |q_r|^{d-1} |\tilde{\omega}|^{d-1} \tilde{D}(\tilde{\omega}^{-1}) \quad \text{for} \quad \omega, q_r \to 0 \tag{7.55}$$

where $\tilde{\omega} = \omega/q_r$ and

$$\tilde{D}(q_r/\omega) = \frac{S_{d-1}}{(2\pi)^{d-1}} \int_0^\infty d\tilde{q}_t \; \tilde{q}_t^{d-2} \; D(q_r/\omega, \tilde{q}_t) \tag{7.56}$$

The latter integral is convergent in $d < 2$ since $D(q_r/\omega, \tilde{q}_t)$ vanishes linearly for $\omega \to 0$ at fixed small $\mathbf{q}$, for forward scattering interactions that satisfy the Landau sum-rule.

We introduce an exponential cutoff $\Lambda_r =: \Lambda$ for *radial* momentum transfers in the expression (5.48) for $L(\omega, q_r)$. Using the asymptotic scaling behavior of $\bar{D}$, one has

$$L(\omega, q_r) = |q_r|^{d-1} e^{-|q_r|/\Lambda} \frac{i|\tilde{\omega}|^{d-1} \tilde{D}(\tilde{\omega}^{-1})}{[\omega - v_F q_r + i0^+ s(\omega)]^2} \tag{7.57}$$

A cutoff for q_r is necessary for short-range interactions in any dimension (including $d = 1$) to make the Fourier transform $L(t, r)$ well-defined. Changing integration variables in the Fourier transformation from ω and q_r to $\tilde{\omega}$ and q_r, the q_r integration can be carried out analytically, leading to

$$L(t, r) = \frac{1}{(2\pi)^2} \int_{-\infty}^\infty d\tilde{\omega} \; \frac{i|\tilde{\omega}|^{d-1} \tilde{D}(\tilde{\omega}^{-1})}{[\tilde{\omega} - v_F + i0^+ s(\tilde{\omega})]^2} \; I_{d,\Lambda}(r - \tilde{\omega} t) \tag{7.58}$$

where the function $I_{d,\Lambda}$ is given by

$$I_{d,\Lambda}(y) = \int_{-\infty}^\infty dq_r |q_r|^{d-2} e^{iq_r y - |q_r|/\Lambda} = \Gamma(d-1) \frac{2\cos[(d-1)\mathrm{arctg}(y\Lambda)]}{(y^2 + \Lambda^{-2})^{(d-1)/2}} \tag{7.59}$$

For $r, t \to 0$, one obtains from (7.58)

$$L_0 = L(0, 0) = \Lambda^{d-1} \tilde{L}_0 \tag{7.60}$$

where

$$\tilde{L}_0 = \frac{\Gamma(d-1)}{2\pi^2} \int_{-\infty}^\infty d\tilde{\omega} \; \frac{i|\tilde{\omega}|^{d-1} \tilde{D}(\tilde{\omega}^{-1})}{[\tilde{\omega} - v_F + i0^+ s(\tilde{\omega})]^2} \tag{7.61}$$

which is a cutoff-independent number. Note that the limit $r, t \to 0$ is unique, i.e. it is not important whether t or r tends to zero first.

Appendix B contains expressions for $L(t, r)$ and L_0 in terms of the spectral function $\tilde{\Delta}$ associated with $\tilde{D}$.

For $r, t \to \infty$, $\tilde{t} = t/r$ fixed and arbitrary, the function $L(t, r)$ scales as

$$L(t, r) \sim |r|^{1-d} \tilde{L}(\tilde{t}) \tag{7.62}$$

where

$$\tilde{L}(\tilde{t}) = \frac{1}{(2\pi)^2}\int_{-\infty}^{\infty} d\tilde{\omega}\, \frac{i|\tilde{\omega}|^{d-1}\tilde{D}(\tilde{\omega}^{-1})}{[\tilde{\omega} - v_F + i0^+ s(\tilde{\omega})]^2}\, \tilde{I}_d(1-\tilde{\omega}\tilde{t}) \tag{7.63}$$

and

$$\tilde{I}_d(y) = \frac{2\Gamma(d-1)}{|y|^{d-1}}\cos[(d-1)\pi/2] \tag{7.64}$$

which is independent of the cutoff Λ.

We now consider the functions $L(0,r)$ and $L(t,0)$, which determine the momentum distribution function and the density of states for single-particle excitations, respectively.

The function $L(t,r)$ has a unique limit for $t \to 0$, which can be written as

$$L(0,r) = \tilde{L}_0\, (r^2 + \Lambda^{-2})^{(1-d)/2}\, \cos[(d-1)\mathrm{arctg}(\Lambda r)] \tag{7.65}$$

where $\tilde{L}_0$ is the constant defined in (7.61). For $|r| \gg \Lambda^{-1}$, one finds

$$L(0,r) \sim \tilde{L}_0\, \cos[(d-1)\pi/2]\, |r|^{1-d} \tag{7.66}$$

The asymptotic behavior of $G(0^{\pm},r)$ for large $|r|$ is thus given by

$$G(0^{\pm},r)\ \sim\ \exp\left[\tilde{L}_0 \cos[(d-1)\pi/2]|r|^{1-d} - \tilde{L}_0\Lambda^{d-1}\right] G_0(0^{\pm},r) \tag{7.67}$$

and the leading large-$|r|$ behavior is

$$G(0^{\pm},r) \to e^{-L_0} G_0(0^{\pm},r) \quad \text{for} \quad |r| \to \infty \tag{7.68}$$

The *momentum distribution* $n_{\mathbf{p}}$ is obtained by Fourier transforming $G(0^-,r)$ as in (5.34), i.e. $n_{\mathbf{p}} = -i\int_{-\infty}^{\infty} dr\, G(0^-,r)\, e^{-ip_r r}$. From (7.68) we see that the discontinuity in $n_{\mathbf{p}}$ is reduced by a factor

$$\bar{Z} = e^{-L_0} \tag{7.69}$$

Here we have to remember that the above propagator and momentum distribution function are actually renormalized quantities, calculated for an effective low-energy action $\bar{S}^{\Lambda}$. A wave function renormalization Z^{Λ} had been performed already in passing from a microscopic description to the effective action in (7.1), while the above $\bar{Z}$ is the additional renormalization due to residual forward scattering. Thus, the total renormalization Z, which is equal to the discontinuity of the unrenormalized (i.e. physical) momentum distribution function, can be written as

$$Z = Z^{\Lambda}\bar{Z} \tag{7.70}$$

In Appendix B it is shown that the constant L_0 is a positive real number, which guarantees that $\bar{Z} \leq 1$, as it should be.

The limit $r \to 0$ in $L(t,r)$ is also unique. In particular, for $|t| \to \infty$ (i.e. $v_F|t| \gg \Lambda^{-1}$) one obtains the scaling behavior

$$L(t,0) \sim \hat{L}|t|^{1-d} \tag{7.71}$$

where

$$\hat{L} = \frac{\Gamma(d-1)}{2\pi^2} \int_{-\infty}^{\infty} d\tilde{\omega}\, \frac{i\tilde{D}(\tilde{\omega}^{-1})}{[\tilde{\omega} - v_F + i0^+ s(\tilde{\omega})]^2}\, \cos[(d-1)\pi/2] \tag{7.72}$$

which is finite and independent of Λ. The asymptotic behavior of $G(t,0)$ is thus

$$G(t,0) \;\sim\; e^{\hat{L}|t|^{1-d} - L_0}\, G_0(t,0) \tag{7.73}$$

In $d > 1$, $L(t,0)$ vanishes for $|t| \to \infty$, i.e. eventually

$$G(t,0) \;\to\; e^{-L_0} G_0(t,0) \quad \text{for} \quad |t| \to \infty \tag{7.74}$$

Fourier transforming $G(t,0)$ as in (5.35) yields the *density of states* (per spin), $N(\xi) = -\pi^{-1} s(\xi)\, \mathrm{Im} \int_{-\infty}^{\infty} dt\, G(t,0) e^{i\xi t}$. Hence, the factor $\bar{Z} = e^{-L_0}$ determines the reduction of the density of states at the Fermi level $\omega = 0$. Since the same factor determined the reduction of the discontinuity, the Fermi velocity v_F is obviously not renormalized by the residual interactions.

7.5. CRITICAL DIMENSIONS AND DIMENSIONAL CROSSOVER

7.5.1. Leading low-energy behavior

The above results imply that to *leading* order in a low-energy expansion, Fermi liquid behavior is not destroyed by residual scattering in any dimension $d > d_c = 1$. This follows from the behavior of the function $L(t,r)$ for large distances, which according to (7.62) scales as

$$L(st, sr) = s^{1-d} L(t,r) \tag{7.75}$$

and hence vanishes for $r, t \to \infty$ in dimensions $d > 1$. Strictly speaking, this conclusion is justified only if $r/v_F t \neq 1$ when taking the large-distance limit, because the scaling function $\tilde{L}(\tilde{t})$ in (7.63) diverges for $v_F \tilde{t} = 1$. Thus, in the special limit $r, t \to \infty$ with $r/v_F t = 1$, the function $L(t,r)$ does not vanish and may even diverge. This behavior can be traced back to the neglect of higher order terms in the small-$\mathbf{q}$ expansion, especially in deriving the asymptotic Ward identities (5.11) and (5.13) for the charge and spin density vertices. As discussed already in the final paragraph of 5.3, in the limit $r, t \to \infty$ with $r/v_F t = 1$ the function $L(t,r)$ is dominated by Fourier components $L(\omega, q_r)$ with $\omega \sim v_F q_r$, where subleading corrections to the asymptotic Ward identities should be important, because they cut off the poles in (5.11) and (5.13). In all other cases one

integrates over many values for the ratio $\omega/v_F q_r$, and corrections are generally negligible. In short, the singular behavior of our result for $G(t,r)$ in the special large-distance limit with $r/t \to v_F$ is an artefact produced by truncating the small-$\mathbf{q}$ expansion at leading order. Since explicit expressions for the corrections are not available, one cannot calculate the behavior in that special limit quantitatively.

Equation (7.75) and the above arguments imply that

$$G(t,r) \;\to\; e^{-L_0} G_0(t,r) \quad \text{for} \quad r,t \to \infty \quad \text{in} \quad d > d_c = 1 \tag{7.76}$$

at least for $r \neq v_F t$, and most plausibly also for $r = v_F t$. Fourier transforming, and "unrenormalizing" the propagator by multiplying with Z^Λ, we obtain

$$G(p) \;\to\; \frac{Z}{\xi - v_F p_r + i0^+ s(\xi)} \quad \text{for} \quad p \to (0, \mathbf{k}_F) \quad \text{in} \quad d > d_c = 1 \tag{7.77}$$

where $Z = Z^\Lambda e^{-L_0}$. By contrast, in one dimension $G(p)$ is given by the very different Luttinger liquid form (see Sec. 6). Hence, the random phase approximation to the self-energy in 7.3 gave the correct criterion on the stability of the quasi-particle pole in d dimensions, while second order perturbation theory led to misleading results.

An earlier analysis by Ueda and Rice [74] also indicated the possible stability of the quasi-particle pole with respect to residual interactions in dimensions above but close to one. Their argument was based on an ϵ-expansion around a one-dimensional system, with $\epsilon = d-1$. The coupling space was restricted to g_1- and g_2-type couplings (in g-ology notation), while g_4 and the generation of other couplings in higher dimensions was ignored. With these simplifying assumptions, Ueda and Rice calculated the flow of g_1 and g_2 to one-loop order in $d = 1+\epsilon$, finding that for $\epsilon > 0$ these couplings scale either to zero, or to strong coupling. The former result was interpreted as a Fermi liquid fixed point, and the latter as a signal for an instability towards ordered phases. Embedding this calculation in our more general context we would say that Ueda and Rice have correctly calculated the flow of Cooper couplings, which indeed flow either to zero or to large values. However the role of forward scattering couplings in $d > 1$, which do not scale to zero but nevertheless do not destroy the quasi-particle pole, has not been addressed in this earlier work.

We now analyze how the wave function renormalization Z vanishes in the limit $d \to 1$. We recall from (7.70) that Z is a product of two factors, namely Z^Λ for arriving from a microscopic model at the effective action $\bar{S}^\Lambda$, and the factor $\bar{Z} = e^{-L_0}$ from residual scattering within $\bar{S}^\Lambda$. Of course only the latter is controlled non-perturbatively by our results. Remember also that Z measures the jump in the momentum distribution function and the weight of the quasi-particle peak in the spectral function. Above we have shown that

$$\bar{Z} = Z/Z^\Lambda = e^{-\tilde{L}_0 \Lambda^{d-1}} \tag{7.78}$$

where $\tilde{L}_0$ is a cutoff independent number given by (7.61). The right hand side of (7.78) is the renormalization due to residual scattering in the effective action, and the expression is valid to all orders in the coupling constants, provided that $\Lambda \ll k_F$. From Eq. (C.12) we see that $\bar{Z}$ is a finite number between zero and one in $d > 1$. For $d \to 1$, however, $\tilde{L}_0$ diverges because the prefactor $\Gamma(d-1) \sim (d-1)^{-1}$ diverges, and consequently $\bar{Z}$ (and hence Z) vanishes, as expected. Quantitatively, one finds

$$\bar{Z} \to e^{-(\eta/\epsilon)\Lambda^\epsilon} \quad \text{for} \quad d \to 1 \tag{7.79}$$

where $\epsilon = d - 1$ is the deviation from one-dimension, and

$$\eta := \lim_{d\to 1} (d-1)\tilde{L}_0 = \frac{1}{2\pi^2} \int_{-\infty}^{\infty} d\tilde{\omega}\, \frac{i|\tilde{\omega}|^{d-1}\tilde{D}(\tilde{\omega}^{-1})}{[\tilde{\omega} - v_F + i0^+ s(\tilde{\omega})]^2} \tag{7.80}$$

is nothing but the anomalous scaling dimension determined by the Luttinger liquid one approaches in the limit $d \to 1$! This latter identification can be obtained directly from the solution of the one-dimensional model (see Appendix E). Alternatively, the identification can be made by ϵ-expanding our result (7.67) for $G(0, r)$: Expanding, for $r > 0$, $\tilde{L}_0 = \eta/\epsilon + \mathcal{O}(1)$, $r^{1-d} = 1 - \epsilon \log r + \mathcal{O}(\epsilon^2)$, $\cos[(d-1)\pi/2] = 1 + \mathcal{O}(\epsilon^2)$, and $\Lambda^{d-1} = 1 + \epsilon \log \Lambda + \mathcal{O}(\epsilon^2)$, one finds $L(0, r) - L_0 = -\eta \log(r\Lambda) + \mathcal{O}(\epsilon)$, and thus

$$G(0, r) \propto r^{-1-\eta} \quad \text{in} \quad d = 1 \tag{7.81}$$

We finally note that the constant $\hat{L}$ in (7.72) becomes equal to $\tilde{L}_0$ for $d \to 1$, and is thus also given by η/ϵ.

7.5.2. Subleading corrections

For $|r| \gg \Lambda^{-1}$ and $|t| \gg (v_F\Lambda)^{-1}$, the (renormalized) propagator can be written as

$$G(t, r) = e^{|r|^{1-d}\tilde{L}(t/r) - L_0} G_0(t, r) \tag{7.82}$$

where the scaling function $\tilde{L}(\tilde{t})$ is given by (7.63), and the constant L_0 by (7.60) and (7.61). For $|r| \gg \Lambda^{-1}$ at $t = 0^{\pm}$, the asymptotic behavior of G is given by (7.67), and for $|t| \gg (v_F\Lambda)^{-1}$ at $r = 0$ by (7.73). In 7.5.1 we have discussed only the *leading* low-energy (or long-distance) behavior of G, obtained in the limit where only the constant L_0 survives in the exponent. At any finite energy (or, distance), the (t, r)-dependent term yields *subleading* corrections. For dimensions close to $d = 1$, these vanish very slowly as a function of energy (or distance), and *in the limit $d \to 1$ they merge with the leading terms*, giving rise to a different (Luttinger-type) leading behavior.

To discuss these subleading corrections, let us write the exact (and unrenormalized) propagator in the vicinity of the quasi-particle pole in a form customary in Fermi liquid theory, as

$$G(p) \;=\; \frac{Z}{\xi - \xi^*_{\mathbf{p}} \pm i\gamma_{\mathbf{p}}} \tag{7.83}$$

where $\xi^*_{\mathbf{p}}$ is the quasi-particle energy and $\gamma_{\mathbf{p}}$ the quasi-particle decay rate (see eq. (4.27)). In three dimensions, subleading corrections to the Fermi liquid fixed point scale quadratically to zero in the low-energy limit, i.e. $\xi^*_{\mathbf{p}} - v_F p_r = \mathcal{O}(p_r^2)$ and $\gamma_{\mathbf{p}} = \mathcal{O}(p_r^2)$ [2]. Perturbation theory and RPA as described in 7.2 and 7.3 yield the same behavior in any dimension $d > 2$. In exactly two dimensions, logarithmic corrections to the quadratic behavior are known to occur [127, 128]. Below two dimensions, perturbative and RPA results have indicated a scaling of corrections with a power d instead of two. This is confirmed by our resummation of forward scattering: We have found large-distance corrections of order $|t|^{1-d}$ or $|r|^{1-d}$, which, by Fourier transform, correspond to low-energy corrections with a scaling power d, i.e.

$$\begin{aligned} \xi^*_{\mathbf{p}} - v_F p_r &= \mathcal{O}(p_r^d) \\ \gamma_{\mathbf{p}} &= \mathcal{O}(p_r^d) \end{aligned} \qquad \text{in} \quad 1 < d < 2 \tag{7.84}$$

Thus, a normal Fermi system with short-range interactions has *two characteristic dimensions*, where the low-energy behavior undergoes qualitative changes. In any dimension above the *critical dimension* $d_c = 1$ one has asymptotically stable quasi-particles, but below $d = 2$ their decay rate and other subleading corrections are bigger than the quadratic behavior known from three-dimensional Fermi systems.

A phase-diagram in the dimensionality-coupling plane summarizing the behavior of the regular normal model can be found in the conclusions (Sec. 10).

8. LONG-RANGE DENSITY-DENSITY INTERACTIONS

So far we have restricted our analysis to Fermi systems with pure short-range interactions, corresponding to bounded coupling functions $g_{kk';q}$ in momentum space. In this and the following section we analyze the low-energy physics of systems with *long-range* interactions. Sec. 9 deals with fermions coupled to an abelian gauge field, which gives rise to an effective long-range current-current interaction between fermions.

Here we consider a class of long-range *density-density* interactions, with coupling functions that diverge in the forward scattering channel. To be specific, we analyze interactions with an arbitrary power-law decay in real space. The corresponding coupling functions in momentum space depend only on the momentum transfer $\mathbf{q}$, not on $\mathbf{k}$ and $\mathbf{k}'$ (nor on spin), where the $\mathbf{q}$-dependence is given by a power-law

$$g(\mathbf{q}) = g_0/|\mathbf{q}|^\gamma \,, \quad g_0 > 0 \,, \quad \gamma > 0 \tag{8.1}$$

In real space, this corresponds to a pair-potential $V(\mathbf{r}) \propto 1/|\mathbf{r}|^{d-\gamma}$. This generalization of the Coulomb interaction has been introduced by Bares and Wen [37], who also first derived the most important physical consequences of such an interaction. In the Coulomb case one has $\gamma = d-1$ and $g_0 = \Gamma(d-1)S_d\, e^2$ in d dimensions, where e is the electron charge. Multiplying $g(\mathbf{q})$ by the bare density of states at the Fermi level, N_F, one may define a dimensionless coupling function

$$\tilde{g}(\mathbf{q}) = N_F\, g(\mathbf{q}) = (\kappa/|\mathbf{q}|)^\gamma \tag{8.2}$$

where $\kappa = (N_F g_0)^{1/\gamma}$ is a generalized Thomas-Fermi screening wave vector [106].

Since $g(\mathbf{q})$ diverges for $\mathbf{q} \to \mathbf{0}$, the importance of scattering processes with small momentum transfers is enhanced with respect to the case of short-range interactions. The screening vector κ sets a scale in momentum space that separates a weak coupling region for $|\mathbf{q}| \gg \kappa$ from a strong coupling region for $|\mathbf{q}| \ll \kappa$. For systems with $\kappa \ll k_F$, only interactions with small momentum transfers $|\mathbf{q}| \ll k_F$ contribute significantly, such that the techniques developed for forward scattering dominated systems derived in Sec. 5 are adequate. In particular, for continuum systems with a free-particle dispersion $\epsilon_{\mathbf{k}} = \mathbf{k}^2/2m$ one has $N_F \propto k_F^{d-2}$ and thus $\kappa \propto k_F^{(d-2)/\gamma}$. Hence, one has $\kappa \ll k_F$ in the high-density limit $k_F \to \infty$ if $d < 2+\gamma$, as for example in the high-density Coulomb gas in any dimension [73].

In the following we discuss the low-energy small-$\mathbf{q}$ density-density response and the low-energy behavior of the single-particle propagator in systems specified by (8.1), concentrating on contributions from scattering processes with $|\mathbf{q}| \ll k_F$. We will rely on the general techniques and results derived in Sec. 5. These techniques have been applied to the present problem in their Ward identity version [38] as well as in bosonization language [73, 106, 129]. The results obtained by Bares and Wen [37] are thereby confirmed and extended.

As a consequence of loop cancellation (see Sec. 5) in forward scattering dominated systems, the leading low-energy small-$\mathbf{q}$ charge density-density response function is given by the RPA form

$$N(q) = \frac{2\Pi_0(\omega, \mathbf{q})}{1 - 2g(\mathbf{q})\Pi_0(\omega, \mathbf{q})} \tag{8.3}$$

This function has a real pole in $|\omega| = \omega_c(\mathbf{q})$, corresponding to a propagating collective excitation. For $\gamma > 0$ and small $\mathbf{q}$ it is clear that $\Pi_0(\omega, \mathbf{q})$ must be small and positive to make $|\omega| = \omega_c(\mathbf{q})$ satisfy the equation $|\mathbf{q}|^\gamma = 2g_0\Pi_0(\omega, \mathbf{q})$ for the position of the pole. This implies that $\omega_c(\mathbf{q})$ is much larger than $v_F|\mathbf{q}|$ for small $\mathbf{q}$. For $|\omega| \gg v_F|\mathbf{q}|$, the particle-hole bubble has the form

$$\Pi_0(q) \sim \frac{N_F}{2d} \frac{v_F^2|\mathbf{q}|^2}{\omega^2}, \quad |\omega| \gg v_F|\mathbf{q}| \tag{8.4}$$

Inserting this in the equation for the pole, one obtains

$$\omega_c(\mathbf{q}) \sim \lambda|\mathbf{q}|^{1-\gamma/2} \quad \text{where} \quad \lambda = v_F(N_F g_0/d)^{1/2} \tag{8.5}$$

We stress that this simple form holds only for small $\mathbf{q}$. Obviously $\omega_c(\mathbf{q})$ is the dispersion of a *gapless* mode as long as $\gamma < 2$. For the Coulomb interaction in $d = 3$ one has $\gamma = 2$, and $\omega_c(\mathbf{q})$ is the well-known plasmon mode with a gap given by the plasma frequency $\lambda = \omega_p$. For $\gamma > 2$, the dispersion relation $\omega_c(\mathbf{q})$ has a gap, too.

We now analyze the low-energy behavior of the single-particle propagator G and related quantities. We focus on the regime $0 < \gamma < 2$, where $\omega_c(\mathbf{q}) \to 0$ for $|\mathbf{q}| \to 0$. We would like to see in particular how strongly the single-particle propagator is affected by the presence of the gapless collective mode, and under which conditions Fermi liquid behavior is destroyed.

The RPA effective interaction associated with $g(\mathbf{q})$ is

$$D^{\sigma\sigma'}_{\mathbf{k}\mathbf{k}'}(q) = D(q) = \frac{g(\mathbf{q})}{1 - 2g(\mathbf{q})\Pi_0(\omega, \mathbf{q})} = \frac{g_0}{|\mathbf{q}|^\gamma - 2g_0\Pi_0(\omega, \mathbf{q})} \tag{8.6}$$

This function has the same denominator as $N(q)$, and thus the same pole in $|\omega| = \omega_c(\mathbf{q})$. It is convenient to decompose the effective interaction as $D(q) = D_c(q) + D_{reg}(q)$, where

$$D_c(q) \;=\; \lambda g_0 |\mathbf{q}|^{1-3\gamma/2} \frac{\omega_c(\mathbf{q})}{\omega^2 - \omega_c^2(\mathbf{q})} \;=\; \lambda^2 g_0 \frac{|\mathbf{q}|^{2-2\gamma}}{\omega^2 - \lambda^2 |\mathbf{q}|^{2-\gamma}} \tag{8.7}$$

contains the pole, while $D_{reg}(q)$ is a regular function which is finite in the limit $q \to 0$. The effect of the regular part $D_{reg}(q)$ on the propagator is qualitatively the same as for short-range interactions. In particular, it cannot destroy the quasi-particle pole in dimensions $d > 1$. Hence, we now concentrate on the singular part $D_c(q)$ only. In contrast to the effective interaction for short-range interactions, $D_c(q)$ does not depend only on the ratio $\mathbf{q}/\omega$. Instead, one has the scaling behavior

$$D_c(q) \mapsto b^{-\gamma} D_c(q) \quad \text{for} \quad \mathbf{q} \mapsto b\,\mathbf{q}\,, \quad \omega \mapsto b^{1-\gamma/2}\omega \tag{8.8}$$

Note that this scaling transformation is anisotropic in $(\omega, \mathbf{q})$-space (the dynamical scaling index $z = 1 - \gamma/2$ differs from one).

Since the interaction $g(\mathbf{q})$ is singular only for $\mathbf{q} \to \mathbf{0}$, it is clear that only scattering processes with a small momentum transfer can have a drastic effect on the propagator. For these processes we can apply the results derived in Sec. 5. We define an angular averaged effective interaction

$$\bar{D}_c^{\Lambda_t}(\omega, q_r) = \frac{S_{d-1}}{(2\pi)^{d-1}} \lambda^2 g_0 \int_0^{\Lambda_t} dq_t \; q_t^{d-2} \frac{(q_r^2 + q_t^2)^{1-\gamma}}{\omega^2 - \lambda^2 (q_r^2 + q_t^2)^{1-\gamma/2}} \tag{8.9}$$

where $\Lambda_t \ll k_F$ is a cutoff for tangential momentum transfers. This quantity obeys the scaling relation

$$\bar{D}_c^{b\Lambda_t}(b^{1-\gamma/2}\omega, b q_r) = b^{d-1-\gamma} \bar{D}_c^{\Lambda_t}(\omega, q_r) \tag{8.10}$$

Applying the results from Sec. 5, the modification of the propagator by scattering processes with $|\mathbf{q}| < \Lambda \ll k_F$ is given by Eqs. (5.28)-(5.33) with the above $\bar{D}_c^{b\Lambda_t}$ and $\Lambda_t = (\Lambda^2 - q_r^2)^{1/2}$. From (5.30) and (8.10) one obtains the scaling behavior at large distance and long time

$$L(s^{1-\gamma/2}t, sr) = s^{1-d+\gamma/2} L(t, r) \tag{8.11}$$

Thus, for dimensions above the *critical dimension*

$$d_c^\gamma = 1 + \gamma/2 \tag{8.12}$$

scattering processes with small momentum transfers do not modify the leading long-distance asymptotics of $G(t, r)$ with respect to the Fermi liquid behavior. By contrast, for $d \leq d_c^\gamma$ Fermi liquid behavior is completely destroyed [38, 73, 129]. This result from

a resummation of forward scattering processes confirms the earlier result obtained by Bares and Wen [37], which was based on an RPA calculation of the self-energy and other approximate methods. Note that d_c^γ is smaller than $1+\gamma$, which is what one might expect from a (very) naive power-counting argument. Counting the power of the bare coupling $g(\mathbf{q})$ misses screening, which weakens the singularity, and does not take into account the modified dynamical scaling, eq. (8.8), imposed by the "plasmon" mode.

For $d < 1+\gamma$ the angular averaged effective interaction becomes independent of Λ_t for $\omega, q_r \to 0$. Defining $\tilde{q}_t = q_t/|q_r|$ and $\tilde{\omega} = \omega/|q_r|^{1-\gamma/2}$, one can write

$$\bar{D}_c^{\Lambda_t}(\omega, q_r) \;\to\; |q_r|^{d-1-\gamma} \tilde{D}_c(\tilde{\omega}) \quad \text{for} \quad \omega, q_r \to 0 \tag{8.13}$$

where the scaling function $\tilde{D}_c(\tilde{\omega})$ is given by

$$\tilde{D}_c(\tilde{\omega}) = \frac{S_{d-1}}{(2\pi)^{d-1}} \lambda^2 g_0 \int_0^\infty d\tilde{q}_t \; \tilde{q}_t^{d-2} \frac{(1+\tilde{q}_t^2)^{1-\gamma}}{\tilde{\omega}^2 - \lambda^2 (1+\tilde{q}_t^2)^{1-\gamma/2}} \tag{8.14}$$

which is independent of tangential cutoffs.

Explicit results have so far been obtained for the propagator $G(0,r)$ at $t = 0$ and especially for the quasi-particle weight Z.

The quasi-particle weight is related to the function $L(t,r)$ via $Z = e^{-L_0}$ where $L_0 = L(0,0)$ (cf. Sec. 7). Using the spectral representation for the effective interaction D (see App. C), one can express L_0 as [106]

$$L_0 = \frac{\lambda g_0}{2} \int \frac{d^d\mathbf{q}}{(2\pi)^d} \frac{|\mathbf{q}|^{1-3\gamma/2}}{[\omega_c(\mathbf{q}) + |\mathbf{v}_{\mathbf{k}_F} \cdot \mathbf{q}|]^2} \;+\; \text{continuum contributions} \tag{8.15}$$

The first term is exclusively due to the collective mode in the effective interactions. For $d \le d_c^\gamma = 1 + \gamma/2$ this term is infrared divergent, i.e. Z vanishes, signalling again the breakdown of Fermi liquid behavior in this case [73]. For $d < 1+2\gamma$ the $\mathbf{q}$-integrals in (8.15) are convergent at large $\mathbf{q}$ even without ultraviolet cutoff; the screening vector κ acts as a natural cutoff in this situation [73]. Hence, for $d_c^\gamma < d < 1+2\gamma$ one obtains a finite *cutoff-independent* expression for Z (with collective mode and continuum contributions). The Coulomb interaction satisfies these conditions in any dimension, and the result obtained for Z from the above resummation of forward scattering is expected to be asymptotically exact in the high-density limit. A simple analytic expression has been derived for the spinless Coulomb gas close to one dimension [73]:

$$Z \sim \sqrt{k_F/\kappa}\, e^{-1/2(d-1)} \quad \text{for} \quad d \to 1 \quad \text{at fixed} \quad \kappa \ll k_F \tag{8.16}$$

In the *marginal* case $d = d_c^\gamma$ the propagator $G(0,r)$ obeys a power law decay with an anomalous dimension [106]

$$\eta = \frac{\sqrt{d}}{2} (\kappa/k_F)^{d-1} \tag{8.17}$$

implying a corresponding power-law behavior of the momentum distribution function near the Fermi surface as in a one-dimensional Luttinger liquid. This power-law has been first obtained by Bares and Wen [37] from a hydrodynamical calculation of the orthogonality catastrophe as well as from an exponentiation of a logarithmic divergence found in the RPA self-energy.

9. FERMIONS COUPLED TO A GAUGE FIELD

The physics of two-dimensional Fermi systems coupled to a $U(1)$ gauge-field has recently become important in two different contexts. One is the theory of the half-filled Landau level proposed by Halperin, Lee and Read [44] and by Kalmeyer and Zhang [45], where the gauge-field is associated with a ficticious magnetic flux attached to the electrons [130, 131, 132]. The other is the theory of high-temperature superconductors, where gauge-fields have been introduced by Baskaran and Anderson [39], and subsequently by Ioffe and Larkin [40] and Lee and Nagaosa [41], to describe "spin liquids", i.e. correlated electron systems with local constraints imposed by strong repulsive forces. The singular behavior of the gauge-field propagator at small q generates strong forward scattering of the fermions in these systems.[25] The main issue is to what extent Fermi liquid behavior is thereby destroyed.

Many years ago Holstein, Norton and Pincus [42] have already pointed out that the transverse component of the electromagnetic field is not screened in a metal and leads to non-Fermi liquid properties even in three-dimensional systems (see also Ref. [43]). However, due to the smallness of the ratio v_F/c in usual metals these effects have a tiny energy scale and are thus hard to observe. By contrast, in the above-mentioned more recent physical contexts, the dimensionless coupling constant is not small. In addition, the reduced dimensionality $d = 2$ enhances the phase space for forward scattering.

The problem of treating the infrared singularities in a Fermi system coupled to a gauge-field has been addressed by numerous methods: Self-consistent approximations [135], bosonization [140, 141], asymptotic Ward identities [63, 38], eikonal expansion [64], $1/N$-expansion [135, 63] and renormalization group [133, 142, 136]. Polchinski [135] and Ioffe, Lidsky and Altshuler [63] have introduced an arbitrary number of fermion species (or "flavor number") N as a control and expansion parameter. Expansions around both limits $N \to 0$ and $N \to \infty$, as carried out in much detail by Altshuler, Ioffe and Millis [63], have helped to assess the regime of validity of several earlier approaches. Our review is restricted to pure systems; for the interesting interplay between disorder and fluctuating gauge-fields the reader should consult the papers by Aronov and Wölfle [137].

[25] A similar strong forward scattering is obtained for Fermi systems near a Pomerantchuk instability towards phase separation [51].

9.1. ACTION

The theory of the *half-filled Landau level* leads to an action [44, 45]

$$S = S_0 + S_{CS} + S_I \tag{9.1}$$

which involves a fermionic field ψ and a $U(1)$ gauge-field a_μ. For fully polarized systems in a strong external magnetic field, ψ is a one-component (or "spinless") fermionic field. In euclidean (imaginary time) representation the various terms in (9.1) are defined as follows. The first term

$$S_0 = \int dx \left\{ \psi^*(x) \left[-\partial_0 + \mu - i a_0 \right] \psi(x) \; - \; \frac{1}{2m} \, \psi^*(x) \left[-i\nabla - \mathbf{a}(x) \right]^2 \psi(x) \right\} \tag{9.2}$$

describes non-relativistic free fermions coupled minimally to the gauge-field $a = (a_0, \mathbf{a})$. The integration extends over the (2+1)-dimensional euclidean space-time $x = (x_0, \mathbf{x})$. The scalar field a_0 and the Chern-Simons term

$$S_{CS} = (2\pi\tilde{\phi})^{-1} \int dx \, i a_0(x) \, \nabla \times \mathbf{a}(x) \tag{9.3}$$

impose the *constraint* $\nabla \times \mathbf{a}(x) = 2\pi\tilde{\phi}\, \psi^*(x)\psi(x)$ that links the flux tubes to the electrons. The number of flux quanta $\tilde{\phi}$ attached to each electron is chosen equal to two in the theory of the half-filled Landau level. With this choice the external magnetic field is cancelled by the average (mean-field) ficticious gauge-field and the Aharonov-Bohm phase of the flux tubes does not transmute the fermionic statistics, i.e. the composite object given by an electron and its flux tube is a fermion. Electron-electron interactions are described by

$$S_I = -\tfrac{1}{2} \int dx dx' \; \psi^*(x)\psi(x) \; v(\mathbf{x}-\mathbf{x}') \; \psi^*(x')\psi(x') \tag{9.4}$$

where $v(\mathbf{x})$ is a Coulomb potential proportional to $e^2/|\mathbf{x}|$, or an effective short-range interaction in systems where the long-range part is screened by charges outside the two-dimensional electron gas, e.g. by a metallic gate. Using the above-mentioned constraint, S_I can also be expressed as a quadratic form in the gauge-field $\mathbf{a}$ instead of a quartic monomial in fermionic fields.

The theory of *spin liquids* constructed in the context of high-T_c superconductors has led to the action [39, 40, 41]

$$S = \int dx \left\{ \psi^*(x) \left[-\partial_0 + \mu \right] \psi(x) \; - \; \frac{1}{2m} \, \psi^*(x) \left[-i\nabla - \mathbf{a}(x) \right]^2 \psi(x) \right\} \tag{9.5}$$

where $\psi = \binom{\psi_\uparrow}{\psi_\downarrow}$ is now a two-component spinor. This action is sometimes supplemented by a pure gauge term

$$S_a = -\frac{1}{4g^2} \int dx \, f_{\mu\nu}^2 \quad \text{where} \quad f_{\mu\nu} = \partial_\mu a_\nu - \partial_\nu a_\mu \tag{9.6}$$

which is generated by integrating out high-energy processes (g is an effective coupling constant). We emphasize that the action S ($+S_a$) is only a part of what is expected to describe the strongly correlated metallic phase in high-T_c materials, since the holon part [39] is missing. Nevertheless, the above action has attracted considerable theoretical interest in its own right.

Note that we use the Coulomb gauge $\nabla \cdot \mathbf{a} = 0$ throughout the section.

9.2. GAUGE-FIELD PROPAGATOR

We will first discuss the structure of the gauge-field propagator within the random phase approximation (i.e. 1-loop order) and will then point out that higher order contributions do not lead to qualitative modifications.

Within RPA, the gauge-field propagator is given by the Feynman diagrams shown in Fig. 9.1, i.e. the polarization insertion Π (or gauge-field self-energy) is calculated only to 1-loop order. In Coulomb gauge the longitudinal component of the vector-field $\mathbf{a}$ vanishes and the polarization tensor $\Pi^{\mu\nu}(q)$ can be expressed in terms of two functions $\Pi^0(q)$ and $\Pi^t(q)$ coupling to the scalar field a_0 and the transverse vector-field $\mathbf{a}$, respectively.

To 1-loop order, the scalar component of the polarisation tensor is given by the density-density correlator of the non-interacting Fermi gas

$$\Pi_0^0(q) \equiv \Pi_0(q) = \int \frac{d^d\mathbf{k}}{(2\pi)^d} \frac{\Theta(\xi_{\mathbf{k}+\mathbf{q}/2}) - \Theta(\xi_{\mathbf{k}-\mathbf{q}/2})}{\omega - (\xi_{\mathbf{k}+\mathbf{q}/2} - \xi_{\mathbf{k}-\mathbf{q}/2}) + i0^+ s(\omega)} \tag{9.7}$$

while the transverse component is given by the transverse current-current correlator

$$\Pi_0^t(q) \;=\; \frac{n_s}{m} \;+\; \int \frac{d^d\mathbf{k}}{(2\pi)^d} \frac{1}{m^2}[\mathbf{k}^2 - (\mathbf{k}\cdot\hat{\mathbf{q}})^2] \frac{\Theta(\xi_{\mathbf{k}+\mathbf{q}/2}) - \Theta(\xi_{\mathbf{k}-\mathbf{q}/2})}{\omega - (\xi_{\mathbf{k}+\mathbf{q}/2} - \xi_{\mathbf{k}-\mathbf{q}/2}) + i0^+ s(\omega)} \tag{9.8}$$

where n_s is the density per spin species. The term n_s/m is due to the tadpole diagram in Fig. 9.1. Note that these are expressions for time-ordered real-frequency quantities. In two dimensions and for small $q = (\omega, \mathbf{q})$ with $|\omega| \ll v_F|\mathbf{q}|$ one obtains [44]

$$\Pi_0^0(q) \sim -\frac{m}{2\pi}\left(1 + i\frac{|\omega|}{v_F|\mathbf{q}|}\right) \tag{9.9}$$

$$\Pi_0^t(q) \sim -i\gamma_0 \frac{|\omega|}{|\mathbf{q}|} + \chi_0 \mathbf{q}^2 \tag{9.10}$$

where $\chi_0 = 1/24\pi m$ is the (Landau) diamagnetic susceptibility of the electron gas (per spin species) and $\gamma_0 = k_F/2\pi$.

Written in a general matrix form, the RPA gauge-field propagator is determined by the equation

$$D^{-1} = D_0^{-1} - \Pi_0 \tag{9.11}$$

which resummes the geometric series in Fig. 9.1. Instead of solving this equation in the $(d+1)$-dimensional basis corresponding to a^μ with $\mu = 0, .., d$, it is easier to decompose the vector part in subspaces corresponding to transverse and longitudinal fluctuations, where the latter vanish due to the Coulomb gauge.

For the half-filled Landau-level one has a 2x2-matrix structure in a space spanned by scalar field a_0 and the transverse vector-field a_t. The RPA equation for D becomes [44]

$$\begin{pmatrix} D^{00} & D^{0t} \\ D^{t0} & D^{tt} \end{pmatrix}^{-1} = \begin{pmatrix} 0 & \frac{i|\mathbf{q}|}{2\pi\tilde{\phi}} \\ -\frac{i|\mathbf{q}|}{2\pi\tilde{\phi}} & \frac{\mathbf{q}^2 v(\mathbf{q})}{(2\pi\tilde{\phi})^2} \end{pmatrix} - \begin{pmatrix} \Pi_0^0(q) & 0 \\ 0 & \Pi_0^t(q) \end{pmatrix} \tag{9.12}$$

where the coupling function $v(\mathbf{q})$ is the Fourier transform of $v(\mathbf{x})$ in (9.4). The off-diagonal elements are due to the Chern-Simons term in the action. Note that the interaction part S_I has been attributed to the "non-interacting" (i.e. Gaussian) gauge-field propagator D_0 by virtue of the constraint linking the electron density operator to $\nabla \times \mathbf{a}$. If $v(\mathbf{x})$ is a Coulomb interaction, one has $v(\mathbf{q}) = 2\pi e^2/\epsilon|\mathbf{q}|$ where ϵ is a dielectric constant.

For small q and $|\omega| \ll v_F|\mathbf{q}|$ the propagator for transverse gauge-field fluctuations has the form

$$D^{tt}(q) \equiv D^t(q) \sim \frac{1}{i\gamma_0|\omega|/|\mathbf{q}| - \chi(\mathbf{q})|\mathbf{q}|^2} \quad \text{where} \quad \chi(\mathbf{q}) = \chi_0 + \frac{1}{2\pi m\tilde{\phi}^2} + \frac{v(\mathbf{q})}{(2\pi\tilde{\phi})^2} \tag{9.13}$$

Obviously $D^t(q)$ is highly singular in the limit $q \to 0$ with $|\omega|/|\mathbf{q}| \to 0$. By contrast, D^{t0} and D^{00} involve extra factors $|\mathbf{q}|$ and $|\omega|/|\mathbf{q}|$, respectively, in the numerator, and are thus comparatively small [44]. If $v(\mathbf{x})$ is a Coulomb interaction, one has $\chi(\mathbf{q}) \sim e^2/2\pi\epsilon\tilde{\phi}^2|\mathbf{q}|$, while in the case of external screening the function $\chi(\mathbf{q})$ tends to a constant $\chi(\mathbf{0}) \equiv \chi$ in the limit $\mathbf{q} \to 0$.

In the spin liquid case only transverse vector fields enter. Hence the gauge-field propagator is purely transverse and given by

$$[D^t(q)]^{-1} = [D_0^t(q)]^{-1} - 2\Pi_0^t(q) \tag{9.14}$$

in RPA. The factor two in front of Π_0^t is due to the spin-degeneracy. In the absence of a pure gauge term one has $[D_0^t(q)]^{-1} = 0$ and thus

$$D^t(q) = -\frac{1}{2\Pi_0^t(q)} \sim \frac{1}{2i\gamma_0|\omega|/|\mathbf{q}| - 2\chi_0|\mathbf{q}|^2} \tag{9.15}$$

where the latter relation holds for small q with $|\omega| \ll v_F|\mathbf{q}|$. Including a gauge term as in (9.6) merely modifies the prefactor of the term proportional to $|\mathbf{q}|^2$ in (9.15) without changing the qualitative asymptotic structure.

To obtain the exact gauge-field propagator, one has to replace the bare bubbles Π_0 in Fig. 9.1 by dressed polarization insertions Π. These insertions can be expanded in

powers of RPA gauge-field propagators. Singular corrections to the bare bubbles can be expected only from scattering processes with small momentum transfers, where the RPA propagator diverges. Now a crucial point is that all these corrections involve fermionic loops with more than two insertions. Hence, the *loop cancellation* theorem derived in Sec. 5 predicts the cancellation of the leading contributions from individual diagrams. Indeed, an explicit evaluation of polarization insertions to 2-loop order by Kim et al. [100] has shown that corrections to the bare bubbles do not change the qualitative small-q behavior (for any ratio $|\omega|/|\mathbf{q}|$) although individual diagrams seem to over-power the 1-loop contributions. For example, two-loop corrections with a purely transverse gauge propagator $D^t(q) \sim [i\gamma|\omega/|\mathbf{q}| - \chi|\mathbf{q}|^2]^{-1}$ yield a total correction [100]

$$\delta \mathrm{Im}\Pi^t(q) \sim \frac{|\omega|}{v_F|\mathbf{q}|} \left[a \, \frac{|\omega|^{2/3}}{\gamma^{1/3}\chi^{2/3}} + b \, \frac{|\omega|}{\chi|\mathbf{q}|} \right] \tag{9.16}$$

for $|\omega| \ll v_F|\mathbf{q}|$ and $|\mathbf{q}| \ll k_F$, where a and b are dimensionless constants. This correction is negligible with respect to the 1-loop result (9.10). The result (9.16) is a sum of self-energy and vertex corrections to the bare bubble. The separate sum over self-energy corrections leads to [100]

$$\delta \mathrm{Im}\Pi_s^t(q) \sim \frac{k_F}{2\pi} \, \frac{|\omega|}{|\mathbf{q}|} \, \frac{(|\omega|/\chi)^{2/3}}{\gamma^{1/3}|\mathbf{q}|} \tag{9.17}$$

which becomes larger than the 1-loop contributions if $|\omega|/|\mathbf{q}|$ vanishes more slowly than $|\omega|^{1/3}$. The leading singularity in (9.17) is obviously cancelled by the vertex corrections.

Several authors have concluded from various arguments that the exact gauge-field propagator has RPA form for small q. The importance of loop cancellation in the gauge problem has been emphasized in particular by Kopietz et al. [62]. Gan and Won [133] and Fröhlich et al. [134] have derived the absence of renormalizations of the RPA propagator from the irrelevance of non-quadratic terms in the effective gauge-field action. Polchinski [135] has derived the stability of the RPA result from a self-consistency argument. He neglected vertex corrections, however, and has therefore been critisized by Kim et al. [100].

In summary, in the limit $q \to 0$ with $|\omega|/|\mathbf{q}| \to 0$ the transverse gauge-field propagator $D^t(q)$ has the singular form

$$D^t(q) = \frac{1}{i\gamma|\omega|/|\mathbf{q}| - \chi|\mathbf{q}|^\zeta} \tag{9.18}$$

where $\zeta = 1$ for half-filled Landau level systems with Coulomb interactions and $\zeta = 2$ for spin liquids and half-filled Landau level systems with short-range interactions. The singularity describes an overdamped mode with a relaxation rate proportional to $|\mathbf{q}|^{1+\zeta}$.

In general the coefficients γ and χ are effective low-energy parameters which include contributions from high-energy fluctuations.

The largest gauge-field fluctuations carry energies $\omega \sim (\chi/\gamma)\,|\mathbf{q}|^{1+\zeta}$. This energy matches with fermionic excitation energies $\xi_{\mathbf{p}+\mathbf{q}/2} - \xi_{\mathbf{p}-\mathbf{q}/2} \sim v_F q_r$ only if $q_r \ll q_t \sim |\mathbf{q}|$, i.e. the dominant scattering processes involve *small momentum transfers which are essentially parallel to the Fermi surface*.[26] Note also that the fermion-gauge-field vertex favors contributions where the transverse field (perpendicular to $\mathbf{q}$) is parallel to the electron current (i.e. normal to the Fermi surface).

9.3. FERMION PROPAGATOR

The fermion (single-particle) propagator G is strongly affected by the gauge-field fluctuations. This is already evident from the behavior of the RPA result for the fermionic self-energy, i.e. the first-order (in D) contribution shown in Fig. 9.2. The singular structure of the propagator D^t for transverse gauge-field fluctuations at small momenta leads to a large decay rate for (putative) quasi-particles. In particular, for $\zeta = 2$ in a two-dimensional system one finds [138, 43]

$$\tau_{\mathbf{k}}^{-1} = |\mathrm{Im}\Sigma(\xi_{\mathbf{k}}, \mathbf{k})| \;\sim\; \xi_0^{1/3}\,|\xi_{\mathbf{k}}|^{2/3} \tag{9.19}$$

where ξ_0 is an energy scale given by $v_F^3/\gamma\chi^2$ (times numerical factors). For general small ξ and $\xi_{\mathbf{k}}$, the self-energy is proportional to [44, 139]

$$\mathrm{Im}\Sigma(\xi, \mathbf{k}) \propto \begin{cases} |\xi|^{2/3} & \text{for} \quad |\xi| > (\chi/\gamma)|\xi_{\mathbf{k}}/v_F|^3 \\ \xi^2/\xi_{\mathbf{k}}^4 & \text{for} \quad |\xi| < (\chi/\gamma)|\xi_{\mathbf{k}}/v_F|^3 \end{cases} \tag{9.20}$$

For $\zeta = 1$ the RPA self-energy is proportional to $\xi\,\log(\epsilon_F/|\xi|)$ [63]. In any case the self-energy correction modifies the non-interacting fermion propagator dramatically, i.e. the low-energy structure of the bare fermionic excitations is completely destroyed. Halperin et al. [44] have proposed to replace the bare fermions by renormalized quasi-particles with a dispersion relation $\xi_{\mathbf{k}}^* \propto s(k_r)|k_r|^{3/2}$ for $\zeta = 2$ and by $\xi_{\mathbf{k}}^* \propto k_r|\log|k_r||$ in the case $\zeta = 1$.

The strong modification of the fermion propagator by the RPA self-energy clearly calls for a controlled treatment of higher order corrections. This problem and especially the choice of an adequate method has been discussed somewhat controversially in the literature, but the results for G found by different authors do not differ very much after all. We will first derive a non-perturbative result for G along the lines of Sec. 5, i.e. a resummation exploiting the smallness of relevant momentum transfers, and will also discuss

[26] Recall from Sec. 5 that $q_r = \mathbf{q}\cdot\hat{\mathbf{p}}$ and $q_t = \sqrt{\mathbf{q}^2 - q_r^2}$ are the radial and tangential components of $\mathbf{q}$, respectively, referred to a momentum $\mathbf{p}$ close to the Fermi surface.

possible flaws of this approach in the gauge problem. We will then address other methods, especially the large-N expansion and scaling (i.e. renormalization group) theories.

We now extend results from Sec. 5, that had been derived for systems with purely fermionic degrees of freedom, to the gauge theory. Asymptotic Ward identities [35, 63] and bosonization [140, 141] yield the same result for G. Here we focus on the Ward identity approach since it allows for a more direct comparison with the other diagrammatic methods. Since spin does not affect the logical structure of the derivation, we consider spinless fermions for simplicity.

The irreducible current vertex in the gauge theory is defined by[27]

$$\Lambda^\mu(p;q) = -\langle j^\mu(q)\, \psi_{p-q/2}\, \psi^*_{p+q/2}\rangle^{irr}_{tr} \tag{9.21}$$

with a current operator

$$\begin{array}{rcl} j^0(q) & \equiv & \rho(q) = \int_k \psi^*_{k-q/2}\psi_{k+q/2} \\ \mathbf{j}(q) & = & \int_k \mathbf{v_k}\, \psi^*_{k-q/2}\, \psi_{k+q/2} + \frac{1}{m}\int_{q'} \rho(q')\, \mathbf{a}(q-q') \end{array} \tag{9.22}$$

The labels "tr" and "irr" indicate truncation of external fermion legs and irreducibility of Feynman diagrams with respect to cutting a single gauge-field propagator, respectively. Charge conservation (or gauge invariance) implies the well-known exact Ward identity

$$q_\mu \Lambda^\mu(p;q) = G^{-1}(p+q/2) - G^{-1}(p-q/2) \tag{9.23}$$

As discussed in detail in Sec. 5.2, the dominance of forward scattering suggests the asymptotic *density-current relation*

$$\mathbf{\Lambda}(p;q) \sim \mathbf{v_p} \Lambda^0(p;q) \tag{9.24}$$

to leading order in a small momentum transfer (or scattering angle) expansion. Note that the $\mathbf{a}$-dependent part of the current operator does not enter here, because the expectation value $\langle \mathbf{a}\, \rho\, \psi\psi^*\rangle$ vanishes. Combining (9.23) and (9.24) leads to

$$\Lambda^\mu(p;q) = \frac{G^{-1}(p+q/2) - G^{-1}(p-q/2)}{\omega - \mathbf{v_p}\cdot\mathbf{q}}\, \lambda^\mu(p) \tag{9.25}$$

where $\lambda^\mu(p) = (1, \mathbf{v_p})$ is the bare current vertex. The dressed current vertex Λ^μ is thus a simple functional of the propagator G.

The fermionic self-energy Σ is related to G, $D^{\mu\nu}$ and Λ^μ by the exact Dyson equation

$$\Sigma(p) = i \int_{p'} G(p')\, \lambda_\mu[(p+p')/2]\, D^{\mu\nu}(q)\, \Lambda_\nu[(p+p')/2; p'-p] \tag{9.26}$$

[27] Note that we work with a real time/frequency representation here, while in Sec. 5 a euclidean representation prevailed.

which is illlustrated diagramatically in Fig. 9.3. Inserting Λ^μ from (9.25) and using the relation $G^{-1} = G_0^{-1} - \Sigma$ yields a linear integral equation for G,

$$(\xi - \xi_{\mathbf{p}})\, G(p) = 1 + \int_{p'} \frac{i D_{\mathbf{p}}(p-p')}{\xi - \xi' - \mathbf{v}_{(\mathbf{p}+\mathbf{p}')/2} \cdot (\mathbf{p}-\mathbf{p}')}\, G(p') \tag{9.27}$$

with a contracted gauge-field propagator

$$D_{\mathbf{p}}(q) = \lambda_\mu(p)\, D^{\mu\nu}(q)\, \lambda_\nu(p) \tag{9.28}$$

This integral equation is the analogon of (5.17) for fermions coupled to a gauge-field.

For purely *transverse* gauge field fluctuations (which yield the most singular contributions) the gauge-field propagator can be written as

$$D^{jj'}(q) = \Big(\delta_{jj'} - \frac{q_j q_{j'}}{|\mathbf{q}|^2}\Big) D^t(q) \quad \text{for} \quad j, j' = 1, ..., d \tag{9.29}$$

while $D^{0\mu} = D^{\mu 0} = 0$. In this case one has

$$D_{\mathbf{p}}(q) = \mathbf{v}_{\mathbf{p}}^2\, \frac{q_t^2}{|\mathbf{q}|^2}\, D^t(q) \tag{9.30}$$

where q_t is the tangential component of $\mathbf{q}$ (with respect to the Fermi surface near $\mathbf{p}$, see Sec. 5). Decomposing the momenta in (9.27) in radial and tangential components as in Sec. 5, one obtains an effective one-dimensional equation of motion of the form (5.25), with a q_t-averaged gauge-field propagator [63]

$$\bar{D}(\omega, q_r) = \frac{S_{d-1}}{(2\pi)^{d-1}}\, v_F^2 \int dq_t\, \frac{q_t^d}{|\mathbf{q}|^2}\, D^t(q) \tag{9.31}$$

The general formal solution of this equation of motion is given by (5.28)-(5.33). Exactly the same formal expression for the fermionic propagator in the gauge theory has been derived by Haldane-bosonization [140, 141], and the equivalence with the Ward identity approach has been pointed out [38, 140]. A similar expression has been obtained within the eikonal approximation [64], which also exploits the smallness of relevant momentum transfers. For fermions with spin the solution has precisely the same form, i.e. spin enters only indirectly via D^t.

We now discuss explicit results following from the above formal solution for G, focussing mainly on the case with the strongest singularities, i.e. for $\zeta = 2$ in two dimensions (relevant for the half-filled Landau level with screened interactions and the spin liquid model). In this case the q_t-averaged gauge-field propagator has the simple asymptotic low-energy form [63]

$$\bar{D}(\omega, q_r) \propto |\omega|^{-1/3} \tag{9.32}$$

Since the singularity is dominated by momenta with $q_r \ll q_t$, the q_r-dependence in $\bar{D}$ is irrelevant. Note that $\bar{D}(\omega, q_r)$ does not depend on any momentum transfer cutoff, since for $\zeta = 2$ the integrand in (9.31) decays rapidly as a function of q_t.

An explicit result for $G(t,r) = \int_{\xi,p_r} G(\xi,p_r)\, e^{ip_r r - i\xi t}$ has been reported by Ioffe et al. [63]:

$$G(t,r) \sim \frac{1}{2\pi}\, \frac{1}{r - v_F t}\, \exp\left[\frac{-|r|}{r_0^{1/3}[|r| - v_F s(r) t]^{2/3}}\right] \tag{9.33}$$

where $s(.)$ is the sign-function and r_0 is a length scale of order $\gamma\chi^2/v_F^2$. Obviously the fermionic density of states $N(\xi)$, given by the imaginary part of the Fourier transform of $G(t,0)$, is not affected by the gauge-field. However, the locus of a particle is smeared over a distance $r \propto t^{2/3}$, the probability to find it along the classical path $r = v_F t$ decays exponentially, and the velocity of a wave packet vanishes at $t \to \infty$ [63]. Hence, a single fermion cannot propagate in the presence of transverse gauge-fields (while a particle-hole pair with a small relative momentum can). The equal-time correlator

$$G(0,r) \sim \frac{1}{2\pi r}\, e^{-|r/r_0|^{1/3}} \tag{9.34}$$

decays exponentially, i.e. the momentum distribution function is analytic at the Fermi surface. This latter result has also been derived via Haldane-bosonization [141]. An analytic momentum distribution function has also been found by Khveshchenko and Stamp [64] within their eikonal method.

Altshuler et al. [63] have also presented an explicit result for $G(\xi, \mathbf{p})$ in the low-energy limit, with a behavior corresponding to overdamped fermions with a characteristic energy that scales as $|p_r|^{3/2}$. In particular, they have found the asymptotic behavior

$$G(\xi,\mathbf{p}) \propto \begin{cases} -1/v_F p_r & \text{for} \quad \xi_0^{1/3}|\xi|^{2/3} \ll v_F|p_r| \\ 1/\xi_0^{1/3}(i\xi)^{2/3} & \text{for} \quad \xi_0^{1/3}|\xi|^{2/3} \gg v_F|p_r| \end{cases} \tag{9.35}$$

which is equivalent to the RPA result. By contrast, the eikonal method seems to yield an exponential decay of $G(\xi, \mathbf{p}_F)$ for $\xi \to 0$ [64].

Kopietz [141] has derived explicit results via bosonization for the equal-time propagator $G(0,r)$ in the case $\zeta = 2$ for general dimensionality d. He obtained Fermi liquid behavior $G(0,r) \propto 1/r$ for all $d > 3$ (with large subleading corrections for $d < 6$), anomalous power-law decay $G(0,r) \propto 1/|r|^{1+\eta}$ with a non-universal exponent η in $d = 3$, and exponential decay

$$G(0,r) \sim \frac{1}{2\pi r}\, e^{-|r/r_0|^{1-d/3}} \tag{9.36}$$

in dimensions $d < 3$. This identification of critical dimensions and the anomalous power-law in $d = 3$ is consistent with earlier renormalization group results by Gan and Wong [133].

Kwon et al. [140] have analyzed the case $\zeta = 1$ in two dimensions within bosonization. They find "marginal Fermi liquid behavior" with a linear energy dependence of the quasi-particle decay rate, and an anomalous power-law for the equal-time propagator $G(0, r)$. We note that for $\zeta = 1$ the q_t-average in (9.31) does not converge at the upper end, i.e. the original assumption that momentum transfers be small is not respected by the solution, as long as no explicit cutoff is introduced.

The validity of results obtained by resumming forward scattering processes in the gauge theories via asymptotic Ward identities or bosonization has been questioned by several authors. A peculiar feature that distinguishes the gauge theories from other forward scattering dominated systems is that the most important processes are characterized by (small) momentum transfers which are essentially parallel to the Fermi surface, i.e. $q_t \gg q_r$. This makes the linearization of fermionic energies such as $\xi_{\mathbf{k}_F+\mathbf{q}} \sim v_F q_r$ used in the resummation techniques problematic, since the neglected correction of order $\mathbf{q}^2 = q_r^2 + q_t^2$ may become comparable to (or even larger than) the linear term $v_F q_r$. For example, corrections to the RPA self-energy in the case $\zeta = 2$ in two dimensions pick up major contributions from energy-momentum transfers with $\omega \sim q_t^3 \sim q_r^{3/2}$, such that the transverse correction $q_t^2/2m$ to the linearized energy $v_F q_r$ is not negligible [135]. On the other hand one should note that these corrections often cancel, such as for example in the symmetrized expression for particle-hole excitation energies $\xi_{\mathbf{k}+\mathbf{q}/2} - \xi_{\mathbf{k}-\mathbf{q}/2} = \mathbf{v}_\mathbf{k} \cdot \mathbf{q}$. This may explain why these corrections do not seem to overpower the loop cancellation property derived in Sec. 5, as confirmed for the gauge theories by the explicit two-loop calculation of Kim et al. [100].

To equip the problem with a suitable expansion parameter, Polchinski [135] and Ioffe et al. [63] have generalized the above gauge theories (including S_a, eq. (9.6)) by allowing for N fermion flavors with arbitrary N. The latter group rescales terms in the bare action such that the RPA gauge-field propagator has the form (9.18) with $\gamma \mapsto N\gamma$ and $\chi \mapsto \chi/N^{1/2}$. The energy scale $\xi_0 \propto v_F^3/\gamma\chi^2$ remains thus independent of N.

According to Ioffe et al. [63] the results obtained by resummation techniques which do not treat tangential momentum transfers correctly are reliable only in the limit $N \to 0$, because in this limit momentum transfers which are almost parallel to the Fermi surface are no longer favorable. Based on estimates of general vertex corrections, Altshuler et al. [63] conclude that the density-current relation (9.24) has important corrections of order $N^{1/2}$ as soon as N differs from zero. They find that the current vertex $\mathbf{\Lambda}$ has a transverse component $\mathbf{\Lambda}_t$ given by

$$\mathbf{\Lambda}_t(p;q) \sim \mathbf{q}_t \, \frac{v_F q_r}{2q_t^2} \left[\sqrt{1 - 2\alpha|\omega|q_t/q_r^2 + i0^+} - 1\right] \Lambda_r(p;q) \tag{9.37}$$

for small N and q_t, where $\alpha = N^{1/2}/(2\pi)^2\gamma\chi$ and $\Lambda_r(p;q) = \hat{\mathbf{p}}\cdot\mathbf{\Lambda}(p;q) \sim v_F\Lambda^0(p;q)$ is the

radial component of $\mathbf{\Lambda}$. Such a correction smears the pole in $\omega = v_F q_r$ in (9.25), which leads to an intrinsic cutoff for exponential singularities such as in the result (9.33) for G.

Another solvable limit is reached for $N \to \infty$ [63, 135]. For large N the vertex corrections in the Dyson equation (9.26) are suppressed by powers of $1/N$, i.e. a self-consistent random phase approximation is exact for $N \to \infty$. Due to its weak momentum dependence, the plain RPA self-energy (calculated from G_0) solves also the self-consistent equation [63, 135]. The propagator $G(\xi, \mathbf{p})$ obtained by this large-N expansion has several features in common with the one obtained for $N \to 0$. These common features hold plausibly for any N. In particular, for $\zeta = 2$ in two dimensions one recovers the low-energy behavior (9.35), with the same characteristic energy scale $\xi \propto |p_r|^{3/2}$. Discrepancies appear for the precise form of the propagator for $\xi \sim |v_F p_r|^{3/2}/\xi_0^{1/2}$. The extrapolation of results obtained in the large-N limit to the physical values $N = 1$ or 2 is of course problematic, especially since the importance of vertex corrections at finite N has been clearly demonstrated in the two-loop calculation by Kim et al. [100].

In addition to the resummation techniques discussed so far, scaling and renormalization group approaches can provide valuable insight into the low-energy structure of Fermi systems coupled to transverse gauge-fields. By a renormalization group analysis of the case $\zeta = 2$, Gan and Wong [133] have identified $d = 3$ as the critical dimension separating Fermi from non-Fermi liquid behavior [133]. At the critical dimension they find that the quasi-particle weight vanishes as ω^η, where η is a non-universal exponent. Chakravarty et al. [142] have constructed an expansion in $\epsilon = 3-d$. In constrast to Gan and Wong these authors obtain an anomalous exponent of order ϵ for $d < 3$, and only logarithmic corrections to Fermi liquid behavior in $d = 3$. Nayak and Wilczek [136] have analyzed the action for the half-filled Landau level (in two dimensions) with a generalized Coulomb interaction defined by a coupling function $v(\mathbf{q}) \propto 1/|\mathbf{q}|^\gamma$. Introducing deviations $1-\gamma$ from the Coulomb case $\gamma = 1$ as a small control parameter, they were able to treat the infrared singularities of the system by a systematic renormalization group procedure. A scenario similar to that of equilibrium critical phenomena near four dimensions was found. The case of Coulomb interactions is characterized by marginally irrelevant (i.e. logarithmic) corrections to Fermi liquid theory [136]. This is consistent with the results from the $1/N$-expansion [63], but inconsistent with the anomalous power-law behavior of the equal-time propagator found by Kwon et al. [140]. If the interactions are shorter-ranged (i.e. $\gamma < 1$), a non-trivial fixed point controls anomalous infrared power-laws, analogous to the Wilson-Fisher fixed point in $4 - \epsilon$ dimensions [136]. In particular, the quasi-particle weight vanishes as $Z \propto \omega^\eta$ and the effective Fermi velocity as $v_F^* \propto \omega^{|\eta_{v_F}|}$, where the exponents are given by $\eta = -\eta_{v_F} = (1-\gamma)/2 + \mathcal{O}[(1-\gamma)^2]$. Whereever comparable, these results

are consistent with those obtained by most other groups. It would be interesting to see whether the full momentum and energy dependence of $G(\xi, \mathbf{p})$ can be calculated within this expansion. For short-range interactions ($\gamma = 0$) the expansion parameter $1-\gamma$ is of course not small, but one may hope (as in the theory of critical phenomena) that the qualitative infrared structure does not change on the way from $\gamma \sim 1$ to $\gamma = 0$.

We finally note that the single-fermion propagator G is not a gauge-invariant quantitiy. Here we will not address the question of how observable physical properties may be related to it. A short discussion of that problem is given for example by Kim et al. [100]. In any case the fermion propagator at fixed gauge is an important theoretical quantity and its behavior has been the main issue in most theoretical works on the coupled fermion-gauge-field system.

9.4. RESPONSE FUNCTIONS

The response of the coupled Fermi gauge-field system is described by the (dressed) polarization tensor $\Pi^{\mu\nu}$. For low frequencies and long wavelengths ($\mathbf{q}$ small) we have already seen in 9.2 that $\Pi^{\mu\nu}$ is not drastically affected by the gauge-field fluctuations. The leading low-energy small-$\mathbf{q}$ response behaves therefore in a Fermi liquid fashion. This is in fact experimentally observed in the case of half-filled Landau level systems (for references, see Kim et al. [100]). A comparison with the non-Fermi liquid properties of high-T_c superconductors is not direct since the holons are not included in the spinon-gauge-field action (9.5).

Pronounced deviations from Fermi liquid behavior appear for response functions with momenta near $2k_F$. In particular, in systems with $\zeta = 2$ the density-density correlators acquire anomalous power-law behavior at $2k_F$, and interaction vertices Γ_{2k_F} with external fields at momenta $2k_F$ are strongly enhanced [63, 135]. Non-Fermi liquid results for $2k_F$-functions in the case $\zeta = 1$ have also been reported [63, 140].

As an example, let us compare the results obtained for Γ_{2k_F} in systems with $\zeta = 2$ by different methods. Neither the leading asymptotic Ward identity (9.25) (equivalent to bosonization) nor the omission of vertex corrections seems to provide the correct result for systems with finite N. The former yields an exponential divergence $\Gamma_{2k_F}(\omega) \propto e^{|\omega_0/\omega|^{1/3}}$, where ω is supposed to be the biggest of all energy variables of the vertex function, and ω_0 is a system dependent energy scale [63]. Neglecting vertex corrections, one finds a function $\Gamma_{2k_F}(\omega)$ that tends to a constant in the low-energy limit [63]. This latter result is exact in the limit $N \to \infty$. However, including vertex corrections via $(1/N)$-corrections, Altshuler et al. [63] have obtained a power-law

$$\Gamma_{2k_F}(\omega) \propto \omega^{-\sigma} \tag{9.38}$$

with an exponent

$$\sigma = \frac{1}{2N} + \frac{\log^3(N)}{2\pi^2 N^2} + \mathcal{O}(N^{-2}) \quad \text{for} \quad N \gg 1 \tag{9.39}$$

A power-law is also obtained by modifying the simple density-current relation (9.24) to (9.37) by including small-N corrections of order $N^{1/2}$. This yields an exponent [63]

$$\sigma = \frac{16\sqrt{2}}{9\pi} \frac{1}{N^{1/2}} + \mathcal{O}(1) \quad \text{for} \quad N \ll 1 \tag{9.40}$$

Since a power-law is obtained for both small and large N, one may be confident that $\Gamma_{2k_F}(\omega)$ obeys a power-law for any finite N.

10. CONCLUSIONS

We have reviewed the low-energy structure of d-dimensional Fermi systems with short-range or long-range interactions, and also of Fermi systems coupled to a gauge-field. The analysis was restricted to pure systems and normal (i.e. not symmetry-broken) phases. As in Fermi liquid theory, the low-energy physics was assumed to be governed by excitations close to a (Fermi) surface in momentum space. Depending on dimensionality and the nature of the interactions, the systems belong to distinct "universality classes": conventional Fermi liquids, unconventional Fermi liquids with enhanced subleading corrections, and various types of non-Fermi liquids such as the one-dimensional Luttinger liquid. The main-focus of this article were non-trivial interaction effects due to (residual) interactions in the *forward scattering* channel, i.e. with small momentum transfers. Without leading to spontanous symmetry breaking or dynamical gap generation, these interactions affect the qualitative low-energy behavior of any Fermi system.

The *renormalization group* à la Wilson provides a well-defined link between microscopic systems and effective low-energy theories (see Sec. 2). The central concept here is a family of effective low-energy actions $\bar{S}^{\Lambda}$, defined on a thin momentum shell of width Λ around the Fermi surface, and (in principle) uniquely determined by the microscopic theory via integration over high energy degrees of freedom. In practice this integration can be carried out only for weak coupling, but our experience with specific real systems and some exactly solvable models shows that the qualitative structure of effective actions emerging from a weak coupling analysis often applies to strong coupling systems, too.

Conservation laws highly constrain the low-energy behavior of a Fermi system. The continuity equations for conserved currents give rise to *Ward identities* which relate different correlation, response and vertex functions. In particular, charge and spin conservation guarantee that the field renormalization Z cancels from the low-energy long-wavelength response functions. This cancellation, which is crucial for Landau's theory of the response of a Fermi liquid to external perturbations [2], holds in non-Fermi liquids, too (see Sec. 3).

All Fermi systems seem to share the common feature that residual interactions with finite momentum transfers (i.e. of order k_F) either drive an instability towards symmetry breaking[28] or do not play any prominant role at all. On the other hand, residual

[28]In one dimension one may rather have dynamical gap-generation without symmetry breaking, since

interactions in the forward scattering channel modify at least the leading low-energy long-wavelength response in normal systems (for example, Landau parameters renormalize the compressibility and magnetic susceptibility). In this sense the low-energy theory of interacting Fermi systems is never asymptotically free, but contains at least marginal interactions to deal with. Fortunately (for the theorist), these residual interactions, which are usually not weak in real systems, are easier to treat than bare interactions with arbitrary momentum transfers in a microscopic theory. There are the following two sources of simplifications.

First, the reduced phase space for virtual excitations in the effective low-energy theory, which is restricted to the thin Λ-shell around the Fermi surface, leads to a suppression of most virtual processes in the perturbation expansion. Real decay or scattering processes, such as the decay of a quasi particle via a single particle-hole excitation, are also restricted by an intrinsic energy cutoff set by the energy of the excitations (Landau's argument). If the dimensionality of the system is not too low and the interactions are not too singular, only a very restricted set of Feynman diagrams determines the leading and sometimes even the subleading low-energy behavior in terms of effective low-energy parameters (e.g. Landau parameters).[29]

Second, forward scattering processes obey a special conservation law: The velocity of each scattering particle is conserved in the process. Equivalently one can say that charge and spin are not only conserved globally, but even locally in arbitrarily small sectors in momentum space. This conservation law holds only asymptotically in the sense that it is exact only in the forward scattering limit $\mathbf{q} \to 0$. Only in special models such as the one-dimensional Luttinger model the velocity conservation is exact even for finite $\mathbf{q}$. As shown in detail in Sec. 5, this asymptotic conservation law gives rise to several simplifications, which become particularly important in systems where the phase-space reduction in the low-energy theory is compensated by singular interactions or low dimensionality. One of these simplifications is *loop cancellation*, i.e. Feynman diagrams containing fermionic loops with more than two insertions cancel each other or at least leading singularities of single diagrams cancel. This implies in particular that the polarization insertion $\Pi(q)$ is not dressed by forward scattering processes for small $\mathbf{q}$. Consequently the random phase approximation describes the effect of forward scattering processes on the leading low-energy long-wavelength response not only in Fermi liquids, but also in systems where quasi-particle excitations are destroyed by forward scattering. Furthermore, the asymptotic velocity conservation yields an asymptotic *density-current relation* between the ir-

the latter is prevented by strong fluctuations

[29]We emphasize that this has been explained already without extensive use of RG machinary some time ago in the literature, see for example Serene and Rainer [77].

reducible density vertex and the irreducible current vertex, i.e. $\mathbf{\Lambda}(p;q) \sim \mathbf{v_p}\Lambda^0(p;q)$. Combining this relation with the exact Ward identity reflecting total charge (or spin) conservation, leads to an *asymptotic Ward identity* that expresses $\Lambda^0(p;q)$ and $\mathbf{\Lambda}(p;q)$ uniquely in terms of the single-particle propagator G. Plugging this into a Dyson equation, we have derived an expression for G which sums the dressing by forward scattering processes to all orders in the coupling constant. An alternative way of exploiting the asymptotic conservation laws obeyed by forward scattering is d-dimensional *bosonization* as proposed by Haldane [68], which leads to the same results.

In a *Fermi liquid* (Sec. 4) residual interactions (within a thin Λ-shell in momentum space) do not affect the leading asymptotic behavior of the single-particle propagator in the low-energy limit. In particular, the wave function renormalization Z^Λ has a finite limit $Z > 0$ for $\Lambda \to 0$. To leading order in a low-energy expansion, the propagator has the form $G(p) = Z/(\xi - v_F^* p_r)$. This implies a discontinuity Z in the momentum distribution function, and the existence of fermionic single-particle excitations, Landau's *quasi-particles*, with a linear dispersion relation and velocity v_F^*. For the single-particle properties residual interactions are important only at next-to-leading order in a low-energy expansion, where they yield a damping of quasi-particle excitations and contribute to a smooth background of incoherent excitations. By contrast, residual interactions contribute already in leading order to low-energy response functions in a Fermi liquid. In particular, the low-energy long-wavelength (small $\mathbf{q}$) charge- or spin-density response is described exactly by an RPA summation with the Landau function $f_{\mathbf{k}_F\mathbf{k}'_F} = \lim_{\Lambda\to 0} \bar{g}^\Lambda_{\mathbf{k}_F\mathbf{k}'_F}(\mathbf{0})$ as interaction (where $\bar{g}^\Lambda_{\mathbf{k}\mathbf{k}'}(\mathbf{q})$ is the renormalized running coupling function). These residual interactions renormalize the compressibility and the spin susceptibility by a finite factor.

In a *one-dimensional Luttinger liquid* (Sec. 6) residual interactions in the forward scattering channel do not only affect response functions, but also the leading low-energy behavior of the single-particle propagator. The wave function renormalization factor Z^Λ vanishes for $\Lambda \to 0$. The propagator obeys an anomalous scaling law $G(sp) = s^{\eta-1}G(p)$ with a non-universal (i.e. system-dependent) anomalous exponent η. As a consequence, the momentum distribution has a continuous power-law behavior $n_\mathbf{k}$–$n_{\mathbf{k}_F} \propto |k_r|^\eta$ near the Fermi surface, and the density of single-particle excitations $N(\xi)$ vanishes as $|\xi|^\eta$ at low energies. In addition to properties related to anomalous scaling, the strong coupling of fermions to collective modes in the Luttinger liquid leads also to "spin-charge separation": an extra fermion inserted into the system decays in collective charge and spin fluctuations which propagate with two different velocities and thus separate. Although residual forward scattering destroys Landau quasi-particles, the low-energy, long-wavelength response of a Luttinger liquid has the same structure as for a Fermi liquid: an RPA sum of bubble

chains with fixed point couplings $f_{\mathbf{k}_F\mathbf{k}'_F} = \lim_{\Lambda\to 0} \bar{g}^{\Lambda}_{\mathbf{k}_F\mathbf{k}'_F}(\mathbf{0})$ as interactions. In contrast to the situation in a Fermi liquid, the β-function for $\bar{g}^{\Lambda}_{\mathbf{k}_F\mathbf{k}'_F}(\mathbf{0})$ does not vanish simply as a consequence of reduced phase space for $\Lambda \to 0$, but due to cancellations imposed by the special conservation laws obeyed by the forward scattering processes. Singular self-energy and vertex corrections cancel in the polarization bubble. The charge and spin density response functions have poles in $\omega = u_c|\mathbf{q}|$ and $\omega = u_s|\mathbf{q}|$, respectively, which describe gapless bosonic collective excitations with a linear energy-momentum relation.

In Sec. 7 we have presented a quantitative analysis of the effects of residual interactions in d-dimensional systems with *short-range* interactions, clarifying in particular how Luttinger liquid behavior in one dimension crosses over to Fermi liquid behavior in higher dimensions as a function of continuous dimensionality. It turned out that a normal Fermi system with short-range interactions has two distinct characteristic dimensions where the low energy behavior undergoes significant changes. Below two dimensions, forward scattering of single-particle excitations by long-wavelength spin- and charge-density fluctuations yields the dominant contribution to the self-energy, and makes the quasi-particle decay-rate scale as $|k_r|^d$ instead of the square law valid for $d > 2$. These scattering corrections do not, however, destroy Fermi liquid quasi-particle behavior until the *critical dimension* $d_c = 1$ is reached, where small-$\mathbf{q}$ scattering completely destroys the quasi-particle pole in the propagator. In Fig. 10.1 we show a schematic phase diagram of Fermi systems with short-range interactions in the dimensionality-coupling plane, where g is a typical coupling in the forward scattering channel (e.g. a certain harmonic of the Landau function). For strong effective couplings $|g| > g_c$ there may be Pomerantchuk instabilities, i.e. strong coupling instabilities such as phase separation, signalled by an infinite compressibility, or ferromagnetism, indicated by a divergent spin susceptibility. At weak coupling in two dimensions, the scenario in Fig. 10.1 is compatible with a rigorous result by Feldman et al. [17], who proved that in the absence of Cooper processes the perturbation expansion with respect to short-range interactions has a finite radius of convergence. Other instabilities (than Cooper or forward scattering driven), such as antiferromagnetism or charge density wave instabilities, may only occur at sufficiently strong coupling or for very special Fermi surfaces (with nesting).

Fermi systems with a pair-potential $V(\mathbf{r}) \propto 1/|\mathbf{r}|^{d-\gamma}$ as a prototype for *long-range* density-density interactions have been reviewed in Sec. 8. For such interactions there is a critical dimension $d_c = 1 + \gamma/2 > 1$ separating Fermi liquid from non-Fermi liquid behavior. In spite of screening effects, for $d \leq d_c$ the forward scattering is so strong that quasi-particles cannot exist. In the marginal case $d = d_c$ the momentum distribution function obeys a power-law behavior as in a Luttinger liquid.

In Sec. 9 we have reviewed the low-energy physics of coupled fermion-gauge-field systems, which have become important in the theory of the half-filled Landau level and also in attempts to describe the "strange" normal metallic phase of high-T_c superconductors. Transverse gauge-fields are not screened and generate forward scattering amplitudes in the Fermi system. As a consequence of loop cancellation, the gauge-field propagator and also the low-energy long-wavelength response of the coupled system can be calculated in random phase approximation. Reliable calculations of the detailed behavior of the fermion propagator $G(\xi, \mathbf{p})$ seem to be difficult, because vertex corrections are not yet fully under control. A peculiar feature that distinguishes the gauge theories from other forward scattering dominated systems is that the most important processes are characterized by (small) momentum transfers which are essentially parallel to the Fermi surface. This makes the linearization of fermionic energies used in resummation techniques such as asymptotic Ward identities or bosonization problematic. As a consequence, there is a broad consensus only on those properties of the fermion propagator which seem to be insensitive to the different assumptions made for the vertex corrections. In particular, it is clear that Landau quasi-particles are destroyed by the gauge-field fluctuations (in physically relevant dimensions), and are replaced by overdamped fermionic excitations with a non-linear dispersion relation. While the long-wavelength response of the coupled fermion-gauge-field system behaves as in a Fermi liquid at low energies, the $2k_F$-response seems to obey anomalous power-laws.

Let us conclude with a few remarks on the normal phase of high temperature superconductors. The research reviewed in this work has shown that a generic normal Fermi system with short-range or Coulomb interactions in dimensions $d > 1$ obeys Fermi liquid behavior, with no weak coupling instabilities besides the Cooper instability. Here "generic" means that the Fermi surface should not have a special shape, and the Fermi velocity should be finite. Singular long-range interactions (stronger that Coulomb) such as in the gauge-theories can destroy Fermi liquid behavior in two and three dimensions, and have therefore been invoked to explain the observed non-Fermi liquid properties of high-T_c superconductors [39, 40, 41]. Nested Fermi surfaces [26] and strong van Hove singularities in the density of states due to peculiarities of the bandstructure [27, 28] have also been argued to be responsible for anomalous low-energy behavior. These suggestions take support from photoemission experiments which indicate extended pieces of the Fermi surface with very small curvature and regions with "flat" energy-bands in the Brillouin zone [6]. Other proposals identify the proximity to an instability (quantum critical point) of antiferromagnetic [48, 49] and/or charge [50, 51, 52] origin as a source of singular scattering, leading to a doping dependent disruption of Fermi liquid behavior. Also these ideas

are (more or less) motivated by experimental evidences. All the proposals (gauge-theory, nesting or Van-Hove scenario, and quantum criticality) involve complicated infrared singularities and strong coupling problems, which have so far prevented the construction of a complete and generally accepted theory.

Acknowledgements

We are very grateful for valuable discussions with P.W. Anderson, N. Andrei, G. Gallavotti, M. Grilli, H. Fukuyama, D. Haldane, R. Hlubina, L.B. Ioffe, D. Khveshchenko, H. Knörrer, P. Kopietz, B. Marston, V. Meden, E. Müller-Hartmann, M. Randeria, D. Rainer, T.M. Rice, M. Salmhofer, K. Schönhammer, E. Trubowitz, and D. Vollhardt. During the course of this work, we have benefitted from the hospitality of the International Center for Theoretical Physics (Trieste), the Institute for Scientific Interchange (Torino) and the Aspen Center for Physics. W.M. was partially supported by the Sonderforschungsbereich No. 341 of the Deutsche Forschungsgemeinschaft. C.C. and C.D. have been supported within the "progetto ricerca avanzata istituto nazionale fisica della materia (INFM)".

APPENDICES

Appendix A: Loop-cancellation for N = 3:

Here we demonstrate the loop-cancellation explicitly for the case $N = 3$. There are two ways of adding an insertion with momentum $q_3 = q$ to a loop with two insertions carrying momenta q_1 and q_2, as shown in Fig. A.1. Using the relation (5.18), the two Feynman diagrams yield

$$\int_k \frac{h(\mathbf{k},\mathbf{k},\mathbf{k};q_1,q_2,q)}{iq_0 - \mathbf{v_k}\cdot\mathbf{q}} [G_0(k-q/2+q_1/2) - G_0(k+q/2+q_1/2)]G_0(k-q/2-q_1/2) +$$

$$\int_k \frac{h(\mathbf{k},\mathbf{k},\mathbf{k};q_1,q_2,q)}{iq_0 - \mathbf{v_k}\cdot\mathbf{q}} [G_0(k-q/2-q_1/2) - G_0(k+q/2-q_1/2)]G_0(k+q/2+q_1/2)$$

to leading order in $\mathbf{q}_\nu$. Note that $q_1+q_2+q = 0$ (momentum conservation). The above expression yields contains four terms. The third term obviously cancels the second one in the above expression. Shifting the integration variable k by $\pm q/2$ and using $\mathbf{v}_{\mathbf{k}\pm\mathbf{q}/2} \approx \mathbf{v_k}$ and $h(\mathbf{k}\pm\mathbf{q}/2,\ldots;q_1,\ldots) \approx h(\mathbf{k},\ldots;q_1,\ldots)$, one finds that the two remaining contributions to the integral cancel each other to leading order in $\mathbf{q}$.

Appendix B: Spectral representation of $\bar{D}$

The effective interaction has a spectral representation of the form

$$D_{\mathbf{p}}(\omega,\mathbf{q}) = \int_{-\infty}^{\infty} d\omega' \frac{\Delta_{\mathbf{p}}(q_0',\mathbf{q})}{\omega - \omega' + i0^+ s(\omega)} \; + \; D_{\mathbf{p}}(\mathbf{q}) \qquad (B.1)$$

where $\Delta_{\mathbf{p}}(\omega,\mathbf{q}) = -\pi^{-1}s(\omega)\mathrm{Im}D_{\mathbf{p}}(\omega,\mathbf{q})$ and $D_{\mathbf{p}}(\mathbf{q})$ is a real function of $\mathbf{q}$. Performing the angular average, one has a spectral representation for $\bar{D}$ (cutoff-dependence not written)

$$\bar{D}(\omega,q_r) = \int_{-\infty}^{\infty} d\omega' \frac{\bar{\Delta}(\omega',q_r)}{\omega - \omega' + i0^+ s(\omega)} \; + \; \bar{D}(q_r) \qquad (B.2)$$

Inserting this in our expression for $L(\omega,q_r)$ and performing the energy-time Fourier transformation by doing a simple contour integral, one finds

$$L(t,q_r) = s(t)\int_{-\infty}^{\infty} d\omega \, \frac{\bar{\Delta}(\omega,q_r)}{(\omega - v_F q_r)^2}\{\Theta(\omega t)e^{-i\omega t} + \Theta(q_r t)[it(\omega - v_F q_r) - 1]e^{-iv_F q_r t}\}$$

$$-i\bar{D}(q_r)\Theta(q_r t)|t|e^{-iv_F q_r t} \qquad (B.3)$$

Note that the double pole in $\omega = v_F q_r$ has been eliminated in this representation of L, since the curly bracket has a double zero in $\omega = v_F q_r$. Note also that $L(t, q_r)$ is continuous in $t = 0$, although it doesn't look so at first sight, and

$$L(0^+, r) = L(0^-, r) = \frac{1}{2\pi} \int_{-\infty}^{\infty} dq_r \int_{-\infty}^{\infty} d\omega \, \Theta(-\omega q_r) \frac{|\bar{\Delta}(\omega, q_r)|}{(\omega - v_F q_r)^2} e^{iq_r r} \tag{B.4}$$

which is manifestly real and positive. In particular,

$$L_0 = \frac{1}{2\pi} \int_{-\infty}^{\infty} dq_r \int_{-\infty}^{\infty} d\omega \, \Theta(-\omega q_r) \frac{|\bar{\Delta}(\omega, q_r)|}{(\omega - v_F q_r)^2} \tag{B.5}$$

is a positive real number.

Using the above spectral representation it is easy to show that the momentum distribution function obtained from our solution for G is real, as it should be for the expectation value of a hermitian operator. From $n_{\mathbf{p}} = -i \int_{-\infty}^{\infty} dr \; G(0^-, r) \; e^{-ip_r r}$ it is easy to see that $n_{\mathbf{p}}$ is real if and only if $G^*(0^-, -r) = -G(0^-, r)$, where the asterisk means complex conjugation. For G_0 this is obviously satisfied. Hence, we must require that

$$L^*(0^-, -r) = L(0^-, r) \tag{B.6}$$

This property follows immediately from the expression for $L(0^-, r)$ in terms of $\bar{\Delta}$.

In systems with *short-range* interactions in dimensions $d < 2$ the scaling behavior (7.55) for $\bar{D}$ implies

$$\bar{\Delta}(\omega, q_r) \sim s(\omega) |\omega|^{d-1} \tilde{\Delta}(q_r/\omega) = s(q_r) |q_r|^{d-1} s(\tilde{\omega}) |\tilde{\omega}|^{d-1} \tilde{\Delta}(\tilde{\omega}^{-1}) \tag{B.7a}$$

and

$$\bar{D}(q_r) = |q_r|^{d-1} \tilde{D} \tag{B.7b}$$

where $\tilde{D}$ is a real q_r-independent number. Note that $\tilde{\Delta}(\tilde{\omega}^{-1}) > 0$ by definition. Inserting this in (B.3), and substituting ω by $q_r \tilde{\omega}$, one can carry out the q_r-integration explicitly, and obtains

$$L(t, r) = \frac{1}{2\pi} \int_{-\infty}^{\infty} d\tilde{\omega} \, \frac{|\tilde{\omega}|^{d-1} \tilde{\Delta}(\tilde{\omega}^{-1})}{(\tilde{\omega} - v_F)^2} \left[I_{d,\Lambda}^{s(\tilde{\omega} t)}(r - \tilde{\omega} t) - s(\tilde{\omega}) I_{d,\Lambda}^{s(t)}(r - v_F t) \right.$$

$$\left. + s(\tilde{\omega}) i |t| (\tilde{\omega} - v_F) I_{d+1,\Lambda}^{s(t)}(r - v_F t) \right] - \frac{i\tilde{D}}{2\pi} \, |t| \, I_{d+1,\Lambda}^{s(t)}(r - v_F t) \tag{B.8}$$

where

$$I_{d,\Lambda}^{\alpha}(y) = \int_{-\infty}^{\infty} dq_r \Theta(\alpha q_r) |q_r|^{d-2} e^{iq_r y - |q_r|/\Lambda} = \Gamma(d-1) \frac{e^{i\alpha(d-1)\mathrm{arctg}(y\Lambda)}}{(y^2 + \Lambda^{-2})^{(d-1)/2}} \tag{B.9}$$

Due to the double zero of the curly bracket for $\tilde{\omega} = v_F$ in (B.8), the $\tilde{\omega}$-integral is well-defined and finite. Replacing it by a principal value integral around v_F changes nothing. Exploiting the relation

$$P\int_{-\infty}^{\infty} d\tilde{\omega}\, \frac{s(\tilde{\omega})|\tilde{\omega}|^{d-1}\tilde{\Delta}(\tilde{\omega}^{-1})}{v_F - \tilde{\omega}} + \tilde{D} \;=\; \mathrm{Re}\tilde{D}(v_F^{-1}) \qquad (B.10)$$

the two terms proportional to $|t|$ in (B.8) can be combined to one term proportional to $\mathrm{Re}\tilde{D}(v_F^{-1})$:

$$L(t,r) = \frac{1}{2\pi} P\int_{-\infty}^{\infty} d\tilde{\omega}\, \frac{|\tilde{\omega}|^{d-1}\tilde{\Delta}(\tilde{\omega}^{-1})}{(\tilde{\omega}-v_F)^2}\, [I_{d,\Lambda}^{s(\tilde{\omega}t)}(r-\tilde{\omega}t) - s(\tilde{\omega}) I_{d,\Lambda}^{s(t)}(r-v_F t)]$$
$$-\frac{i\mathrm{Re}\tilde{D}(v_F^{-1})}{2\pi}\, |t|\, I_{d+1,\Lambda}^{s(t)}(r-v_F t) \qquad (B.11)$$

Note that $\tilde{D}(v_F^{-1}) = 0$ in the one-dimensional Luttinger model, i.e. the t-linear term vanishes in that case.

The constant $\tilde{L}_0$ defined in (7.61) can be written as

$$\tilde{L}_0 = \frac{\Gamma(d-1)}{\pi} \int_{-\infty}^{0} d\tilde{\omega}\, |\tilde{\omega}|^{d-1} \frac{\tilde{\Delta}(\tilde{\omega}^{-1})}{(\tilde{\omega}-v_F)^2} \qquad (B.12)$$

Note that the logarithmic "ultraviolet" divergence of $\tilde{L}_0$ for $d \to 2$ does of course *not* imply that $Z \to 0$ for $d \to 2$, since the range of applicability of the leading small-Λ behavior in $d < 2$ shrinks for increasing dimensionality. The divergence of $\tilde{L}_0$ probably signals the expected substitution of leading terms proportional to Λ^{d-1} in $d < 2$ by terms of order $\Lambda \log \Lambda$ in $d = 2$.

Appendix C: Bubbles for a quadratic dispersion relation

Here we list explicit formulae for particle-hole and particle-particle bubbles in d-dimensional systems with the usual *quadratic* (i.e. not linearized) dispersion relation $\epsilon_{\mathbf{k}} = \mathbf{k}^2/2m$. For this $\epsilon_{\mathbf{k}}$ exact analytic expressions can be obtained for the spectral functions associated with the bubbles for arbitrary excitation energies ω.

We introduce dimensionless momentum and energy variables $\bar{\mathrm{q}} = |\mathbf{q}|/k_F$, $\bar{\mathrm{p}} = |\mathbf{p}|/k_F$ and $\bar{\omega} = \omega/v_F k_F = \omega/2\epsilon_F$, where $v_F = k_F/m$ and $\epsilon_F = k_F^2/2m$.

Let us start with the *particle-hole* bubble $\Pi_0(\omega, \mathbf{q})$, as defined in (7.8). One has to distinguish several regimes in the $(\bar{\mathrm{q}}, \bar{\omega})$-plane, which are separated by the functions $\pm\bar{\omega}_+(\bar{\mathrm{q}})$ and $\pm\bar{\omega}_-(\bar{\mathrm{q}})$, where

$$\bar{\omega}_\pm(\bar{\mathrm{q}}) = \pm\bar{\mathrm{q}} + \bar{\mathrm{q}}^2/2 \qquad (C.1)$$

For the spectral function associated with the particle-hole bubble one finds [37]

$$\Delta^{ph}(\omega,\mathbf{q}) = \frac{S_{d-1}}{(2\pi)^d}\frac{1}{d-1}\frac{k_F^{d-1}}{v_F}\frac{1}{\bar{\mathrm{q}}^d}\Big[(\bar{\mathrm{q}}^2-(\bar\omega-\bar{\mathrm{q}}^2/2)^2)^{\frac{d-1}{2}}-(\bar{\mathrm{q}}^2-(\bar\omega+\bar{\mathrm{q}}^2/2)^2)^{\frac{d-1}{2}}\Big] \quad (C.2a)$$

for $0<\bar\omega<-\bar\omega_-(\bar{\mathrm{q}})$ with $\bar{\mathrm{q}}\in[0,2]$,

$$\Delta^{ph}(\omega,\mathbf{q}) = \frac{S_{d-1}}{(2\pi)^d}\frac{1}{d-1}\frac{k_F^{d-1}}{v_F}\frac{1}{\bar{\mathrm{q}}^d}\Big[\bar{\mathrm{q}}^2-(\bar\omega-\bar{\mathrm{q}}^2/2)^2\Big]^{\frac{d-1}{2}} \quad (C.2b)$$

for $|\bar\omega_-(\bar{\mathrm{q}})|<\bar\omega<\bar\omega_+(\bar{\mathrm{q}})$, and

$$\Delta^{ph}(\omega,\mathbf{q}) = 0 \quad (C.2c)$$

for $\bar\omega>\bar\omega_+(\bar{\mathrm{q}})$ with $\bar{\mathrm{q}}\in[0,2]$, or $0<\bar\omega<\bar\omega_-(\bar{\mathrm{q}})$ with $\bar{\mathrm{q}}>2$. The corresponding results for $\bar\omega<0$ can be obtained from $\Delta^{ph}(\omega,\mathbf{q})=-\Delta^{ph}(-\omega,\mathbf{q})$.

We now turn to the *particle-particle* bubble $K_0(\omega,\mathbf{p})$, defined in (7.20). One has to distinguish several regimes in the $(\bar{\mathrm{p}},\bar\omega)$-plane, which are separated by the functions $\bar\omega_+(\bar{\mathrm{p}})$, $\bar\omega_-(\bar{\mathrm{p}})$ and $\bar\omega_0(\bar{\mathrm{q}})$, where

$$\bar\omega_\pm(\bar{\mathrm{q}}) = \pm\bar{\mathrm{q}}+\bar{\mathrm{q}}^2/2 \quad \text{and} \quad \bar\omega_0(\bar{\mathrm{q}}) = (\bar{\mathrm{q}}/2)^2-1 \quad (C.3)$$

For the spectral function associated with the *particle-particle* bubble we have derived the expressions

$$\Delta^{pp}(\omega,\mathbf{p}) = \frac{S_{d-1}}{(2\pi)^d}\frac{k_F^{d-1}}{v_F}\left[\bar\omega-\bar\omega_0(\bar{\mathrm{p}})\right]^{\frac{d-2}{2}}\int_0^{x(\bar\omega,\bar{\mathrm{p}})}(1-x^2)^{\frac{d-3}{2}} \quad \text{where} \quad x(\bar\omega,\bar{\mathrm{p}}) = \frac{\bar\omega/\bar{\mathrm{p}}}{\sqrt{\bar\omega-\bar\omega_0(\bar{\mathrm{p}})}} \quad (C.4a)$$

for $\bar\omega_-(\bar{\mathrm{p}})<\bar\omega<\bar\omega_+(\bar{\mathrm{p}})$,

$$\Delta^{pp}(\omega,\mathbf{p}) = \frac{S_{d-1}}{(2\pi)^d}\frac{k_F^{d-1}}{v_F}\,s(\bar\omega)\left[\bar\omega-\bar\omega_0(\bar{\mathrm{p}})\right]^{\frac{d-2}{2}}\int_0^1(1-x^2)^{\frac{d-3}{2}} = \frac{s(\bar\omega)k_F^{d-1}/v_F}{2^d\pi^{d/2}\Gamma(d/2)}\left[\bar\omega-\bar\omega_0(\bar{\mathrm{p}})\right]^{\frac{d-2}{2}} \quad (C.4b)$$

for $\bar\omega_0(\bar{\mathrm{p}})<\bar\omega<\bar\omega_-(\bar{\mathrm{p}})$ or $\bar\omega>\bar\omega_+(\bar{\mathrm{p}})$, and

$$\Delta^{pp}(\omega,\mathbf{p}) = 0 \quad (C.4c)$$

for $\bar\omega<\bar\omega_0(\bar{\mathrm{p}})$.

The division of energy-momentum space in different regimes does not depend on dimensionality. An illustration of these regimes and explicit expressions for Π_0 and K_0 in $d=1,2,3$ dimensions (including the real parts) can be found in a paper by Fukuyama et al. [30].

Appendix D: Perturbative self-energy in one dimension

In the one-dimensional g-ology model (see Sec. 6) without umklapp processes and couplings $g_{i\perp} =: g_i \neq 0$, $g_{i\parallel} = 0$ for $i = 1, 2, 4$, the Feynman diagram in Fig. 7.2 yields the contribution

$$-\frac{g_1^2 + g_2^2}{16\pi v_F^2}(\xi - \xi_{\mathbf{p}})\Theta(|\xi| - |\xi_{\mathbf{p}}|) - \frac{g_4^2}{8\pi v_F^2} s(\xi_{\mathbf{p}})\xi^2 \delta(\xi - \xi_{\mathbf{p}}) \qquad (D.1)$$

to $\mathrm{Im}\Sigma(\xi, \mathbf{p})$. For $\xi_{\mathbf{p}} = 0$, this yields, via Kramers-Kronig, a contribution

$$\frac{g_1^2 + g_2^2}{8\pi^2 v_F^2}\, \xi \log|\xi/v_F\Lambda| \qquad (D.2)$$

to the real part $\mathrm{Re}\Sigma(\xi, \mathbf{p})$, where Λ is a band-width cutoff. For interactions between opposite spins (only) these are the only second order contributions to the self-energy (except trivial constant Hartree terms).

Appendix E: Limit d → 1 for the propagator

It is instructive to recover the one-dimensional solution for G by taking the limit $d \to 1$ in the general d-dimensional result (B.11). In $d = 1$, the effective interaction of two particles with parallel spins near the same Fermi point has the form

$$D(\omega, q_r) = (\omega - v_F q_r) \sum_{\nu=c,s} \left[\frac{A_\nu}{\omega - u_\nu q_r + i0^+ s(\omega)} + \frac{B_\nu}{\omega + u_\nu q_r + i0^+ s(\omega)} \right] \qquad (E.1)$$

Since there is no transverse component q_t in one dimension, D and $\bar{D}$ are identical. The spectral weight of the dynamical part of D is obviously given by

$$\Delta(\omega, q_r) = q_r \sum_\nu [(u_\nu - v_F)A_\nu \delta(\omega - u_\nu q_r) - (u_\nu + v_F)B_\nu \delta(\omega + u_\nu q_r)] \qquad (E.2)$$

while the ω-independent part is a constant $D(q_r) = \sum_\nu (A_\nu + B_\nu) = g_4^{\sigma\sigma}$. All these functions are scale-invariant. The scaling function $\tilde{\Delta}$ is obtained as

$$\tilde{\Delta}(\tilde{\omega}^{-1}) = \sum_\nu [(u_\nu - v_F)A_\nu \delta(\tilde{\omega} - u_\nu) + (u_\nu + v_F)B_\nu \delta(\tilde{\omega} + u_\nu)] \qquad (E.3)$$

while $\tilde{D}(v_F^{-1})$ is obviously zero. To obtain $L(t, r)$, we must expand the function $I^\alpha_{d,\Lambda}(y)$ around $d = 1$:

$$I^\alpha_{d,\Lambda}(y) = \frac{1}{d-1} - \log(y + i\alpha/\Lambda) + i\alpha\pi/2 + \mathcal{O}(d-1) \qquad (E.4)$$

Inserting all this in our expression (B.11) for $L(t, r)$, performing the $\tilde{\omega}$-integrals (trivial due to δ-functions), and collecting all terms, one gets

$$L(t, r) \;\to\; \frac{\eta}{d-1} \;+\; \log(r - v_F t + is(t)/\Lambda)$$

$$-\sum_{\nu}\Big[(1/2+\eta_\nu/2)\log(r-u_\nu t+is(t)/\Lambda)+(\eta_\nu/2)\log(r+u_\nu t-is(t)/\Lambda)\Big] \qquad (E.5)$$

where $\eta_\nu = B_\nu/\pi(u_\nu+v_F) = 2-A_\nu/\pi(u_\nu-v_F)$ and $\eta=\sum_\nu \eta_\nu$. Finally, L_0 is determined by

$$L_0 = L(0,0) = \frac{\eta}{d-1} + \eta\log\Lambda \qquad (E.6)$$

Thus the constant $\eta/(d-1)$ is cancelled in $L(r,t)-L_0$, and we are left with the well-known result for the Luttinger model in one dimension [20], discussed already in Sec. 6.

References

[1] L.D. Landau, Sov. Phys. JETP **3**, 920 (1956); ibid **5**, 101 (1957).

[2] A particularly thorough discussion of Fermi liquid theory is given by P. Nozières, *Theory of Interacting Fermi Systems* (Benjamin, Amsterdam 1964).

[3] D. Vollhardt and P. Wölfle, *The superfluid phases of Helium 3* (Taylor and Francis, London 1990).

[4] For reviews on heavy fermion systems, see G. Czycholl, Phys. Rep. **143**, 277 (1986); P. Fulde, J. Keller, and G. Zwicknagl, in *Solid State Physics*, Vol. 41, ed. by H. Ehrenreich and D. Turnbull (Academic, San Diego 1988); P.A. Lee, T.M. Rice, J.W. Serene, L.J. Sham, and J.W. Wilkins, Comments Cond. Matt. Phys. **12**, 99 (1986).

[5] See, for example, Proceedings of the International Conference on *Materials and Mechanisms of Superconductivity: High Temperature Superconductors IV*, ed. P. Wyder (North Holland 1994).

[6] For a review on the Fermi surface structure of high-T_c superconductors see, for example, W.E. Pickett, H. Krakauer, R.E. Cohen and D.J. Singh, Science **255**, 46 (1992); Z.X. Shen and D.S. Dessau, Phys. Rep. **253**, 1 (1995).

[7] P. W. Anderson, Phys. Rev. Lett. **64**, 1839 (1990); Phys. Rev. Lett. **65**, 2306 (1990).

[8] C. M. Varma, P. B. Littlewood, S. Schmitt-Rink, E. Abrahams and A. E. Ruckenstein, Phys. Rev. Lett. **63**, 1996 (1989).

[9] B. Andraka and A.M. Tsvelik, Phys. Rev. Lett. **67**, 2886 (1991).

[10] B. Andraka and G.R. Stewart, Phys. Rev. B **47**, 3208 (1993).

[11] C.L. Seaman et al., Phys. Rev. Lett. **67**, 2882 (1991).

[12] H. v. Löhneysen, T. Pietrus, G. Portisch, H.G. Schlager, A. Schröder, M. Sieck, and T. Trappmann, Phys. Rev. Lett. **72**, 3262 (1994).

[13] L. Degiorgi, H.R. Ott, and F. Hulliger, Phys. Rev. B **52**, 42 (1995); L. Degiorgi and H.R. Ott, J. Phys.: Cond. Matt. (in press).

[14] A.A. Abrikosov, L.P. Gorkov, and I.E. Dzyaloshinski, *Methods of Quantum Field Theory in Statistical Physics* (Dover, New York 1975).

[15] J. Feldman and E. Trubowitz, Helv. Phys. Act. **63**, 156 (1990); **64**, 213 (1991); **65**, 679 (1992).

[16] G. Benfatto and G. Gallavotti, Phys. Rev. **B42**, 9967 (1990); J. Stat. Phys. **59**, 541 (1990).

[17] J. Feldman, J. Magnen, V. Rivasseau, and E. Trubowitz in *The State of Matter*, M. Aizenmann and H. Araki eds, Advanced Series in Mathematical Physics Vol. **20** (World Scientific, 1994).

[18] R. Shankar, Physica A **177**, 530 (1991).

[19] R. Shankar, Rev. Mod. Phys. **66**, 129 (1994).

[20] For an extensive early review on 1D and quasi-1D electronic systems, see J. Solyom, Adv. Phys. **28**, 201 (1979).

[21] S. Tomonaga, Prog. Theor. Phys. **5**, 544 (1950); J.M. Luttinger, J. Math. Phys. **4**, 1154 (1963).

[22] D.C. Mattis and E.H. Lieb, J. Math. Phys. **6**, 304 (1965).

[23] For a comprehensive up-to-date review on one-dimensional Fermi systems, see J. Voit, Rep. Prog. Phys. **58**, 977 (1995).

[24] For a recent review on one-dimensional Fermi systems (including a discussion of the one-dimensional Hubbard model), see H.J. Schulz, preprint (1995).

[25] F.D.M. Haldane, J. Phys. C **14**, 2585 (1981); Phys. Rev. Lett. **45**, 1358 (1980); **47**, 1840 (1981); Phys. Lett. **81A**, 153 (1981).

[26] A. Viroztek and J. Ruvalds, Phys. Rev. B **42**, 4064 (1990).

[27] P.C. Pattnaik, C.L. Kane, D.M. Newns, and C.C. Tsuei, Phys. Rev. B **45**, 5714 (1992).

[28] D.M. Newns, C.C. Tsuei, R.P. Huebener, P.J.M. van Bentum, P.C. Pattnaik, and C.C. Chi, Phys. Rev. Lett. **73**, 1695 (1994).

[29] R. Hlubina and T.M. Rice, Phys. Rev. B **51**, 9253 (1995).

[30] H. Fukuyama, Y. Hasegawa, and O. Narikiyo, J. Phys. Soc. Jpn. **60**, 2013 (1991).

[31] J.R. Engelbrecht and M. Randeria, Phys. Rev. B **45**, 12419 (1992).

[32] P.C.E. Stamp, J. Phys. I (France) **3**, 625 (1993).

[33] W. Metzner and C. Castellani, Int. J. Mod. Phys. B (1995).

[34] J. Feldman, H. Knörrer, D. Lehmann, and E. Trubowitz, in *Constructive Physics*, V. Rivasseau (ed.), Springer Lecture Notes in Physics (Springer, 1995).

[35] C. Castellani, C. Di Castro and W. Metzner, Phys. Rev. Lett. **72**, 316 (1994).

[36] P.C.E. Stamp, Phys. Rev. Lett. **68**, 2180 (1992).

[37] P. Bares and X.G. Wen, Phys. Rev. B **48**, 8636 (1993).

[38] C. Castellani and C. Di Castro, Physica C **235-240**, 99 (1994); A. Maccarone, M.Sc. thesis at Università "La Sapienza", Roma (1994).

[39] G. Baskaran and P.W. Anderson, Phys. Rev. B **37**, 580 (1988).

[40] L.B. Ioffe and A.I. Larkin, Phys. Rev. B **39**, 8988 (1989).

[41] P.A. Lee and N. Nagaosa, Phys. Rev. B **46**, 5621 (1992).

[42] T. Holstein, R. Norton, and P. Pincus, Phys. Rev. B **8**, 2649 (1973).

[43] M. Reizer, Phys. Rev. B **39**, 1602 (1989).

[44] B.I. Halperin, P.A. Lee and N. Read, Phys. Rev. B **47**, 7312 (1993).

[45] V. Kalmeyer and S.C. Zhang, Phys. Rev. B **46**, 9889 (1992).

[46] H. von Löhneysen, J. Phys.: Cond. Matt. (in press).

[47] F. Steglich et al., Proceedings of Euroconference on Correlations and Unconventional Quantum Liquids (Evora 1996).

[48] T. Moriya, Y. Takahashi, and K. Ueda, J. Phys. Soc. Jpn. **59**, 2905 (1990).

[49] P. Monthoux and D. Pines, Phys. Rev. B **49**, 4261 (1994) and references therein.

[50] V.J. Emery and S. Kivelson, Physica **209C**, 597 (1993).

[51] C. Castellani, C. Di Castro and M. Grilli, Phys. Rev. Lett. **75**, 4650 (1995).; A. Perali, Castellani, C. Di Castro and M. Grilli, Phys. Rev. B, in press (1996).

[52] C.M. Varma, preprint (1996).

[53] P. Nozières and A. Blandin, J. Phys. **41**, 193 (1980).

[54] D.L. Cox, Phys. Rev. Lett. **59**, 1240 (1987).

[55] N. Andrei and C. Destri, Phys. Rev. Lett. **52**, 364 (1984).

[56] P.B. Wiegmann and A.M. Tsvelick, Z. Phys. B **54**, 201 (1985).

[57] I. Affleck and A.W.W. Ludwig, Nucl. Phys. **B360**, 641 (1991); Phys. Rev. Lett. **67**, 161 (1991); A.W.W. Ludwig and I. Affleck, Phys. Rev. Lett. **67**, 3160 (1991).

[58] Q. Si, G. Kotliar, and A. Georges, Phys. Rev. B **46**, 1261 (1992).

[59] Q. Si and G. Kotliar, Phys. Rev. Lett. **70**, 3143 (1993); Phys. Rev. B **48**, 13881 (1993).

[60] Q. Si, M.J. Rozenberg, G. Kotliar, and A.E. Ruckenstein, Phys. Rev. Lett. **72**, 2761 (1994).

[61] I.E. Dzyaloshinskii and A.I. Larkin, Sov. Phys. JETP **38**, 202 (1974).

[62] P. Kopietz, J. Hermisson, and K. Schönhammer, Phys. Rev. B **52**, 10877 (1995).

[63] L.B. Ioffe, D. Lidsky, and B.L. Altshuler, Phys. Rev. Lett. **73**, 472 (1994); B.L. Altshuler, L.B. Ioffe, and A.J. Millis, Phys. Rev. B **50**, 14048 (1994).

[64] D.V. Khveshchenko and P.C.E. Stamp, Phys. Rev. Lett. **71**, 2118 (1994); Phys. Rev. B **49**, 5227 (1994).

[65] A. Luther and I. Peschel, Phys. Rev. B **9**, 2911 (1974).

[66] D.C. Mattis, J. Math. Phys. **15**, 609 (1974).

[67] A. Luther, Phys. Rev. B **19**, 320 (1979).

[68] F.D.M. Haldane, Helv. Phys. Acta, **65**, 152 (1992); Proceedings of the International School of Physics "Enrico Fermi", Course CXXI, eds. R.A. Broglia and J.R. Schrieffer (North Holland, Amsterdam, 1994).

[69] A. Houghton and J.B. Marston, Phys. Rev. B **48**, 7790 (1993).

[70] A. Houghton, H.-J. Kwon, and J.B. Marston, Phys. Rev. B **50**, 1351 (1994); H.-J. Kwon, A. Houghton, and J.B. Marston, Phys. Rev. B **52**, 8002 (1995).

[71] A.H. Castro Neto and E. Fradkin, Phys. Rev. Lett. **72**, 1393 (1994); Phys. Rev. B **49**, 10877 (1994).

[72] D.V. Khveshchenko, Phys. Rev. B **49**, 16893 (1994); Phys. Rev. B **52**, 4833 (1995).

[73] P. Kopietz and K. Schönhammer, Z. Phys. B **100**, 561 (1996).

[74] K. Ueda and T.M. Rice, Phys. Rev. B **29**, 1514 (1984).

[75] K.G. Wilson, Rev. Mod. Phys. **47**, 773 (1975); K.G. Wilson and J.B. Kogut, Phys. Rep. **12**, 7 (1974), and references therein.

[76] For a review, see P.A. Lee and T.V. Ramakrishnan, Rev. Mod. Phys. **57**, 287 (1985); C. Castellani, C. Di Castro and G. Strinati in *Lecture Notes in Physics* **268** (Springer 1986); A.M. Finkelstein, Sov. Sci. Rev. A. Phys. **14**, 1 (1990); D. Belitz and T.R. Kirkpatrick, Rev. Mod. Phys. **66**, 261 (1994).

[77] See, in particular, the review articles by J.W. Serene and D. Rainer, Phys. Rep. **101**, 221 (1983); D. Rainer, Prog. Low Temp. Phys. **X**, 371 (1986).

[78] G. Benfatto, G. Gallavotti, and V. Mastropietro, Phys. Rev. B **42**, 9967 (1990); G. Benfatto, G. Gallavotti, A. Procacci, and B. Scoppola, Comm. Math. Phys. **160**, 93 (1994).

[79] G. Benfatto and G. Gallavotti, *Renormalization Group* (Princeton University Press 1995).

[80] J. Feldman, H. Knörrer, D. Lehmann and E. Trubowitz in *Constructive Physics*, ed. V. Rivasseau, Springer Lecture Notes in Physics (Springer 1995).

[81] T. Chen, J. Fröhlich, and M. Seifert, in Proceedings of the Les Houches Summer School 1994.

[82] J. Polchinski in Proceedings of 1992 Theoretical Advanced Studies Institute in Elementary Particle Physics, eds. J. Harvey and J. Polchinski (World Scientific, Singapore, 1993).

[83] J.W. Negele and H. Orland, *Quantum Many-Particle Systems* (Addison-Wesley 1988).

[84] V.N. Popov, *Functional Integrals and Collective Excitations* (Cambridge University Press).

[85] J. Feldman, M. Salmhofer and E. Trubowitz, preprint (1995).

[86] W. Metzner and C. Di Castro, Phys. Rev. B **47**, 16107 (1993).

[87] W. Kohn and J.M. Luttinger, Phys. Rev. Lett. **15**, 524 (1965).

[88] E. Trubowitz, private communication.

[89] D. Fay and A. Layzer, Phys. Rev. Lett. **20**, 187 (1968).

[90] S. Küchenhoff and P. Wölfle, Phys. Rev. B **38**, 935 (1988).

[91] M.Yu. Kagan and A.V. Chubukov, JETP Lett. **47**, 614 (1988); M.A. Baranov, A.V. Chubukov and M.Yu. Kagan, Int. J. Mod. Phys. B **6**, 2471 (1992).

[92] T. Bohr, Nordita preprint 81/4, *Lectures on the Luttinger Model* (1981).

[93] G.Y. Chitov and D. Sénéchal, Phys. Rev. B **52**, 13487 (1995).

[94] G. Baym and C. Pethick, *Landau Fermi Liquid Theory* (Wiley, New York 1991).

[95] M. Eschrig, J. Heym, and D. Rainer, J. Low Temp. Phys. **95**, 323 (1994).

[96] See, for example, J.M. Ziman, *Electrons and Phonons* (Oxford University Press 1960).

[97] I.Ja. Pomerantchuk, Sov. Phys. JETP **8**, 361 (1958).

[98] L.N. Cooper, Phys. Rev. **104**, 1189 (1956).

[99] W. Metzner, unpublished notes (1995).

[100] Y.B. Kim, A. Furusaki, X.G. Wen, and P.A. Lee, Phys. Rev. B **50**, 17917 (1994).

[101] J.A. Hertz and M.A. Klenin, Phys. Rev. B **10**, 1084 (1974).

[102] J. Feldman, H. Knörrer, R. Sinclair and E. Trubowitz, preprint (ETH Zürich, 1996).

[103] J. Fröhlich, R. Götschmann and P.A. Marchetti, J. Phys. A **28**, 1169 (1995).

[104] H.C. Fogedby, J. Phys. C **9**, 3757 (1976).

[105] D.K.K. Lee and Y. Chen, J. Phys. A **21**, 4155 (1988).

[106] P. Kopietz, Habilitationsschrift, Georg-August University Göttingen (1995).

[107] E.H. Lieb and F.Y. Wu, Phys. Rev. Lett. **20**, 1445 (1968).

[108] H. Frahm and V.E. Korepin, Phys. Rev. B **42**, 10553 (1990); Phys. Rev. B **43**, 5653 (1991).

[109] F.D.M. Haldane and Y. Tu (unpublished).

[110] N. Kawakami and S.K. Yang, Phys. Lett. A **148**, 359 (1990).

[111] H.J. Schulz, Phys. Rev. Lett. **64**, 2831 (1990).

[112] M. Ogata and H. Shiba, Phys. Rev. B **41**, 2326 (1990).

[113] A. Parola and S. Sorella, Phys. Rev. Lett. **64**, 1831 (1990); S. Sorella, A. Parola, M. Parrinello and E. Tosatti, Europhys. Lett. **12**(8), 721 (1990).

[114] K. Penc, F. Mila and H. Shiba, Phys. Rev. Lett. **75**, 894 (1995).

[115] See, in particular, J.M.P. Carmelo and P. Horsch, Phys. Rev. Lett. **68**, 871 (1992).

[116] H.U. Everts and H. Schulz, Sol. State Comm. **15**, 1413 (1974).

[117] V.J. Emery in *Highly Conducting One-Dimensional Solids*, edited by J.T. Devreese, R.P. Evrard, and V.E. van Doren (Plenum, New York 1979).

[118] For a review, see C. Di Castro and G. Jona-Lasinio in *Phase Transitions and Critical Phenomena*, edited by C. Domb and M.S. Green (Academic, London, 1976), Vol.6.

[119] For an introduction to the field theoretic renormalization group, see D. Amit, *Field Theory, the Renormalization Group and Critical Phenomena* (World Scientific, Singapore, 1984).

[120] C. Di Castro and W. Metzner, Phys. Rev. Lett. **67**, 3852 (1991).

[121] A. Luther and V.J. Emery, Phys. Rev. Lett. **33**, 589 (1974).

[122] K. Johnson, Nuovo Cimento **20**, 773 (1961).

[123] V. Meden and K. Schönhammer, Phys. Rev. B **46** (1992).

[124] J. Voit, Phys. Rev. B **47**, 6740 (1993).

[125] N.D. Mermin, Phys. Rev. **159**, 161 (1967).

[126] H. Fukuyama and M. Ogata, J. Phys. Soc. Japan. **63**, 3923 (1995); H. Fukuyama and M. Ogata in *Spectroscopy of Mott Insulator and Correlated Metals*, ed. A. Fujimori and Y. Tokura (Springer Verlag 1995), p. 34.

[127] C. Hodges, H. Smith, and J.W. Wilkins, Phys. Rev. B **4**, 302 (1971).

[128] P. Bloom, Phys. Rev. B **12**, 125 (1975).

[129] A. Houghton, H.-J. Kwon, J.B. Marston, and R. Shankar, J. Phys. C **6**, 4909 (1994).

[130] K. Jain, Phys. Rev. Lett. **63**, 199 (1989); Phys. Rev. B **41**, 7653 (1990); Adv. Phys. **41**, 105 (1992).

[131] A. Lopez and E. Fradkin, Phys. Rev. B **44**, 5246 (1991); Phys. Rev. Lett. **69**, 2126 (1992).

[132] F. Wilczek, *Fractional Statistics and Anyon Superconductivity* (World Scientific, Singapore, 1990).

[133] J. Gan and E. Wong, Phys. Rev. Lett. **71**, 4226 (1993).

[134] J. Fröhlich, R. Götschmann, and P.A. Marchetti, Commun. Math. Phys. **173**, 417 (1995).

[135] J. Polchinski, Nucl. Phys. B **422**, 617 (1994).

[136] C. Nayak and F. Wilczek, Nucl. Phys. B **417**, 359 (1994); **430**, 534 (1994).

[137] A.G. Aronov and P. Wölfle, Phys. Rev. Lett. **72**, 2239 (1994); Phys. Rev. B **50**, 16574 (1994).

[138] P.A. Lee, Phys. Rev. Lett. **63**, 680 (1989).

[139] B. Blok and H. Monien, Phys. Rev. B **47**, 3454 (1993).

[140] H.-J. Kwon, A. Houghton, and J.B. Marston, Phys. Rev. Lett. **73**, 284 (1994); preprint (1995).

[141] P. Kopietz, Phys. Rev. B **53**, 12761 (1996).

[142] S. Chakravarty, R.E. Norton, and O.F. Syljuåsen, Phys. Rev. Lett. **74**, 1423 (1995).

Figure Captions

Fig. 2.1: Diagrammatic representation of the bare interaction $g_{\mathbf{kk'};\mathbf{q}}$ with momentum labels according to the notations used in this paper.

Fig. 2.2: Mode elimination in momentum space. Fermi fields ψ_k, ψ_k^* with $d(\mathbf{k}, \partial\mathcal{F}) > \Lambda$ are integrated out. The remaining effective theory is defined on a shell of width 2Λ around the Fermi surface in momentum space.

Fig. 2.3: Examples for Feynman diagrams contributing to the self-energy. Shaded boxes connected to $2n$ lines represent renormalized effective n-particle interactions.

Fig. 2.4: Contribution to the effective three-particle interaction (in terms of bare interactions); the internal line must carry a momentum outside the Λ-shell.

Fig. 2.5: Scattering processes on the Fermi surface in two dimensions.

Fig. 2.6: Diagrams contributing to effective two-particle interactions in second order perturbation theory. The shaded squares represent (antisymmetrized) effective two-particle interactions.

Fig. 4.1: The two-particle vertex Γ in terms of the particle-hole irreducible vertex Γ_{irr}. The internal lines correspond to dressed propagators G.

Fig. 5.1: Dyson equation relating the vertex part Γ to irreducible components.

Fig. 5.2: Fermionic loop with N insertions.

Fig. 5.3: Diagrams contributing to the RPA effective interaction D.

Fig. 5.4: The current-current response function $J^{\mu\nu}$ in terms of D.

Fig. 5.5: Perturbation expansion of the current vertex $\Lambda^\mu(p;q)$.

Fig. 5.6. Two cancelling contributions to $\Lambda^\mu(p;q)$.

Fig. 5.7: Dyson equation relating the self-energy Σ to the irreducible density vertex Λ^0, the effective interaction D and the propagator G.

Fig. 6.1: Band structure of the g-ology model.

Fig. 6.2: Interaction terms in the g-ology model; "-" and "+" indicate the left and right Fermi point, respectively.

Fig. 7.1: Particle-hole bubble $\Pi_0(q)$ for small q as a function of $\tilde{\omega} = \omega/|\mathbf{q}|$ in dimensions $d = 1.1$, 1.5, 2 and 3.

Fig. 7.2: Feynman diagram contributing to the second order self-energy.

Fig. 7.3: $\mathrm{Im}\Sigma(\xi, \mathbf{p})$ from second order perturbation theory as a function of ξ for fixed $p_r = 0.1k_F$ in dimensions $d = 1$, 1.5, 2, and 3 ($g = k_F = v_F = 1$).

Fig. 7.4: Spectral function $\rho(\xi, \mathbf{p})$ from second order perturbation theory as a function of ξ for fixed $p_r = 0.1k_F$ in $d = 1.5$ ($g = 2$, $k_F = v_F = 1$).

Fig. 7.5: Effective interaction $D^{\sigma\sigma}(\omega, \mathbf{q})$ as a function of $\tilde{\omega} = \omega/|\mathbf{q}|$ for a constant coupling $g = 2$ between opposite spins in dimensions $d = 1.1$, 1.5 and 2.

Fig. 7.6: Feynman diagram representing the RPA self-energy.

Fig. 7.7: RPA-result for $\mathrm{Im}\Sigma(\xi, \mathbf{p})$ as a function of ξ at fixed $p_r = 0.1k_F$ for a constant coupling $g = 2$ between opposite spins in $d = 1.5$ dimensions ($k_F = v_F = 1$).

Fig. 9.1: The gauge-field propagator D in random phase approximation: The polarization insertion Π is approximated by its 1-loop result Π_0.

Fig. 9.2: The fermion self-energy within RPA, where D is the RPA gauge-field propagator, and $\lambda^\mu(k) = (1, \mathbf{v_k})$ the bare fermion-gauge-field vertex.

Fig. 9.3: Dyson equation relating the fermion self-energy Σ to the fermion propagator G, the gauge-field propagator $D^{\mu\nu}$ and the irreducible fermion-gauge-field vertex Λ^μ.

Fig. 10.1: Phase-diagram of Fermi systems with short-range interactions in the dimensionality-coupling plane, where g is a typical renormalized coupling in the forward scattering channel.

Fig. A.1: Two loops with $N = 3$ insertions.


d
3
2
1
0
conventional
Fermi liquid
$\Gamma_k \propto k_r^2$
unconventional
Fermi liquid
$\Gamma_k \propto k_r^d$
strong
coupling
(Pomerantchuk)
instabilities
Luttinger liquid
g_c
g

Fig. 10.1



Fig. 2.1



Fig. 4.1


$\Gamma^{\mu}_{\sigma}(p;q) = \Lambda'^{\mu}_{\sigma}(p;q) +$
$J^{\mu \mathbf{k'}\sigma'}(q)$
$g_{\mathbf{k'k''}}{}^{\sigma'\sigma''}(q)$
$\Lambda^{\mathbf{k''}\sigma''}_{\sigma}(p;q)$

Fig. 5.1



Fig. 6.1



Fig. 7.1



Fig. 9.1



Fig. A.1



Fig. 2.2



Fig. 5.2



Fig. 6.2



Fig. 7.2



Fig. 9.2

a)



b)



c)



Fig. 2.3



Fig. 5.3



Fig. 7.3



Fig. 9.3



Fig. 2.4



Fig. 5.4


d = 1.5
ρ(ξ,p)
energy ξ
0
20
40
60
80
100
120
0
0.05
0.1
0.15
0.2



$g_F(\theta)$ $g_E(\theta)$ $g_C(\phi)$

Fig. 2.5



Fig. 5.5







Fig. 7.5



Fig. 2.6



Fig. 5.6



Fig. 7.6



Fig. 5.7


d = 1.5
- Im Σ(ξ,p)
energy ξ
0.01
0.008
0.006
0.004
0.002
0
-0.2
-0.15
-0.1
-0.05
0
0.05
0.1
0.15
0.2